



Structure of Everninomicin-2

By A. K. GANGULY,* S. SZMULEWICZ, O. Z. SARRE, and V. M. GIRIJAVALLABHAN
(*Chemical Research Department, Schering Corporation, Bloomfield, New Jersey 07003*)

Summary The structure of everninomicin-2 (**2**) has been elucidated and a method of conversion of everninomicin D (**1**) into (**2**) is discussed.

(6) and evertetetrose (8). Based on the fact that the monomethyl ether of everninomicin-2 on solvolysis yields (6), (7), and (8) and that the ^{13}C n.m.r. spectrum of everninomicin-2 (2) shows the presence of two orthoester carbon atoms, we propose structure (2) for everninomicin-2 and structure (3) for its monomethyl ether.

At this point the conversion of everninomicin D (**1**) into everninomicin-2 (**2**) was considered. This involved hydrolysis of only one of the several glycosidic bonds present in the molecule and more importantly the labile orthoester linkages had to be kept intact. It was conceived that nitroso-everninomicin D (**4**) on treatment with triethyl phosphite or triphenylphosphine could be converted into a nitrene (see formula) which would rearrange with bond migration (one of the three possibilities shown) to an enamine which should on principle hydrolyse the required glycoside bond yielding everninomicin-2 (**2**).



Everninomicin D (**1**) on reduction⁷ with aluminium amalgam in aqueous ethanol yielded hydroxylamino-everninomicin D (**5**) as a colourless crystalline solid, $C_{66}H_{101}Cl_2NO_{34}$, m.p. 185–186 °C, $[\theta]_{255} (-17,400)$, ν_{\max} 2.9 and 5.72 μm , no nitro-group absorption. It gave a positive colour reaction⁸ with triphenyltetrazolium chloride for a hydroxylamino-group. Compound (**5**) was unstable to acid and was oxidised readily in air to nitrosoeverninomicin D.⁴ For preparative purposes hydroxylaminoeverninomicin D (**5**) was oxidized in tetrahydrofuran solution using sodium hypobromite to nitrosoeverninocin D (**4**), a blue amorphous solid, $[\theta]_{255} (-15,000)$. Nitrosoeverninomicin D (**4**) was refluxed in benzene solution with 1.1 mol. equiv. of triphenylphosphine until the blue colour disappeared (*ca.* 15 min). The reaction mixture was evaporated to dryness and the residue chromatographed on silica gel to yield everninomicin-2 (**2**) (identical on comparison with a

† Solvolysis in this communication refers to treatment of the compound with methanolic toluene-*p* sulphonic acid.

sample obtained from a natural source). The overall yield of (2) from (1) was *ca.* 30%. oxylaminoeverninomicin D (5) gave the highest blood level when administered intramuscularly to dogs.⁹

Compounds (1), (2), (4), and (5) possessed equal *in vitro* activity against gram-positive bacteria. However, hydr-

(Received, 6th April 1976; Com. 370.)

¹ M. J. Weinstein, G. M. Luedemann, E. M. Oden, and G. H. Wagman, *Antimicrob. Agents Chemotherapy*, 1964, 24.

² A. K. Ganguly and A. K. Saksena, *J. Antibiotics*, 1975, 28, 707.

³ A. K. Ganguly and S. Szmulewicz, *J. Antibiotics*, 1975, 28, 710.

⁴ A. K. Ganguly, O. Z. Sarre, D. Greeves, and J. Morton, *J. Amer. Chem. Soc.*, 1975, 97, 1982.

⁵ A. K. Ganguly, O. Z. Sarre, and S. Szmulewicz, *Chem. Comm.*, 1971, 746.

⁶ W. D. Ollis and C. Smith, *J.C.S. Chem. Comm.*, 1974, 882. After we completed our work, the paper on the structural elucidation of flambolactone appeared. As the structure of flambolactone was elucidated following similar procedures outlined by us (refs. 4 and A. K. Ganguly, O. Z. Sarre, D. Greeves, and J. Morton, *J. Amer. Chem. Soc.*, 1973, 95, 942), for a related compound and the constants for compound (5) and the mono-*O*-methyl flambolactone were so similar, a direct comparison of the two samples for establishing their identity was felt unnecessary.

⁷ A. K. Ganguly and O. Z. Sarre, U.S.P. 3,915,956.

⁸ G. A. Snow, *J. Chem. Soc.*, 1954, 2589.

⁹ Unpublished work, G. Miller and J. A. Waitz, Schering Corporation.