

Purines. XLIII.¹⁾ A Total Synthesis of the Marine Sponge Base 6-Imino-1,9-dimethyl-8-oxopurine

Tozo FUJII,* Tohru SAITO, and Shigeji MORI

Faculty of Pharmaceutical Sciences, Kanazawa University, Takara-machi, Kanazawa 920, Japan. Received February 19, 1990

The first total synthesis of the 8-oxopurine **8**, a constituent of the marine sponge *Hymeniacidon sanguinea* GRANT, has been achieved via two alternative routes starting from 8-bromo-9-methyladenine (**11**), which is obtainable from 9-methyladenine (**10**) by bromination. The first route includes methylation of **11** with MeI to give the 1-methylated product **13**·HI, conversion of **13**·HI into the free base **13**, and treatment of **13** with NaOAc in boiling AcOH to produce **8** in 25% overall yield (from **10**). The second route includes treatment of **11** with boiling 1N aqueous NaOH and methylation of the resulting 8-oxo derivative **12** with MeI, affording **8** in 63% overall yield (from **10**). The rearranged isomer **15** and the *N*⁶-acetyl derivative **9** have also been synthesized.



Keywords marine sponge 8-oxopurine; adenine bromination; 8-bromoadenine methylation; nucleophilic aromatic substitution; 8-oxoadenine; Dimroth rearrangement; acetylation

More than a dozen naturally occurring unusual purine derivatives, based mainly on the adenine nucleus, have so far been reported from various marine sponges. These sponge purines include 1-methyladenine (spongopurine) (**1**)²⁾ and 7,9-dimethylguaninyl betaine (herbipoline) (**2**)³⁾ from the siliceous giant sponge *Geodia gigas*, 2-methoxyadenosine (spongosine) (**3**) from the Caribbean sponge *Cryptotethia crypta*,⁴⁾ 1-methylisoguanosine (**4**) from the Tasman Sea sponge *Tedania digitata*,⁵⁾ agelasine (**5a**) from the Barbadian sponge *Agelas dispar*,⁶⁾ agelasines A—F (**5b—g**) from the Okinawan sponge *Agelas nakamurai*,^{7,8)} ageline A (agelasine F⁷⁾) (**5g**) and ageline B (**5h**) from a Pacific sponge *Agelas* sp.,⁹⁾ and agelasimine A (**6**) and agelasimine B (**7**) from the orange sponge *Agelas mauritiana*.^{10,11)} Many of these purines exhibit a wide range of interesting biological activities such as maturation-inducing hormonal activity on oocytes of the starfish *Asterias amurensis*, *Asterias rubens*, *Marthasterias glacialis*, or *Asterina pectinifera* (by **1**)^{12,13)}; muscle relaxant,

antiinflammatory, and other pharmacological activities (by **4**)^{5,14)}; antimicrobial activity (by **5b—e**, **5g**, and **5h**)^{7,8)}; cytotoxicity (by **6** and **7**)¹⁰⁾; ichthyotoxicity (by **5g** and **5h**)⁹⁾; inhibitory effect on Na⁺,K⁺-ATPase (by **5b—g**)^{7,8)}; and inhibition of adenosine transfer into rabbit erythrocytes, Ca²⁺-channel antagonistic action, and α₁ adrenergic blockade (by **6** and **7**).¹⁰⁾

In 1985, Cimino *et al.*¹⁵⁾ reported the isolation of a new purine (**8**) and known 1-methyladenine (**1**), although both only in the form of the acetyl derivative (**9** and acetyl-spongopurine), from the English Channel sponge *Hymeniacidon sanguinea* GRANT. While the new acetyl derivative **9** was fully characterized by means of spectroscopic and X-ray crystallographic analyses, the parent base **8** remained unknown because of the difficulty in separating **8** and **1** from each other at the free base level.¹⁵⁾ This was why we tried to secure the base **8** itself by synthesis in the present study. A brief account of the results recorded here has been published in a preliminary form.¹⁶⁾

The synthesis of the target compound **8** started with bromination of 9-methyladenine (**10**),¹⁷⁾ as shown in Chart 1. Treatment of **10** with Br₂ in 0.5M acetate buffer (pH 4) in a manner similar to that reported by Ikehara's group¹⁸⁾






Chart 1

afforded the 8-bromo derivative **11** in 87% yield. It is well known that an alkyl group at the 9-position of adenine itself, as in the case of the methyl group in **10**, orients further alkylation to the 1-position to form 1,9-dialkyladenine (type **16**).¹⁹⁾ If such directivity in alkylation holds in the case of the 8-bromo derivative **11**, the desired 1,9-dimethyl derivative **13** would be accessible from **11** by methylation. In practice, methylation of **11** with MeI in AcNMe₂ at 50 °C for 3.5 h produced **13**·HI in 99% yield. On treatment with aqueous Na₂CO₃, the salt **13**·HI was converted into the free base **13** in 80% yield. The structural assignment of **13** was derived from its ultraviolet (UV) spectra in 95% (v/v) aqueous EtOH and in H₂O at various pH's, which were similar to those^{19h)} of 1,9-dialkyladenines (type **16**), and from the Dimroth rearrangement of **13**·HI to the N⁶,9-dimethyl isomer **14**.²⁰⁾ The rearrangement of **13**·HI was effected in 1 N aqueous NaOH at 55 °C for 35 min, giving **14** in 88% yield. The proton nuclear magnetic resonance (¹H-NMR) spectrum of **14** in dimethyl sulfoxide-*d*₆ exhibited signals at δ 2.96 [3H, dull d, $J=4$ Hz, C(6)-NHMe], 3.66 [3H, s, N(9)-Me], 7.83 [1H, br, C(6)-NHMe], and 8.21 [1H, s, C(2)-H]. The observation of the presence of a vicinal interproton coupling with $J=4$ Hz in the C(6)-NHMe system supported the correctness of the 6-methylamino structure of **14**. Under more vigorous conditions (1 N aqueous NaOH, reflux, 1.5 h), the 8-bromo derivative **14** was further converted into the 8-oxo derivative **15** in 72% yield. The 8-oxo form of **15** was inferred from its infrared (IR) spectrum in a Nujol mull, which showed two NH absorption bands at 3370 and 3190 cm⁻¹ and a carbonyl absorption band at 1695 cm⁻¹.²¹⁾

On the other hand, treatment of **13** with NaOAc in boiling AcOH for 5 h furnished the target compound **8** (mp > 300 °C) and the N⁶-acetyl derivative **9** in 36% and 34% yields, respectively. The UV, IR, and ¹H-NMR spectral data obtained from the synthetic **9** were in agreement with those reported¹⁵⁾ for a "natural" sample. The correctness of the 1,9-dimethyl structure of **8** was confirmed by acetylation (Ac₂O/pyridine, reflux, 10 min) leading to **9** (81% yield) and by the Dimroth rearrangement (1 N aqueous

NaOH, reflux, 1 h) to form **15** (90% yield), which was identical with a sample derived from **14** (*vide supra*). However, attempts to obtain **8** from **9** by selective hydrolysis were all unsuccessful. As in the case of **9**, the existence of **8** in the 8-oxo form was indicated by the appearance of a carbonyl absorption band at 1694 cm⁻¹ in the IR spectrum taken in a Nujol mull.²¹⁾ The transformation of the 8-bromo function into the 8-oxo function achieved in the above step **13**→**8**+**9** has precedents in that Ikehara's group²²⁾ prepared 8-oxoadenosine derivatives from 8-bromoadenosine derivatives under similar reaction conditions. In this transformation, a large portion of the substrate **13** must be protonated in AcOH solution,²³⁾ giving **13**·H⁺ as shown in Chart 2. The nucleophilic aromatic substitution at C(8) of **13**·H⁺ would then be facilitated by the positive charge in the ring system. The resulting 8-acetoxy derivative **17**·H⁺ may be regarded as an activated acetic ester and would produce acetic anhydride and the 8-oxo derivative **8** on attack by acetate ion, which is present in large quantity. Part of both products thus generated would then react to effect acetylation at the N⁶ atom of **8**, and this would account for the formation of the N⁶-acetyl derivative **9** as a by-product. In a similar conversion of an 8-bromoadenosine derivative into the corresponding 8-oxoadenosine derivative, Ikehara and Kaneko^{22c)} observed partial acetylation of the amino and hydroxy groups. Their observation may also be explained in terms of a sequence of processes analogous to that depicted in Chart 2.

The usual regioselectivity found in the above methylation of the 8-bromo derivative **11** led us to expect the same in methylation of the 8-oxo derivative **12**. In order to check this possibility, **12** was prepared in 97% yield from **11** by treatment with boiling 1 N aqueous NaOH for 1.5 h. The assignment of the 8-oxo structure to **12** was based on its IR spectrum in the solid state, which displayed a carbonyl absorption band at 1712 cm⁻¹.²¹⁾ On methylation with MeI in AcNMe₂ at 50 °C for 7 h, **12** gave, after basification, the desired compound **8** in 75% yield. Thus, this second three-step approach to **8** from **10** through **11** and **12** was




Chart 2

found to be more straightforward, simpler to operate, and more efficient (63% overall yield from **10**) than the first four-step approach **10**→**11**→**13**·HI→**13**→**8** (25% overall yield from **10**).

In conclusion, the present results have made it possible to characterize fully the title compound **8** itself, in advance of the yet unrealized isolation of this substance from natural sources. Interestingly, the known regioselectivity in methylation of 9-methyladenine (**10**) also holds in the cases of the 8-bromo and 8-oxo derivatives (**11** and **12**), and this has permitted a concise total synthesis of **8** for the first time. It is also interesting to note that the Dimroth rearrangement of the 8-oxopurine **8** to **15** seems to require more vigorous reaction conditions than that of the 8-bromopurine **13**·HI to **14**. In view of the broad spectrum of biological activities exhibited by the unusual purine derivatives from various marine sponges (*vide supra*), we are now in the process of testing the synthetic 8-oxopurine **8** and related compounds for biological and pharmacological activities.

Experimental

General Notes All melting points were taken on a Yamato MP-1 capillary melting point apparatus and are corrected. Thin-layer chromatography (TLC) was performed on Merck silica gel 60 F₂₅₄ plates (0.25-mm thickness), Merck aluminum oxide F₂₅₄ (type E) plates (0.25 mm), or Funakoshi Avicel SF-2020F plates, and spots were located under UV light (254 nm). Spectra reported herein were recorded on a Hitachi model 320 UV spectrophotometer [on solutions in MeOH, 95% (v/v) aqueous EtOH, 0.1 N aqueous HCl (pH 1), 0.005 M phosphate buffer (pH 7), and 0.1 N aqueous NaOH (pH 13)], a JASCO A-202 IR spectrophotometer, a Hitachi M-80 mass spectrometer, or a JEOL JNM-FX-100 NMR spectrometer at 25 °C with Me₄Si as an internal standard. Elemental analyses were performed by Mr. Y. Itatani and his associates at Kanazawa University. The following abbreviations are used: br=broad, d=doublet, DMSO=dimethyl sulfoxide, m=multiplet, q=quartet, s=singlet, sh=shoulder.

8-Bromo-9-methyladenine (11) 3-Methyladenine (**10**)^{17a,d} (18.64 g, 0.125 mol) was dissolved in 0.5 M NaOAc–AcOH buffer (pH 4) (625 ml) by application of heat. The resulting solution was cooled to room temperature, and a solution of Br₂ (39.95 g, 0.25 mol) in H₂O (2.44 l) was added dropwise with stirring over a period of 100 min. After the mixture had been stirred at room temperature for 5 h, the reaction was quenched by adding a solution of NaHSO₃ (18.8 g, 0.18 mol) in H₂O (400 ml). The resulting mixture was brought to pH 5 with 10% aqueous Na₂CO₃ and then kept in a refrigerator for 2 d. The precipitate that resulted was filtered off, washed with H₂O, and dried to give **11** (24.68 g, 87%) as a pale yellowish solid, mp 265–267 °C (dec.). Recrystallization from EtOH yielded an analytical sample as faintly yellowish prisms, mp 274–275 °C (dec.) [lit.^{18b}] mp 229 °C (dec.) for **11**·1/4H₂O; MS *m/z*: 229, 227 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 266.5 nm (ϵ 15000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 265 (16900); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7)

267 (16000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 267 (16000); IR $\nu_{\text{max}}^{\text{Nujol}}$ cm⁻¹: 3350, 3175 (NH₂); NMR (DMSO-*d*₆) δ : 3.65 [3H, s, N(9)-Me], 7.33 (2H, br, NH₂), 8.12 [1H, s, C(2)-H]. Anal. Calcd for C₆H₆BrN₅: C, 31.60; H, 2.65; N, 30.71. Found: C, 31.49; H, 2.58; N, 30.53.

9-Methyl-8-oxoadenine (12) A stirred mixture of **11** (5.02 g, 22 mmol) and 1 N aqueous NaOH (220 ml) was heated under reflux for 1.5 h. After cooling, the reaction mixture was brought to pH 7 by addition of 10% aqueous HCl and then cooled in an ice bath for 1.5 h. The pale brownish crystals that deposited were filtered off, washed with cold H₂O, and dried to give **12** (3.52 g, 97%), mp > 300 °C. Recrystallization from H₂O afforded an analytical sample as colorless prisms, mp > 300 °C; MS *m/z*: 165 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 271 nm (ϵ 13000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 270 (sh) (10300), 280 (10700); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 271 (13000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 280 (14900); IR $\nu_{\text{max}}^{\text{Nujol}}$ cm⁻¹: 3415, 3250, 3135 (NH₂ and NH), 1712 (CO); NMR (DMSO-*d*₆) δ : 3.22 [3H, s, N(9)-Me], 6.35 (2H, dull, NH₂), 8.01 [1H, s, C(2)-H], 10.09 [1H, dull, N(7)-H]. Anal. Calcd for C₆H₇N₅O: C, 43.64; H, 4.27; N, 42.40. Found: C, 43.65; H, 4.24; N, 42.47.

6-Imino-1,9-dimethyl-8-oxopurine (8) A mixture of **12** (1.98 g, 12 mmol) and MeI (8.52 g, 60 mmol) in AcNMe₂ (20 ml) was stirred at 50 °C for 7 h. The reaction mixture was cooled in an ice bath for 30 min, and the precipitate that resulted was filtered off, washed successively with a little EtOH and ether, and dried to furnish a colorless solid (3.54 g), mp 281–283 °C (dec.), presumed to be the hydriodide **8**·HI. The solid was dissolved in warm H₂O (100 ml), and the resulting aqueous solution was made alkaline (pH 9) with 10% aqueous Na₂CO₃ and kept in a refrigerator overnight. The colorless needles that deposited were filtered off, washed with a little H₂O, and dried over P₂O₅ at 2 mmHg and 75 °C for 6 h to provide the free base **8** (1.62 g, 75%), mp > 300 °C. This sample was identical (by comparison of the IR spectrum and TLC mobility) with the anhydrous one obtained from **13** by nucleophilic aromatic substitution of the 8-bromo group (*vide infra*).

6-Acetyl-8-oxopurine (9) A stirred mixture of **8** (896 mg, 5 mmol) and Ac₂O (610 mg, 6 mmol) in pyridine (15 ml) was heated under reflux for 10 min. After cooling, the reaction mixture was kept in an ice bath for 1 h, and the precipitate that resulted was filtered off, washed successively with a little AcOEt and EtOH, and dried to yield **9** (899 mg, 81%) as a faintly yellowish solid, mp 247.5–249.5 °C (dec.). Recrystallization from AcOEt gave an analytical sample as colorless needles, mp 248–250 °C (dec.) (lit.¹⁵) mp 245–246 °C; MS *m/z*: 221 (M⁺); UV $\lambda_{\text{max}}^{\text{MeOH}}$ 236 nm (ϵ 17300), 326 (15600); $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 238 (19600), 290 (sh) (4800), 328 (17200); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 226 (23000), 302.5 (15600); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 234 (19300), 319 (17200); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 227 (21000), 316.5 (14200); IR $\nu_{\text{max}}^{\text{Nujol}}$ cm⁻¹: 3400 (br, NH), 1711 (endocyclic CO), 1655 (NCOME); $\nu_{\text{max}}^{\text{CHCl}_3}$ (at 0.05 M concentration) cm⁻¹: 3320 (br, NH), 1730 (endocyclic CO), 1645 (NCOME), 1580, 1500, 1400, 1380; NMR (DMSO-*d*₆) δ : 2.09 (3H, s, COMe), 3.28 (3H, s, NMe), 3.68 (3H, s, NMe), 8.61 [1H, s, C(2)-H], 9.5–10.6 [1H, br, N(7)-H]; NMR (CDCl₃) δ : 2.26 (3H, s, COMe), 3.43 (3H, s, NMe), 3.82 (3H, s, NMe), 8.11 [1H, s, C(2)-H]. Anal. Calcd for C₉H₁₁N₅O₂: C, 48.87; H, 5.01; N, 31.66. Found: C, 48.73; H, 5.00; N, 31.59. The MS, UV (MeOH), IR (CHCl₃), and ¹H-NMR (CDCl₃) spectra of this sample were in agreement with those reported¹⁵) for a “natural” sample of **9**.

8-Bromo-6-imino-1,9-dimethylpurine Hydriodide (13·HI) A mixture of **11** (4.10 g, 18 mmol) and MeI (12.77 g, 90 mmol) in AcNMe₂ (30 ml) was

stirred at 50 °C for 3.5 h. After cooling, the reaction mixture was kept in an ice bath for 30 min, and the precipitate that resulted was filtered off, washed with a little EtOH, and dried to give **13**·HI (6.60 g, 99%) as a colorless solid, mp 243—245 °C (dec.). Recrystallization from 95% (v/v) aqueous EtOH yielded an analytical sample as colorless prisms, mp 244—246 °C (dec.); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 265 nm (ϵ 15700); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 265 (16500); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 265 (16500); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 265 (13900), 300 (sh) (4200); NMR (DMSO- d_6) δ : 3.76 and 3.79 (3H each, s, NMe's), 8.71 [1H, s, C(2)-H], 9.17 and 10.00 (1H each, br, =NH₂⁺). *Anal.* Calcd for C₇H₈BrN₅·HI: C, 22.72; H, 2.45; N, 18.93. Found: C, 22.72; H, 2.40; N, 18.90.

8-Bromo-6-imino-1,9-dimethylpurine (13) A solution of **13**·HI (4.07 g, 11 mmol) in warm H₂O (60 ml) was brought to pH 11 by addition of 10% aqueous Na₂CO₃, and the mixture was immediately cooled in an ice bath. The colorless crystals that deposited were filtered off, washed with a little H₂O, and dried to give **13** (2.12 g, 80%), mp 204—219 °C. Recrystallization from AcOEt afforded an analytical sample as colorless prisms, mp 216.5—218 °C; MS m/z : 243, 241 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 264 nm (ϵ 13300), 290.5 (sh) (4900); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 265 (15900); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 265 (15900); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 264 (13700), 295 (sh) (4800); IR $\nu_{\text{max}}^{\text{Nujol}}$ 3255 cm⁻¹ (NH); NMR (DMSO- d_6) δ : 3.44 and 3.59 (3H each, s, NMe's), 7.23 (1H, br, NH), 8.09 [1H, s, C(2)-H]. *Anal.* Calcd for C₇H₈BrN₅: C, 34.73; H, 3.33; N, 28.93. Found: C, 34.69; H, 3.20; N, 29.17.

Conversion of 13 into 8 and 9 Anhydrous NaOAc (dried over P₂O₅ at 2 mmHg and 75 °C for 6 h) (6.15 g, 75 mmol) was dissolved in AcOH (150 ml) by application of heat, and then **13** (1.21 g, 5 mmol) was added. The mixture was heated under reflux for 5 h, and the reaction mixture was concentrated *in vacuo* to leave a pale brownish residue, which was dissolved in H₂O (150 ml). The resulting solution was passed through a column of Dowex 50W-X8 (H⁺) (175 ml), and the column was eluted with H₂O (ca. 1 l) until the eluate became neutral, followed by 3% aqueous NH₃ (ca. 1.4 l). The ammoniacal eluate was concentrated *in vacuo* to leave a pale yellowish solid (1.16 g). After having been dried over P₂O₅ at 2 mmHg and 75 °C for 6 h, the solid was continuously extracted with AcOEt for 24 h by using a Soxhlet extractor. The AcOEt extract was concentrated *in vacuo*, and the residue was recrystallized from AcOEt to give **9** (378 mg, 34%) as colorless needles, mp 248—249.5 °C (dec.). This sample was identical (by comparison of the IR spectrum and TLC behavior) with the one obtained from **8** by acetylation (*vide supra*).

On the other hand, the insoluble solid left in the above AcOEt extraction was recrystallized from EtOH-H₂O (90:1, v/v), and the almost colorless crystals that resulted were filtered off and dried over P₂O₅ at 2 mmHg and 75 °C for 15 h to furnish **8** (319 mg, 36%), mp > 300 °C. Further recrystallization from EtOH and drying in the same manner produced an analytical sample of **8** as colorless needles, mp > 300 °C; MS m/z : 179 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 291.5 nm (ϵ 12300); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 221 (28000), 278 (10400); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 220 (24500), 285 (12000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 280 (14600), 310 (sh) (4800); IR $\nu_{\text{max}}^{\text{Nujol}}$ cm⁻¹: 3380, 3270, 3230 (NH's), 1694 (CO); NMR (CF₃CO₂D) δ : 3.66 and 4.10 (3H each, s, NMe's), 8.58 [1H, s, C(2)-H]. *Anal.* Calcd for C₇H₈N₅O: C, 46.92; H, 5.06; N, 39.09. Found: C, 46.77; H, 5.10; N, 38.92. This sample was identical (by comparison of the IR spectrum and TLC mobility) with the one obtained from **12** by methylation (*vide supra*) and was found to be slightly hygroscopic, forming a hydrate, mp > 300 °C. On the basis of elemental analysis, the hydrate was estimated to contain ca. 1.75 equivalent mole of H₂O of crystallization.

8-Bromo-N⁶,9-dimethyladenine (14) A mixture of **13**·HI (370 mg, 1 mmol) and 1 N aqueous NaOH (5 ml) was stirred at 55 °C for 35 min. The reaction mixture was kept in a refrigerator overnight, and the colorless prisms that deposited were filtered off, washed with H₂O, and dried to give **14** (213 mg, 88%), mp 185—186.5 °C. Recrystallization from AcOEt provided an analytical sample as colorless prisms, mp 186.5—188 °C; MS m/z : 243, 241 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 273 nm (ϵ 17000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 270 (20300); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 274 (17900); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 274 (17700); IR $\nu_{\text{max}}^{\text{Nujol}}$ 3335 cm⁻¹ (NH); NMR (see the text). *Anal.* Calcd for C₇H₈BrN₅: C, 34.73; H, 3.33; N, 28.93. Found: C, 34.53; H, 3.19; N, 29.00.

N⁶,9-Dimethyl-8-oxoadenine (15) i) From **14**: A stirred mixture of **14** (968 mg, 4 mmol) and 1 N aqueous NaOH (40 ml) was heated under reflux for 1.5 h. After cooling, the reaction mixture was neutralized by addition of 10% aqueous HCl, concentrated *in vacuo* to a small volume, and kept in a refrigerator overnight. The colorless precipitate that resulted was filtered off, washed with H₂O, and dried to afford crude **15** (770 mg), mp > 300 °C. Recrystallization from EtOH produced **15** (514 mg, 72%) as colorless needles, mp > 300 °C. Further recrystallization from EtOH yielded an analytical sample, mp > 300 °C; MS m/z : 179 (M⁺); UV $\lambda_{\text{max}}^{95\% \text{ EtOH}}$ 274 nm (ϵ 16500); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 1) 275 (14100); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 7) 274 (17000); $\lambda_{\text{max}}^{\text{H}_2\text{O}}$ (pH 13) 284 (18000); IR (see the text); NMR (DMSO- d_6) δ : 2.94 (3H, d,

$J = 5$ Hz, NHMe), 3.22 [3H, s, N(9)-Me], 6.39 (1H, q, $J = 5$ Hz, NHMe), 8.10 [1H, s, C(2)-H], 10.07 [1H, dull, N(7)-H]. *Anal.* Calcd for C₇H₈N₅O: C, 46.92; H, 5.06; N, 39.09. Found: C, 46.79; H, 5.09; N, 39.36.

ii) From **8**: A stirred mixture of **8** (1.08 g, 6 mmol) and 1 N aqueous NaOH (30 ml) was heated under reflux for 1 h. The reaction mixture was worked up in a manner similar to that described above for **15** under method (i), giving **15** (977 mg, 90%) as a colorless solid, mp > 300 °C. Recrystallization from EtOH produced a pure sample as colorless needles, mp > 300 °C. This sample was identical (by comparison of the IR spectrum and TLC behavior) with the one obtained by method (i).

Acknowledgment This work was supported by a Grant-in-Aid for Scientific Research (No. 01571148) from the Ministry of Education, Science and Culture, Japan. We are also grateful to Dr. Guido Cimino (Naples) for a personal communication about the location of a sample and spectral charts of compound **9** isolated from the natural source.

References and Notes

- 1) Paper XLII in this series, T. Fujii and T. Saito, *Chem. Pharm. Bull.*, **38**, 1886 (1990).
- 2) a) D. Ackermann and P. H. List, *Naturwissenschaften*, **48**, 74 (1961); b) *Idem*, *Z. Physiol. Chem.*, **323**, 192 (1961).
- 3) a) D. Ackermann and P. H. List, *Z. Physiol. Chem.*, **309**, 286 (1957); b) *Idem*, *ibid.*, **318**, 281 (1960).
- 4) a) W. Bergmann and D. C. Burke, *J. Org. Chem.*, **21**, 226 (1956); b) W. Bergmann and M. F. Stempien, Jr., *ibid.*, **22**, 1575 (1957).
- 5) a) R. J. Quinn, R. P. Gregson, A. F. Cook, and R. T. Bartlett, *Tetrahedron Lett.*, **21**, 567 (1980); b) A. F. Cook, R. T. Bartlett, R. P. Gregson, and R. J. Quinn, *J. Org. Chem.*, **45**, 4020 (1980).
- 6) E. Cullen and J. P. Devlin, *Can. J. Chem.*, **53**, 1690 (1975).
- 7) a) H. Nakamura, H. Wu, Y. Ohizumi, and Y. Hirata, *Tetrahedron Lett.*, **25**, 2989 (1984); b) H. Wu, H. Nakamura, J. Kobayashi, M. Kobayashi, Y. Ohizumi, and Y. Hirata, *Bull. Chem. Soc. Jpn.*, **59**, 2495 (1986).
- 8) H. Wu, H. Nakamura, J. Kobayashi, Y. Ohizumi, and Y. Hirata, *Tetrahedron Lett.*, **25**, 3719 (1984).
- 9) R. J. Capon and D. J. Faulkner, *J. Am. Chem. Soc.*, **106**, 1819 (1984).
- 10) R. Fathi-Afshar and T. M. Allen, *Can. J. Chem.*, **66**, 45 (1988). For a detailed study on the pharmacological activities of agelasimines A and B, see R. Fathi-Afshar, T. M. Allen, C. A. Krueger, D. A. Cook, A. S. Clanachan, R. Vriend, H. P. Baer, and C. E. Cass, *Can. J. Physiol. Pharmacol.*, **67**, 276 (1989).
- 11) The structure of a purino-diterpene compound isolated from an acetylated mixture of a crude methanolic extract of the same sponge has been determined by X-ray analysis: T. Nakatsu, D. J. Faulkner, G. K. Matsumoto, and J. Clardy, *Tetrahedron Lett.*, **25**, 935 (1984).
- 12) a) H. Kanatani, H. Shirai, K. Nakanishi, and T. Kurokawa, *Nature* (London), **221**, 273 (1969); b) M. Dorée, P. Guerrier, and N. J. Leonard, *Proc. Natl. Acad. Sci. U.S.A.*, **73**, 1669 (1976); c) M. Yoshikuni, K. Ishikawa, M. Isobe, T. Goto, and Y. Nagahama, *ibid.*, **85**, 1874 (1988).
- 13) 1-Methyladenine (**1**) has also been isolated from the starfish *Asterias amurensis*.^{12a)}
- 14) D. Jamieson and P. Davis, *Eur. J. Pharmacol.*, **67**, 295 (1980).
- 15) G. Cimino, A. De Giulio, S. De Rosa, S. De Stefano, R. Puliti, C. A. Mattia, and L. Mazzarella, *J. Nat. Prod.*, **48**, 523 (1985).
- 16) T. Fujii, T. Saito, and S. Mori, *Heterocycles*, **27**, 1145 (1988).
- 17) a) T. Fujii, S. Sakurai, and T. Uematsu, *Chem. Pharm. Bull.*, **20**, 1334 (1972); b) K. K. Ogilvie, S. L. Beaucage, and M. F. Gillen, *Tetrahedron Lett.*, **1978**, 1663; c) *Idem*, *ibid.*, **1978**, 3203; d) M. Hedayatullah, *J. Heterocycl. Chem.*, **19**, 249 (1982).
- 18) a) M. Ikehara and M. Kaneko, *Tetrahedron*, **26**, 4251 (1970); b) M. Ikehara and Y. Ogiso, *J. Carbohydr. Nucleosides, Nucleotides*, **1**, 401 (1974).
- 19) a) P. Brookes and P. D. Lawley, *J. Chem. Soc.*, **1960**, 539; b) H. G. Windmueller and N. O. Kaplan, *J. Biol. Chem.*, **236**, 2716 (1961); c) J. W. Jones and R. K. Robins, *J. Am. Chem. Soc.*, **85**, 193 (1963); d) N. J. Leonard and T. Fujii, *Proc. Natl. Acad. Sci. U.S.A.*, **51**, 73 (1964); e) J. A. Montgomery and H. J. Thomas, *J. Org. Chem.*, **30**, 3235 (1965); f) E. P. Lira, *J. Heterocycl. Chem.*, **5**, 863 (1968); g) M. Rasmussen and N. J. Leonard, *J. Am. Chem. Soc.*, **89**, 5439 (1967); h) T. Fujii and T. Saito, *Chem. Pharm. Bull.*, **33**, 3635 (1985); i) T. Fujii, T. Saito, and N. Terahara, *ibid.*, **34**, 1094 (1986).

- 20) For the Dimroth rearrangement of 1,9-disubstituted adenines to *N*⁶,9-disubstituted adenines, see refs. 19*h* and 19*i* and earlier references cited therein.
- 21) For the appearance of this type of carbonyl absorption in the 1690—1720 cm⁻¹ region, see *a*) J. Altman and D. Ben-Ishai, *J. Heterocycl. Chem.*, **5**, 679 (1968); *b*) T. Fujii and T. Saito, *Chem. Pharm. Bull.*, **21**, 1954 (1973); *c*) Y. Maki, K. Kameyama, M. Suzuki, M. Sako, and K. Hirota, *J. Chem. Res. (M)*, **1984**, 3601.
- 22) *a*) M. Ikehara, H. Tada, and K. Muneyama, *Chem. Pharm. Bull.*, **13**, 1140 (1965); *b*) M. Ikehara, H. Tada, and M. Kaneko, *Tetrahedron*, **24**, 3489 (1968); *c*) M. Ikehara and M. Kaneko, *Chem. Pharm. Bull.*, **18**, 2401 (1970).
- 23) The acid dissociation constant of the conjugate acid of **13** has been determined by us to be 8.54 ± 0.02 at 40°C and ionic strength 1.0. The details will be reported elsewhere shortly.