


Nitro- and Nitroso-derivatives of 3-Methylene-1,2-dithioles

By R. J. S. BEER* and R. J. GAIT

(The Robert Robinson Laboratories, University of Liverpool, Liverpool L69 3BX)

Summary In ω -nitroso-3-methylene-1,2-dithioles, the nitroso-group takes precedence over other groups (*e.g.* nitro-) in adopting the position *cis* to the dithiole S-S link.

STRUCTURE (I) was assigned to the product obtained by the action of nitrous acid on 2,5-diphenyl-6a-thiathiophthen (II) (and on 5-phenyl-1,2-dithiol-3-ylideneacetophenone)



R = H; λ_{\max} 314,464 nm (ϵ 10,000, 26,800)
 R = Me; λ_{\max} 316,468 nm (ϵ 10,500, 23,600)
 R = Br; λ_{\max} 318,477 nm (ϵ 6440, 15,600)

on the basis of spectroscopic evidence.¹ Subsequent study of the structure by X-ray crystallography² has shown that

there is a very short "contact" (2.034 Å) between a sulphur atom and the oxygen atom of the nitroso-group, and it is clear that the dimensions suggested by a single classical structure of type (I) are considerably modified in the actual molecule.

The three nitroso-compounds previously reported,¹ (I), (IIIa), and (IIIb), and two further compounds, (IIIc) and (IIId), obtained by nitrosation of 2-dimethylamino-5-phenyl-6a-thiathiophthen and of 2,5-bismethylthio-6a-thiathiophthen respectively, all have similar absorption spectra, with maxima at *ca.* 330 and *ca.* 420 nm. We infer that the five compounds are of the same structural type, *i.e.* derivatives of the system (IV).

We now report the preparation of structurally related nitro-compounds, (V; R = H) and (V; R = Me), obtained by condensation of 3-methylthio-5-phenyl-1,2-dithiolium methosulphate with nitromethane and nitroethane in acetic acid-pyridine. Compound (V; R = H), which is also produced by the action of mercuric acetate on 2-methylthio-3-nitro-5-phenyl-6a-thiathiophthen,¹ reacts readily with bromine in benzene solution giving the mono-substitution product (V; R = Br). The position of bromination can be inferred by comparison of n.m.r. spectra.

The three nitro-compounds clearly have similar structures, but the product obtained by nitrosation of compound (V; R = H) shows absorption maxima at 327 and 415 nm, closely resembling the nitroso-compounds (I) and (IIIa—d); it is therefore formulated as (IIIe).

This structure is confirmed by the X-ray crystallographic

study described in the following communication,³ which also gives definitive structural data for the nitro-compound (V; R = Br). It is of particular interest that the S–O contact in the nitro-compound is considerably longer than the S–O distances in the two nitroso-compounds for which data are available.

We conclude that the nitroso-group takes precedence over other groups (CO·Ph, CS·SMe, CS·NMe₂, NO₂) in competition for the apparently advantageous position adjacent to the dithiole S–S link.

(Received, January 15th, 1970; Com. 065.)

¹ R. J. S. Beer, D. Cartwright, R. J. Gait, R. A. W. Johnstone, and S. D. Ward, *Chem. Comm.*, 1968, 688.

² P. L. Johnson and I. C. Paul, *J. Amer. Chem. Soc.*, 1969, **91**, 781.

³ K. I. G. Reid and I. C. Paul, following communication.