The Photolysis of Schneider's 1-Phenyliminopyridinium Ylides

By VICTOR SNIECKUS* and GORDON KAN

(Department of Chemistry, University of Waterloo, Waterloo, Canada)

Summary Photolysis of the pyridinium ylides (4) and (5) produces, besides high yields of aniline and (8) and (9)respectively, the additional products (10) [from (4) in benzene] and (12) [from (5) in diethylamine] which suggest a phenyl nitrene intermediate in these reactions.

RECENT investigations in the field of 1-iminopyridinium actions: (i) ring expansion to 1(1H), 2-diazepines $[(1) \rightarrow$ ylide photochemistry has uncovered three types of re-(2)];¹⁻³ (ii) rearrangement to 2-amino-pyridines [3,5-dimethyl- $(1a) \rightarrow (3a)$];^{3b} and (iii) photo-Curtius rearrangement [(1b) \rightarrow pyridine + MeCNO].^{3a} We now report on the photolysis of the 1-phenylpyridinium ylides (4) and (5), two of the first known examples of this class of compounds.4,5 Our work offers new information on some structural and mechanistic aspects of N-N ylide photochemistry, provides a new method for the generation of phenyl nitrene and thereby an alternate mechanistic probe with which to study the nature of its electronic state,⁶ and presents the first opportunity for direct comparison of the less complicated photochemical behaviour of an N-N ylide (5) with that of a similarly substituted, isoelectronic N-oxide (2,4,6-triphenylpyridine N-oxide).7

The ylide (4) was known to undergo a Sommelet-type rearrangement⁸ in many solvents at ambient temperatures



Photolysis of 1-aryliminopyridinium ylides^a

Compound				Products, % yield ^e			
	Reaction conditions		Solvent ^b	(8)	PhNH ₂	(10)	(12)
(4)	0—5°, 0—5°, 5—10°,	17 hr. 18 hr. 18 hr.¢	CH2Cl2 Et2NH C6H6	$16.5 \\ 33 \\ 12$	17d 35·5 59	24	
(5)	8—12°, 8—12°, 8—12°, 8—12°,	2 hr. 2 hr., N ₂ 2 hr., O ₂ 4 hr.	CH2Cl2 Et2NH Et2NH C6H6	(9) 79 73 70 79	12·5ª 89 91·5 24		$\frac{2}{2}$

^a Irradiations (40 ml) were conducted in a Rayonet Reactor using 3500 Å lamps in a Pyrex vessel similar to the one described by G. T. Tisue, S. Linke, and W. Lwowski, J. Amer. Chem. Soc., 1967, 89, 6303. Concentrations were ca. 2.5×10^{-3} M in starting ylides. ^b All solvents were dried and distilled before use.

⁶ Compounds (8) and (9) were obtained border dot. ⁶ Compounds (8) and (9) were obtained by preparative thick-layer or column chromatography (silica gel) and were characterized by comparison with authentic samples: (8);¹⁵ (9).⁺⁴ Yields of aniline and (12) were determined by g.l.c. analysis using naphthalene as an internal standard on a 6 ft. \times 1 in. column of 25% Carbowax 20 M and 3% KOH on Chromosorb W60 \times 80 mesh. For the purpose of direct comparison of retention times, compound (12) was prepared by the known procedure.¹³ ^d The low yield is due to formation of aniline hydrochloride in this solvent.

• A higher temperature was required to avoid freezing the benzene solution. not be prevented and 10.5% of (7) was isolated. Under these conditions, thermal rearrangement could

to yield the pyridine derivative (7).^{4b,5} This thermal dark reaction † was completely avoided by carrying out the photolysis at $0-5^{\circ}$. The Table summarizes the results obtained from the photolysis of (4) in several solvents.[‡] Rather than ring expanding to a 1-aryl-1(1H), 2-diazepine (6),⁹ compound (4) undergoes fragmentation in methylene

chloride or diethylamine solution to yield 2-methyl-4,6diphenylpyridine (8) and aniline as the only isolable products. In benzene solution, an additional product, the aldehyde (10) [m.p. 110-113°; i.r. (CHCl₃) 1695 cm⁻¹ (C=O); n.m.r. (CDCl₃) τ -0.1 (s, 1H, CHO), 2.3 (m, 2H, pyridine protons), 1.9-2.5 (m, 10H, aromatic protons)] is

[‡] The fact that (4) is light-sensitive had received brief comment (see ref. 5).

[†] For example, ylide (4) on standing in methylene chloride at room temperature in the dark gave 44% yield of (7). The n.m.r. spectrum (CDCl₃) of (7) [τ 1.60 (m, 2H, pyridine protons), 1.88–3.18 (m, 14H, aromatic protons), 5.11 (br s, 2H, exchanged with D_2O , NH_2), 5.67 (s, 2H, PhCH₂–] confirmed Schneider's original far-sighted assignment.^{4b}

CHEMICAL COMMUNICATIONS, 1970

produced.§ By analogy with our observations^{3a} on the ylide (1b), a reasonable mechanism for the decomposition of (4) is via heterolytic N-N bond cleavage to produce (8) and phenyl nitrene. In agreement with reactions characteristic of aryl nitrenes,¹⁰ the formation of aniline and aldehyde (10) is observed. The latter product presumably arises via C-H insertion of singlet phenyl nitrene 68,11 to form (11) followed by hydrogen abstraction and hydrolysis of the resulting Schiff base on chromatography.12 That the formation of (10) may involve a cage mechanism receives support from the observation that the photolysis of a mixture of (8) and phenyl azide does not yield (10). Further evidence for the intermediacy of phenyl nitrene was obtained from the much cleaner photochemical breakdown of ylide (5).

As seen from the Table, the photolysis¶ of ylide (5) also results in fragmentation to 2,4,6-triphenylpyridine (9) in high yield while the amounts of aniline obtained are again strongly dependent on solvent. Furthermore, irradiation in diethylamine under nitrogen produced minor but reproducible amounts of 2-diethylamino-3H-azepine (12) indicative of the intermediacy of phenyl nitrene.13,14 Photofragmentation was not quenched and (12) was still produced when the irradiation was carried out under oxygen. Although attempts to sensitize the reaction have been complicated by the intense absorption of ylide (5) in the u.v. and visible region, the above results are suggestive of the reaction proceeding via an excited singlet ylide and

$$(5) \xrightarrow{4\partial, Et_2 \text{NH}} (9) + \left[\swarrow N \right] \longrightarrow \bigotimes_{N} N Et_2$$

$$(12) \xrightarrow{+ PhNH_2}$$

singlet phenyl nitrene species. Comparative evidence which points to the involvement of singlet phenyl azide and singlet phenyl nitrene in the photolysis of phenyl azide is available.^{6a,14} The failure to detect triplet phenyl nitrene in the photolysis of (5) at 77° k in methylene chloride-Fluorolube by e.s.r. measurements supports our tentative conclusions.

Support of this work by the National Research Council of Canada is gratefully acknowledged. Thanks are due to D. Burkholder and Dr. J. M. McIntosh for exploratory work.

(Received, December 2nd, 1969; Com. 1838.)

§ The structure of (10) was established by oxidation to the known 4,6-diphenylpyridine-2-carboxylic acid: C. Gastaldi, Gazzetta, 1922, 52, 174.

 \P In contrast to ylide (4), (5) was found to be stable in all solvents at room temperature in the dark.

 J. Streith and J.-M. Cassal, Bull. Soc. chim. France, 1969, 2175 and references therein.
 S. Sasaki, K. Kanematsu, and A. Kaheki, Chem. Comm., 1969, 432; S. Sasaki et al., J. Org. Chem., in the press.
 (a) V. Snieckus, Chem. Comm., 1969, 831; (b) A. Balasubramanian, J. M. McIntosh, and V. Snieckus, J. Org. Chem., in the press.
 (a) W. Schneider and F. Seebach, Ber., 1921, 54, 2285; (b) W. Schneider and K. Weiss, *ibid.*, 1928, 61, 159; (c) W. Schneider, 1924, 438, 115; (d) K. Dimroth and K. H. Wolf, in "Newer Methods of Preparative Organic Chemistry," ed. W. Foerst, Academic Proce New York and London 1964, vol. 114, p. 257. Academic Press, New York and London, 1964, vol. III, p. 357. ⁵ K. Dimroth, G. Arnoldy, S. von Eicken, and G. Schiffler, Annalen, 1967, 604, 221.

⁶ For recent work, see (a) R. A. Odum and A. M. Aaronson, J. Amer. Chem. Soc., 1969, 91, 5680 and references therein; (b) R. J. Sundberg, R. H. Smith, jun., and J. E. Bloor, *ibid.*, p. 3392.
⁷ P. L. Kumler and O. Buchardt, Chem. Comm., 1968, 1321. Review: C. Kaneko, J. Synthetic Org. Chem., Japan, 1968, 26, 758.
⁸ A. W. Johnson, "Ylid Chemistry," Academic Press, New York and London, 1966, p. 268.
⁹ A single example of a 1-alkyl triphenyldiazepine has been recorded: E. Klingsberg, Abstracts of the American Chemical Society Macting Scattering 1067 of 268.

¹⁰ For recent reviews, see J. H. Boyer in "Mechanisms of Molecular Migrations," ed. B. S. Thyagarajan, Interscience, New York, 1969, vol. 2, p. 296 and G. L'Abbe, *Chem. Rev.*, 1969, 69, 345.
¹¹ G. Smolinsky and B. I. Feuer, J. Amer. Chem. Soc., 1964, 86, 3085.
¹² Intramolecular examples of such sequences are known: G. Smolinsky, J. Amer. Chem. Soc., 1961, 83, 2489.
¹³ W. H. F. Dearing and R. A. Odum Tetrahedron. 1966, 22, 81.

- ¹³ W. von E. Doering and R. A. Odum, *Tetrahedron*, 1966, 22, 81. ¹⁴ J.-S. Splitter and M. Calvin, *Tetrahedron Letters*, 1968, 1445.
- ¹⁵ W. Zecher and F. Krohnke, Chem. Ber., 1961, 94, 698.