

- and H. A. O. Hill, *Chem. Commun.*, 34 (1970); (c) T. Sakurai, J. P. Fox, and L. L. Ingraham, *Inorg. Chem.*, **10**, 1105 (1971); (d) G. Tauscher, R. Dreos, G. Costa, and M. Green, *J. Chem. Soc., Chem. Commun.*, 413 (1973); (e) J. H. Espenson and R. Russell, *Inorg. Chem.*, **13**, 7 (1974).
- (4) W. C. Troglor, R. C. Stewart, and L. G. Marzilli, *J. Am. Chem. Soc.*, **96**, 3697 (1974).
- (5) (a) L. M. Ludwick and T. L. Brown, *J. Am. Chem. Soc.*, **91**, 5188 (1969); (b) T. L. Brown, L. M. Ludwick, and R. S. Stewart, *ibid.*, **94**, 384 (1972); (c) R. J. Guschl and T. L. Brown, *Inorg. Chem.*, **12**, 2815 (1973); (d) R. J. Guschl, R. S. Stewart, and T. L. Brown, *ibid.*, **13**, 417 (1974); (e) R. J. Guschl and T. L. Brown, *ibid.*, **13**, 959 (1974).
- (6) T. W. Swaddle, *Coord. Chem. Rev.*, **14**, 217 (1974).
- (7) D. J. McLennan, *Acc. Chem. Res.*, **9**, 281 (1976).
- (8) D. D. Perrin, W. L. F. Armarego, and D. R. Perrin, "Purification of Laboratory Chemicals", Pergamon Press, London, 1966.
- (9) (a) G. N. Schrauzer and R. J. Windgassen, *J. Am. Chem. Soc.*, **88**, 3738 (1966); (b) G. N. Schrauzer, *Inorg. Synth.*, **11**, 61 (1968).
- (10) R. N. Moore and R. K. Ziegler, Report No. LA-2367, plus Addenda, Los Alamos Scientific Laboratory, Los Alamos, N.Mex., 1959.
- (11) F. J. Kezdy, J. Jaz, and A. Bruylants, *Bull. Soc. Chim. Belg.*, **67**, 687 (1958).
- (12) N. Yamazaki and Y. Hohokabe, *Bull. Chem. Soc. Jpn.*, **44**, 63 (1971), and references therein.
- (13) G. J. Palenik, D. A. Sullivan, and D. V. Naik, *J. Am. Chem. Soc.*, **98**, 1177 (1976), and references therein.
- (14) L. Seibles, Ph.D. Thesis, University of Cincinnati, 1976.
- (15) C. J. Weschler and E. Deutsch, *Inorg. Chem.*, **12**, 2682 (1973).
- (16) (a) D. E. C. Corbridge, *Top. Phosphorus Chem.*, **6**, 235 (1969); (b) W. C. Troglor, L. A. Epps, and L. G. Marzilli, *Inorg. Chem.*, **14**, 2748 (1975).
- (17) (a) H. A. O. Hill and K. G. Morallee, *J. Chem. Soc. A*, 554 (1969); (b) H. A. O. Hill, K. G. Morallee, and G. Pellizer, *J. Am. Chem. Soc.*, **94**, 276 (1972).
- (18) W. C. Troglor, R. C. Stewart, L. A. Epps, and L. G. Marzilli, *Inorg. Chem.*, **13**, 1564 (1974).
- (19) J. P. Fox, R. Banninger, R. T. Proffitt, and L. L. Ingraham, *Inorg. Chem.*, **11**, 2379 (1972).
- (20) (a) For a compilation of relevant references see T. G. Appleton, H. C. Clark, and L. E. Mauzer, *Coord. Chem. Rev.*, **10**, 335 (1973); (b) G. Pellizer, G. R. Tauszik, and G. Costa, *J. Chem. Soc., Dalton Trans.*, 317 (1973).
- (21) Supplementary material.
- (22) (a) F. Basolo and R. G. Pearson, "Mechanisms of Inorganic Reactions", 2nd ed, Wiley, New York, N.Y., 1967; (b) R. G. Wilkins, "The Study of Kinetics and Mechanisms of Reactions of Transition Metal Complexes", Allyn and Bacon, Boston, 1972.
- (23) (a) J. H. Espenson and R. G. Russell, *Inorg. Chem.*, **13**, 7 (1974); (b) G. Tauscher, R. Dreos, and G. Costa, *J. Chem. Soc., Chem. Commun.*, 413 (1973).
- (24) (a) A. A. Frost and R. G. Pearson, "Kinetics and Mechanism", 2nd ed, Wiley, New York, N.Y., 1966, pp 123-159. (b) S. W. Benson, "The Foundations of Chemical Kinetics", McGraw-Hill, New York, N.Y., 1960, pp 493-547. (c) From ref 22b, pp 181-184, using the relationship $K_{os} = [4\pi Na^3/3000] \exp(-U(a)/kT)$, where $U(a)$ is given by $(-|Ze|\mu' \cos \theta)Da^2$, with θ generally taken to be 180° and $a = 4 \text{ \AA}$.
- (25) A. L. McClellan, "Tables of Experimental Dipole Moments", W. H. Freeman and Co., San Francisco, Calif., 1963.
- (26) W. L. Jolly, "The Synthesis and Characterization of Inorganic Compounds", Prentice-Hall, Englewood Cliffs, N.J., 1970, p 99.
- (27) Reference 22b, p 183.
- (28) J. M. Pratt and R. G. Thorp, *Adv. Inorg. Chem. Radiochem.*, **12**, 375 (1969).
- (29) (a) D. H. McDaniel and H. C. Brown, *J. Org. Chem.*, **23**, 420 (1958); (b) J. March, "Advances in Organic Chemistry. Reactions, Mechanisms and Structure", McGraw-Hill, New York, N.Y., 1968, pp 238-245; (c) J. Hine, "Physical Organic Chemistry", 2nd ed, McGraw-Hill, New York, N.Y., 1962, pp 85-93; (d) J. Miller, "Aromatic Nucleophilic Substitution", Elsevier, London, 1968, pp 78-88; (e) J. E. Leffler and E. Grunwald, "Rates and Equilibria of Organic Reactions", Wiley, New York, N.Y., 1963, p 172.

Contribution from the Laboratoire de Chimie Minérale Moléculaire, Equipe de Recherche Associée au CNRS, Institut de Mathématiques et Sciences Physiques, Parc Valrose, 06034, Nice Cedex, France

Alkoxyfluorophosphoranes. 3. Synthesis and Characterization of Some Di-, Tri-, and Tetraalkoxyfluorophosphoranes

DANIEL D. POULIN, CLAIRE DEMAY, and JEAN G. RIESS*

Received November 23, 1976

AIC60841O

Several dialkoxyfluorophosphoranes $\text{PhPF}_2(\text{OR})_2$ ($\text{R} = \text{CH}_2\text{CCl}_3$, CH_2CHCl_2 , $\text{CH}(\text{CH}_3)(\text{CN})$, CH_2CF_3) and $\text{MePF}_2(\text{OCH}_2\text{CF}_3)_2$ have been isolated and characterized by NMR, IR, and mass spectrometry and elemental analysis. Their formation can be facilitated by the disproportionation reaction $2\text{R}'\text{PF}_3(\text{OR}) \rightarrow \text{R}'\text{PF}_2(\text{OR})_2 + \text{R}'\text{PF}_4$. These compounds are stable with respect to both further disproportionation and decomposition into phosphonates. The tetraalkoxyphosphorane $\text{PhP}(\text{OCH}_2\text{CF}_3)_4$ has also been isolated. Trialkoxyfluorophosphoranes $\text{PhPF}(\text{OR})_3$ ($\text{R} = \text{CH}_2\text{CCl}_3$, CH_2CHCl_2 , $\text{CH}(\text{CH}_3)(\text{CN})$, CH_2CF_3) and $\text{MePF}(\text{OCH}_2\text{CF}_3)_3$ have been identified in mixtures but could not be separated from the other members of the $\text{R}'\text{PF}_{4-n}(\text{OR})_n$ series.

Introduction

Phosphoranes containing simultaneously fluorine atoms and alkoxy groups were generally found to be unstable with respect to their decomposition into phosphoryl compounds according to^{1,2}




We have shown³ that they could be stabilized by introducing electron-withdrawing substituents in the alkoxy moiety, and we have recently reported the preparation of a series of stable monoalkoxyfluorophosphoranes⁴ through the general reaction⁵



We now wish to report the preparation and isolation of several stable dialkoxydifluorophosphoranes of type $\text{R}'\text{PF}_2(\text{OR})_2$ and of the tetraalkoxyphosphorane $\text{PhP}(\text{OCH}_2\text{CF}_3)_4$ as well as the identification of several trialkoxyfluorophosphoranes of type $\text{R}'\text{PF}(\text{OR})_3$ which, although they could

not be isolated in a pure form, were unambiguously characterized both by spectral determinations and by their transformation into the corresponding dialkoxydifluorophosphoranes.

Previous studies in the literature which produced nonionic multialkoxyfluorophosphoranes stable at room temperature appear to be limited to^{6,7}



On the other hand, the related aryloxyfluorophosphoranes were found to be more stable than their alkoxy analogues.^{8,9}

Experimental Section

All manipulations of fluorophosphoranes were carried out under a dry nitrogen atmosphere. Reactions involving CH_3PF_4 were carried

Table I. NMR Parameters for Di-, Tri-, and Tetraalkoxyphosphoranes of Types R'PF₂(OR)₂ (I), R'PF(OR)₃ (II), and R'P(OR)₄ (III)^a

Type	R	R'	δ_{CH_3} ^a	δ_{CH_2}	δ_{CH}	$\delta_{\text{F(P)}}$	δ_{P}	$^3J_{\text{HCCH}}$	$^3J_{\text{HCOP}}$	$^3J_{\text{HCCF}}$	$^2J_{\text{HCP}}$	$^3J_{\text{HCPF}}$	$^1J_{\text{FP}}$
I	CH ₂ CCl ₃	C ₆ H ₅		4.62		38.0	59.1		7.5				824
I	CH ₂ CCl ₂ H	C ₆ H ₅		4.34	5.78	41.9	58.0	5.4	10.5				822
I	CH(CH ₃)(CN)	C ₆ H ₅	1.95		5.55	39.8	58.8	7.0					826
I	CH ₂ CF ₃	C ₆ H ₅		4.27		43.2	58.5		10.3	8.3			808
II	CH ₂ CF ₃	C ₆ H ₅		4.19		40.5	57.2		8.4	8.3			820
III	CH ₂ CF ₃	C ₆ H ₅		4.01			58.7		5.0	8.3			
I	CH ₂ CF ₃	CH ₃	1.55	4.11		30.1 ^b	45.4		10.5	7.9	21.2	12.7	819
II	CH ₂ CF ₃	CH ₃	1.52	4.00		34.0			8.3	7.9	19.5	14.2	813
III	CH ₂ CF ₃	CH ₃	1.47	3.96					5.6	7.9	17.1		

^a ¹H, ¹⁹F chemical shifts in ppm relative to internal Me₄Si and CFCl₃, ³¹P relative to external H₃PO₄; coupling constants in Hz. ^b Chemical shift relative to external CFCl₃.

out using standard vacuum techniques in a system constructed with Pyrex glass with stopcocks lubricated with Apiezon N grease. C₆H₅PF₄, CH₃PF₄, and the alkoxyethyl ethers were prepared as previously reported.⁴

The ¹H and ¹⁹F spectra were routinely recorded with a JEOL C-60 HL instrument of samples containing an approximate 10% solution of the compound in CDCl₃. The ³¹P spectra were measured on neat liquids. Infrared spectra were recorded on films between KBr plates with a Perkin-Elmer 577 spectrometer. Mass spectra were measured with a CEC 21-130 instrument. The purity of all compounds has been checked by elemental analysis and by their spectrochemical properties which are given in Tables I and II.

Preparation of Bis(2,2-dichloroethoxy)difluorophenylphosphorane, C₆H₅PF₂(OCH₂CCl₂H)₂. (CH₃)₃SiOCH₂CCl₂H (4.29 g, 22.9 mmol) was added dropwise to C₆H₅PF₄ (2.07 g, 11.2 mmol) at -30 °C. The reaction mixture was allowed to warm slowly to room temperature and was then heated to 70 °C for 30 min. (CH₃)₃SiF (2.06 g, 21.6 mmol) was trapped at -75 °C. C₆H₅PF₂(OCH₂CCl₂H)₂ (3.61 g, 86%) was recovered as a colorless liquid by distillation (70 °C (10⁻² mmHg)).

Anal. Calcd for C₁₀H₁₁Cl₄F₂O₂P: C, 32.11; H, 2.97; F, 10.16; Cl, 37.92; P, 8.28. Found: C, 32.15; H, 3.02; F, 10.42; Cl, 38.02; P, 8.23.

Under more drastic conditions (90 °C, 6 h) and with 3 equiv of the silyl ether, the formation of 80% C₆H₅PF(OCH₂CCl₂H)₃ (δ_{F} 36.1 ppm, J_{FP} = 821 Hz) was observed with 20% C₆H₅PF₂(OCH₂CCl₂H)₂.

Preparation of Bis(1-cyanoethoxy)difluorophenylphosphorane, C₆H₅PF₂[OCH(CH₃)(CN)]₂. (CH₃)₃SiOCH(CH₃)(CN) (3.07 g, 25.9 mmol) was added dropwise to C₆H₅PF₄ (2.29 g, 12.2 mmol) at -30 °C. The reaction mixture was allowed to warm slowly to room temperature and was then heated to 85 °C for 30 min. (CH₃)₃SiF (2.13 g, 23.3 mmol) was trapped at -75 °C. C₆H₅PF₂[OCH(CH₃)(CN)]₂ (3.18 g, 91%) was recovered from the reaction vessel as a yellow viscous nonvolatile liquid.

Anal. Calcd for C₁₂H₁₃F₂N₂O₂P: C, 50.35; H, 4.58; F, 13.28; N, 9.79; P, 10.82. Found: C, 50.74; H, 4.58; F, 13.01; N, 9.91; P, 10.77.

Under more drastic reaction conditions (100 °C, 1 h) the formation of ca. 5% C₆H₅PF[OCH(CH₃)(CN)]₃ (δ_{F} 34.8 ppm, J_{FP} = 824 Hz) was observed together with 95% C₆H₅PF₂[OCH(CH₃)(CN)]₂.

Preparation of Bis(2,2,2-trifluoroethoxy)difluorophenylphosphorane, C₆H₅PF₂(OCH₂CF₃)₂. (CH₃)₃SiOCH₂CF₃ (2.08 g, 12.1 mmol) was added dropwise to C₆H₅PF₄ (1.07 g, 5.82 mmol) at 0 °C. The reaction mixture was allowed to warm slowly to room temperature and to react for 30 h at this temperature. The (CH₃)₃SiF produced was allowed to escape from the reaction and was not collected. C₆H₅PF₂(OCH₂CF₃)₂ (1.81 g, 90%), a colorless liquid, was purified by removing the last traces of (CH₃)₃SiF and (CH₃)₃SiOCH₂CF₃ under vacuum 5 × 10⁻⁴ mm (20 °C).

Anal. Calcd for C₁₀H₉F₈O₂P: C, 34.90; H, 2.63; F, 44.17; P, 9.00. Found: C, 35.03; H, 2.61; F, 43.25; P, 9.16.

Preparation of Bis(2,2,2-trifluoroethoxy)difluoromethylphosphorane, CH₃PF₂(OCH₂CF₃)₂. (CH₃)₃SiOCH₂CF₃ (4.46 g, 25.9 mmol) and CH₃PF₄ (1.57 g, 13.0 mmol) were allowed to react in a sealed tube at 45 °C for 65 h. Vacuum fractionation afforded CH₃PF₂(OCH₂CF₃)₂ (2.81 g, 77%), as a stable colorless liquid collected at -45 °C, and (CH₃)₃SiF (2.66 g) which was trapped at -116 °C. A small amount of nonvolatile and as yet unidentified white solid (0.18 g) remained in the reaction tube.

Anal. Calcd for C₃H₇F₈O₂P: C, 21.29; H, 2.50; F, 53.88; P, 10.98. Found: C, 21.55; H, 2.51; F, 52.62; P, 10.99.

Preparation of Tetrakis(2,2,2-trifluoroethoxy)phenylphosphorane, C₆H₅P(OCH₂CF₃)₄. (CH₃)₃SiOCH₂CF₃ (26.8 g, 156 mmol) was

added dropwise to C₆H₅PF₄ (2.13 g, 11.6 mmol) and the mixture was allowed to react at 70 °C for 63 h. The (CH₃)₃SiF produced was distilled out of the reaction system but not collected. The unreacted (CH₃)₃SiOCH₂CF₃ was pumped away (5 × 10⁻⁴ mmHg (22 °C)) and C₆H₅P(OCH₂CF₃)₄ (3.19 g, 54%) was collected and purified by distillation (64 °C, 5 × 10⁻⁴ mmHg). The compound is a white solid at room temperature (mp 62 °C).

Anal. Calcd for C₁₄H₁₃F₁₂O₄P: C, 33.35; H, 2.60; F, 45.22; P, 6.14. Found: C, 33.55; H, 2.53; F, 44.07; P, 6.33.

Preparation of Bis(2,2,2-trichloroethoxy)difluorophenylphosphorane, C₆H₅PF₂(OCH₂CCl₃)₂. (CH₃)₃SiOCH₂CCl₃ (3.97 g, 17.9 mmol) was allowed to react with C₆H₅PF₄ (0.77 g, 4.2 mmol) in a sealed tube (~25 cm³) for 40 h at 100 °C. Vacuum fractionation of the volatile products afforded (CH₃)₃SiOCH₂CCl₃ (2.81 g, 12.7 mmol) which was collected at -45 °C and (CH₃)₃SiF (0.51 g, 5.5 mmol) which was trapped at -116 °C. Thus only 1.2 equiv of silyl ether had been consumed; previous work⁴ has shown that the monoalkoxy compound is formed under these conditions. In a second step, the colorless nonvolatile liquid which remained in the reaction tube and consisted mainly of C₆H₅PF₂(OCH₂CCl₃) was pumped in for 12 h at 60 °C and C₆H₅PF₄ (0.28 g, 1.5 mmol) was recovered as the volatile product. C₆H₅PF₂(OCH₂CCl₃)₂ (1.01 g, 93%) was isolated from the reaction flask as a colorless liquid.

Anal. Calcd for C₁₀H₉Cl₆F₂O₂P: C, 27.12; H, 2.03; Cl, 48.04; P, 6.99. Found: C, 27.24; H, 2.13; Cl, 48.23; P, 7.08.

Under more drastic reaction conditions (150 °C, 40 h) the formation of 5% C₆H₅PF(OCH₂CCl₃)₃ (δ_{F} 33.5, J_{FP} = 806 Hz) was observed along with 95% C₆H₅PF₂(OCH₂CCl₃)₂.

Preparation of Mixtures of the Tri- and Tetraalkoxyphosphoranes CH₃PF(OCH₂CF₃)₃ and CH₃P(OCH₂CF₃)₄. Attempts to prepare CH₃PF(OCH₂CF₃)₃ and CH₃P(OCH₂CF₃)₄ always resulted in mixtures of the two products. A typical example using a 6:1 molar ratio of (CH₃)₃SiOCH₂CF₃ and CH₃PF₄ afforded, after 75 h at 75 °C, a mixture containing ca. 90% CH₃PF(OCH₂CF₃)₃ and 10% CH₃P(OCH₂CF₃)₄ (by NMR; no dialkoxydifluorophosphorane detected). An 8:1 molar ratio of the reagents resulted, after 154 h at 100 °C, in a ca. 1:1 mixture of the tri- and tetrasubstituted products.

Results and Discussion

Dialkoxydifluorophosphoranes of type R'PF₂(OR)₂ (I) (R' = Ph or Me, R = CH₂CCl₃, CH₂CHCl₂, CH(CH₃)(CN), CH₂CF₃) were isolated in good yields through reaction 2 and their thermal stability was found to be good (no changes were observed in any of these compounds after 6 months at room temperature, and they could be heated for 2 h at 100 °C without noticeable change). This further illustrates the stabilizing effect of electron-attracting substituents in the alkoxy groups which thus opposes the tendency of compounds of type I to decompose into phosphonates, according to eq 1.

These dialkoxydifluorophosphoranes were also found to be stable with respect to their disproportionation according to



i.e., eq 3 lies to the left and the mono- and trialkoxy compounds were not detectable by NMR. This contrasts with the behaviour of the monoalkoxyfluorophosphoranes, which were found to undergo such reactions easily^{2,4}




Table II. Mass Spectra of Di- and Tetraalkoxyphosphoranes [m/e , assignments (relative intensity)]


$\text{PhPF}_2(\text{OCH}_2\text{CCl}_3)_2$	$\text{PhPF}_2(\text{OCH}_2\text{CHCl}_2)_2$	$\text{PhPF}_2[\text{OCH}(\text{CH}_3)(\text{CN})]_2$	$\text{PhPF}_2(\text{OCH}_2\text{CF}_3)_2$	$\text{MePF}_2(\text{OCH}_2\text{CF}_3)_2$	$\text{PhP}(\text{OCH}_2\text{CF}_3)_4$
440 $\text{C}_{10}\text{H}_9\text{Cl}_2\text{F}_2\text{O}_2\text{P}$ (1)	353 $\text{C}_{10}\text{H}_9\text{FCl}_2\text{O}_2\text{P}$ (3)	286 $\text{C}_{12}\text{H}_{13}\text{F}_3\text{N}_2\text{O}_2\text{P}$ (2)	344 $\text{C}_{10}\text{H}_9\text{F}_2\text{O}_2\text{P}$ (7)	267 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (3)	406 $\text{C}_{11}\text{H}_{12}\text{F}_2\text{O}_3\text{P}$ (15)
421 $\text{C}_{10}\text{H}_9\text{Cl}_2\text{F}_2\text{O}_2\text{P}$ (1)	315 $\text{C}_{10}\text{H}_9\text{FCl}_2\text{O}_2\text{P}$ (2)	267 $\text{C}_{12}\text{H}_{13}\text{FN}_2\text{O}_2\text{P}$ (2)	325 $\text{C}_{10}\text{H}_9\text{F}_2\text{O}_2\text{P}$ (20)	263 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (70)	405 $\text{C}_{11}\text{H}_{12}\text{F}_2\text{O}_3\text{P}$ (100)
405 $\text{C}_{10}\text{H}_9\text{Cl}_2\text{F}_2\text{O}_2\text{P}$ (1)	276 $\text{C}_{10}\text{H}_9\text{FCl}_2\text{O}_2\text{P}$ (7)	216 $\text{C}_{12}\text{H}_{13}\text{FNOP}$ (60)	267 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (33)	239 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (3)	385 $\text{C}_{11}\text{H}_{12}\text{F}_2\text{O}_3\text{P}$ (3)
293 $\text{C}_8\text{H}_7\text{Cl}_3\text{F}_2\text{OP}$ (21)	259 $\text{C}_8\text{H}_7\text{F}_2\text{Cl}_2\text{OP}$ (38)	165 $\text{C}_8\text{H}_8\text{NOP}$ (96)	246 $\text{C}_8\text{H}_7\text{F}_8\text{OP}$ (9)	183 $\text{C}_8\text{H}_7\text{F}_8\text{OP}$ (100)	325 $\text{C}_{10}\text{H}_9\text{F}_2\text{O}_2\text{P}$ (17)
274 $\text{C}_8\text{H}_7\text{Cl}_3\text{FOP}$ (1)	161 $\text{C}_8\text{H}_7\text{Cl}_2\text{F}_2\text{OP}$ (100)	163 $\text{C}_8\text{H}_8\text{NOP}$ (96)	245 $\text{C}_8\text{H}_7\text{F}_8\text{OP}$ (100)	181 $\text{C}_8\text{H}_7\text{F}_8\text{OP}$ (2)	322 $\text{C}_{10}\text{H}_9\text{F}_2\text{O}_2\text{P}$ (9)
163 $\text{C}_6\text{H}_5\text{F}_2\text{OP}$ (100)	160 $\text{C}_6\text{H}_5\text{FO}_2\text{P}$ (32)	162 $\text{C}_8\text{H}_7\text{FNO}_2\text{P}$ (28)	185 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (16)	161 $\text{C}_8\text{H}_7\text{F}_8\text{OP}$ (3)	301 $\text{C}_7\text{H}_7\text{F}_5\text{O}_3\text{P}$ (7)
161 $\text{C}_6\text{H}_5\text{F}_2\text{OP}$ (12)	143 $\text{C}_6\text{H}_5\text{FOCl}_2\text{P}$ (24)	158 $\text{C}_8\text{H}_7\text{FOP}$ (11)	165 $\text{C}_8\text{H}_7\text{F}_8\text{O}_2\text{P}$ (10)	101 $\text{F}_2\text{O}_2\text{P}$ (81)	224 $\text{C}_8\text{H}_7\text{F}_5\text{O}_2\text{P}$ (6)
131 $\text{C}_2\text{H}_5\text{Cl}_2$ (9)	105 $\text{H}_2\text{F}_2\text{Cl}_2\text{P}$ (25)	103 $\text{C}_8\text{H}_8\text{NOP}$ (15)	163 $\text{C}_6\text{H}_5\text{F}_2\text{OP}$ (20)	99 $\text{C}_2\text{H}_5\text{F}_2\text{O}$ (1)	176 $\text{C}_8\text{H}_7\text{F}_5\text{OP}$ (12)
103 $\text{H}_2\text{F}_2\text{O}_2\text{P}$ (17)	97 $\text{C}_2\text{H}_5\text{Cl}_2$ (21)	96 $\text{C}_8\text{H}_7\text{NO}$ (58)	162 $\text{C}_6\text{H}_5\text{F}_2\text{OP}$ (13)	83 $\text{CH}_3\text{F}_2\text{P}$ (43)	143 $\text{C}_6\text{H}_5\text{FOP}$ (28)
96 $\text{C}_2\text{H}_5\text{Cl}_2$ (18)	96 $\text{C}_2\text{H}_5\text{Cl}_2$ (32)	94 CHF_2NP (18)	143 $\text{C}_6\text{H}_5\text{FOP}$ (7)	81 C_2F_3 (2)	141 $\text{C}_6\text{H}_5\text{O}_2\text{P}$ (26)
77 C_6H_5 (30)	94 C_2Cl_2 (15)	78 $\text{C}_8\text{H}_8\text{OP}$ (17)	96 $\text{CH}_3\text{FO}_2\text{P}$ (50)	69 CF_3 (3)	133 $\text{C}_2\text{H}_5\text{F}_3\text{OP}$ (27)
61 $\text{C}_2\text{H}_5\text{Cl}$ (6)	77 C_6H_5 (34)	77 C_6H_5 (82)	83 CH_3FOP (24)	64 CH_2FP (2)	83 CH_3FOP (16)
51 HFP (11)	61 CH_3OP (15)	76 CH_3FNP (16)	77 C_6H_5 (42)	61 CH_2OP (2)	77 C_6H_5 (62)
50 FP (6)	51 HFP (23)	56 $\text{C}_2\text{H}_5\text{ON}$ (28)	69 CF_3 (12)	51 HFP (1)	69 CF_3PF_2 (16)
	50 FP (12)	54 $\text{C}_3\text{H}_6\text{N}$ (100)	61 CH_2OP (18)		51 HFP , CF_2H (36)
		51 HFP (50)	51 HFP (31)		
		50 FP (24)			

There, the use of the electron-attracting substituents allows the retention of the five-coordinate character of the phosphorus atom, i.e., their decomposition into phosphonates can be avoided, but does not oppose the scrambling reaction 4. That such a phenomenon is not observed for the dialkoxyfluorophosphoranes, I, can be attributed to the reluctance of the alkoxy group to assume an apical position and reflects the difference in apicophilicity between fluorine atoms and alkoxy groups.

Profit can even be gained from reaction 4 in the preparation of the dialkoxyfluorophosphoranes, I. Thus a sample of pure $\text{PhPF}_2(\text{OCH}_2\text{CCl}_3)_2$ was obtained by first preparing the monoalkoxyfluorophosphorane and then shifting equilibrium 4 to the left by continuously pumping off the more volatile PhPF_4 .

When the substitution series $\text{R}'\text{PF}_{4-n}(\text{OR})_n$ is examined, one observes a marked increase in the difficulty of the substitution between $n = 2$ and $n = 3$; i.e., the reaction conditions have to become much more drastic if one wants to add an alkoxy group in the apical position. Thus, for example, the dialkoxy compound $\text{PhPF}_2(\text{OCH}_2\text{CCl}_3)_2$ was formed quantitatively from a 2:1 molar ratio of reactants after 30 min at 70 °C while the trialkoxy compound $\text{PhPF}(\text{OCH}_2\text{CCl}_3)_3$ was formed only in 20% yield from a 3:1 ratio after 10 h at 75 °C. The dialkoxy compound $\text{MePF}_2(\text{OCH}_2\text{CF}_3)_2$ was obtained in 90% yield from a 2:1 molar ratio of reactants after 47 h at room temperature while the trialkoxy compound $\text{MePF}(\text{OCH}_2\text{CF}_3)_3$ was obtained in 50% yield after heating a 6:1 molar ratio of the reactants for 72 h at 74 °C. Similarly the dialkoxy compound $\text{PhPF}_2(\text{OCH}_2\text{CF}_3)_2$ was formed in a 90% yield from a 2:1 molar ratio of reactants after 48 h at room temperature while it took 40 days at the same temperature to obtain a 33% yield of the trialkoxy compound from a 3:1 molar ratio of the reactants.

It appears difficult to stop the substitution process at the trialkoxy compounds, $\text{R}'\text{PF}(\text{OR})_3$, II, the conditions for tetra substitution being too similar. We were thus unable to obtain them in a pure form, but only in mixtures with the tetraalkoxy derivatives, although these mixtures were free of phosphonates. They were characterized by NMR and, in the case of $\text{PhPF}(\text{OCH}_2\text{CF}_3)_3$, by the addition to the mixture of the amount of $\text{R}'\text{PF}_4$ which is needed to convert all the constituents of the mixture into the dialkoxydifluoro compound according to



The completely substituted tetraalkoxy derivative $\text{R}'\text{P}(\text{OCH}_2\text{CF}_3)_4$ was identified in a mixture (with the trialkoxy compound), for $\text{R}' = \text{Me}$, and isolated in the case of $\text{R}' = \text{Ph}$.

No evidence for the formation of ionic species either directly or upon aging was encountered in these series. All the compounds were soluble in pentane, benzene, CHCl_3 , CH_2Cl_2 , and CH_3CN .

The NMR parameters are reported in Table I. The ^{19}F and ^{31}P chemical shifts and J_{PF} coupling constants are in agreement with those found previously for monoalkoxyfluorophosphoranes.⁴ It is noteworthy that although J_{HCOF} is quite large (5–10 Hz) further J_{HCOF} coupling is not resolved. For the trifluoroethoxyphosphoranes, for which we now have two complete substitution series $\text{R}'\text{PF}_n(\text{OCH}_2\text{CH}_3)_{4-n}$ ($\text{R}' = \text{Ph}$ and Me ; $n = 0-3$), one observes a systematic increase in J_{HCOF} with n and at the same time a gradual downfield shift for the methylene proton signals. The four alkoxy groups of $\text{PhP}(\text{OCH}_2\text{CF}_3)_4$ remain equivalent in the ^1H NMR and give a well-resolved doublet of quadruplets down to -50 °C; these lines coalesce in one broad symmetric hump at ca. -70 °C suggesting that the alkoxy groups become nonequivalent at

lower temperatures and that their exchange between the apical and equatorial sites of a trigonal bipyramid is rapid above this temperature.

The mass spectra (Table II) show a weak or nondetectable molecular ion. The fragments corresponding to the loss of F and OR are important. It is noteworthy that the loss of OR is generally extensive, while the loss of R is not detected; this is the opposite to what has been observed previously in the case of the analogous monoalkoxyfluorophenyl (or methyl)-phosphoranes, for which the breaking of the OR bond was an important feature.⁴

The infrared spectra show the expected $\nu_{as}(\text{P-F})$ bands in the range of 920–800 cm^{-1} . These relatively low values are consistent with apical fluorines in a trigonal-bipyramidal structure¹⁰ and comparable to those found in series of compounds R_3PF_2 having strongly electroattracting R groups.¹¹ The P–O–C linkage gives rise to three absorption bands near 1130, 1020, and 780 cm^{-1} .¹² The Ph–P linkage produces a characteristic vibration near 1450 cm^{-1} .¹² No vibrations assignable to $\nu(\text{P=O})$ stretching vibrations were detected.

Acknowledgment. The authors are grateful to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this work.

Registry No. $\text{C}_6\text{H}_5\text{PF}_2(\text{OCH}_2\text{CCl}_3)_2$, 43227-68-5; $\text{C}_6\text{H}_5\text{PF}_2(\text{OCH}_2\text{CCl}_2\text{H})_2$, 63325-03-1; $\text{C}_6\text{H}_5\text{PF}_2(\text{OCH}(\text{CH}_3)(\text{CN}))_2$, 55509-54-1; $\text{C}_6\text{H}_5\text{PF}_2(\text{OCH}_2\text{CF}_3)_2$, 63325-04-2; $\text{C}_6\text{H}_5\text{PF}(\text{OCH}_2\text{CF}_3)_3$, 63325-05-3; $\text{C}_6\text{H}_5\text{P}(\text{OCH}_2\text{CF}_3)_4$, 63325-06-4; $\text{CH}_3\text{PF}_2(\text{OCH}_2\text{CF}_3)_2$, 63325-07-5; $\text{CH}_3\text{PF}(\text{OCH}_2\text{CF}_3)_3$, 63325-08-6; $\text{CH}_3\text{P}(\text{OCH}_2\text{CF}_3)_4$, 63325-09-7; $(\text{CH}_3)_3\text{SiOCH}_2\text{CCl}_2\text{H}$, 51209-79-1; $(\text{CH}_3)_3\text{SiOCH}(\text{CH}_3)(\text{CN})$, 41309-99-3; $(\text{CH}_3)_3\text{SiOCH}_2\text{CF}_3$, 56859-55-3; $(\text{CH}_3)_3\text{SiOCH}_2\text{CCl}_3$, 41262-38-8; $\text{C}_6\text{H}_5\text{PF}_4$, 666-23-9.

References and Notes

- (1) R. Schmutzler, *J. Chem. Soc.*, 4551 (1964).
- (2) D. U. Robert, G. N. Flatau, A. Cambon, and J. G. Riess, *Tetrahedron*, **29**, 1877 (1973).
- (3) D. U. Robert, G. N. Flatau, C. Demay, and J. G. Riess, *J. Chem. Soc., Chem. Commun.*, 1127 (1972).
- (4) J. G. Riess and D. U. Robert, *Bull. Soc. Chim. Fr.*, 425 (1975).
- (5) R. Schmutzler, "Halogen Chemistry", Vol. 2, V. Gutman Ed., Academic Press, New York, N.Y. 1967.
- (6) E. D. Young and W. B. Fox, *Inorg. Nucl. Chem. Lett.*, **7**, 1033 (1971).
- (7) I. L. Knunyants, V. V. Tynlenova, E. Y. Pervova, and R. N. Sterlin, *Izv. Akad. Nauk SSSR, Ser. Khim.*, **10**, 1797 (1964).
- (8) S. C. Peake, M. Fild, M. J. C. Hewson, and R. Schmutzler, *Inorg. Chem.*, **10**, 2723 (1971).
- (9) L. Kolditz, K. Lehmann, W. Wiecker, and A. R. Grimmer, *Z. Anorg. Allg. Chem.*, **360**, 259 (1968).
- (10) E. L. Muetterties, W. Mahler, and R. Schmutzler, *Inorg. Chem.*, **2**, 613 (1963).
- (11) J. Grosse and R. Schmutzler, *Phosphorus*, **4**, 49 (1974).
- (12) D. E. C. Corbridge, *Top. Phosphorus Chem.*, **6**, (1969).

Contribution from the Department of Chemistry,
University of Alberta, Edmonton, Alberta, Canada T6G 2G2

Silicon-Transition Metal Chemistry. 9.¹ Synthesis, Properties, and Structure of the Rhenium Acylsilane Complex $\text{fac-Re}(\text{CO})_3(\text{diphos})[\text{C}(\text{O})\text{SiPh}_3]$

J. R. ANGLIN, H. P. CALHOUN, and W. A. G. GRAHAM*

Received November 9, 1976

AIC60815Q

The reaction of $[\text{Re}(\text{CO})_4(\text{diphos})]\text{ClO}_4$ with Ph_3SiLi yields the acylsilane complex $\text{fac-Re}(\text{CO})_3(\text{diphos})[\text{C}(\text{O})\text{SiPh}_3]$ (**1**). A stereospecific synthesis of the ^{13}C -enriched precursor $[\text{fac-Re}(\text{CO})_3(^{13}\text{CO})(\text{diphos})]\text{ClO}_4$ is given. When **1** is prepared from this material, the majority of the enrichment remains mutually *cis* to phosphorus, although there is some scrambling to positions *trans* to phosphorus. Treatment of **1** and the unisolated SiMePh_2 analogue with Et_3OPF_6 yields the cationic carbene species $[\text{fac-Re}(\text{CO})_3(\text{diphos})[\text{C}(\text{OEt})\text{SiPh}_3]]^+$ and $[\text{fac-Re}(\text{CO})_3(\text{diphos})[\text{C}(\text{OEt})\text{SiMePh}_2]]^+$, respectively. The thermal decarbonylation of **1** yields *mer-Re* $(\text{CO})_3(\text{diphos})\text{SiPh}_3$, with loss of CO mutually *cis* to phosphorus. The crystal and molecular structure of **1** has been determined from three-dimensional x-ray data collected by counter techniques. The material crystallizes in the space group $P2_1/n$ of the monoclinic system with $a = 20.675$ (4) Å, $b = 11.372$ (2) Å, $c = 18.271$ (2) Å, and $\beta = 108.04$ (9)°. The observed density 1.54 (1) g cm^{-3} may be compared with 1.554 g cm^{-3} calculated for $Z = 4$. The structure was refined by a full-matrix least-squares procedure to an R value of 0.0401 for 3604 observations having $I/\sigma(I) > 3.0$. The structure consists of discrete monomers of slightly distorted octahedral geometry having facial substitution. The $\text{Ph}_3\text{SiC}(\text{O})$ ligand is bonded to rhenium through the acyl carbon and contains a planar $\text{SiC}(\text{O})\text{Re}$ group. This plane roughly bisects the P–Re–P angle and is such that the SiPh_3 group is remote from the diphos ligand. Bond distances in the acyl group are Si–C = 1.969 (10) Å, Re–C = 2.199 (9) Å, and O–C = 1.23 (1) Å. Bond angles are Re–C–Si = 127.9 (5)°, Re–C–O = 124.4 (6)°, and Si–C–O = 107.7 (6)°.

Introduction

Transition metal-acyl complexes are important organometallic intermediates that have deservedly received much study.^{2,3} Related carbamoyl,⁴ alkoxycarbonyl,⁴ and formyl⁵ complexes are also known, in which the metal-bonded carbonyl carbon is attached not to carbon but to N, O, or H, respectively. However, little attention has apparently been directed toward the study of transition metal-acyl derivatives of the heavier group 4 elements, although acylsilanes, acylgermanes, and acylstannanes are well-known.⁶

Transition metal-acyl complexes are most commonly prepared as shown in eq 1 and 2; however, it has not been



feasible to extend these routes to the synthesis of acyl com-

plexes of the heavier group 4 elements. The carbonyl insertion reaction (eq 1) has not been observed to occur in bonds between metals and heavier group 4 elements. The appropriate acyl halides and anhydrides for eq 2 are unknown for heavier group 4 elements, and our attempts to prepare such materials have proved unsuccessful.

Accordingly, we adopted the route (eq 3) where a transition



metal-carbonyl cation is treated with the appropriate lithium reagent, $\text{R}_3\text{M}'\text{Li}$. We reported recently the synthesis of $\text{fac-Re}(\text{CO})_3(\text{diphos})[\text{C}(\text{O})\text{SiPh}_3]$ (**1**) by this route and noted its remarkable spectroscopic properties.⁷ The cation $[\text{Re}(\text{CO})_4(\text{diphos})]^+$ had been chosen for this work because (i) rhenium acyls are known to have good thermal and photolytic stability,³ (ii) the related cation $[\text{trans-Re}(\text{CO})_4(\text{PPh}_3)_2]^+$ is