

slope of $\ln a$ vs. t plots. Reactions were typically first order for several half lives, and the correlation coefficients were 0.995 or better. Second-order rate constants were obtained from the slope of the pseudo-first-order rate constant vs. the diene concentration. Three diene concentrations were used to obtain each second-order rate constant. The second-order rate constants were obtained in triplicate.

In a typical experiment 10 mL of diene solution was placed in a rubber capped reaction flask containing a stirring bar. The flask was put in a thermostated bath and stirred for 30-60 min. At the same time the bath water was circulated through a jacket especially designed to house the IR cell (Precision Cells, Inc., part no. 20710). The reaction flask was connected to the IR cell by Teflon tubing. With the IR instrument set at 2128 cm^{-1} and in the time drive mode about 0.1 mL of approximately 1 M Hfe in the reaction solvent was injected into the reaction flask. The reaction mixture was then forced through the IR cell by inert-gas pressure supplied by a nitrogen or argon tank while the absorbance vs. time data was obtained. Rate data was obtained for 60-300 s. Kinetics data were obtained at 20, 30, and 40°C for activation

energy studies. Carbon monoxide was used in place of the inert gas in each step of the experiment to test its retarding effect on the rate. Similarly rates were measured by using a solvent that was successively 0.1, 0.2, and 0.3 M in trichlorosilane.

Procedure b. For the slower reactions, reaction mixtures were made up in a constant temperature bath and aliquots were periodically transferred to the IR cell.

Acknowledgment. The author wishes to thank the University of Missouri-Kansas City School of Graduate Studies for financial support and Ms. Tohien Huang, Ms. To-Nguyen Huang, and Mr. Joel Sartwell for technical assistance.

Registry No. Hfe, 33361-70-5; Dfe, 91002-46-9; 1,3-butadiene, 106-99-0; 2-methyl-1,3-butadiene, 78-79-5; 2,3-dimethyl-1,3-butadiene, 513-81-5; *trans*-1,3-pentadiene, 2004-70-8; *cis*-1,3-pentadiene, 504-60-9; 1,3-cyclohexadiene, 592-57-4; 3-methylene-cyclohexene, 1888-90-0; (*E,E*)-2,4-hexadiene, 5194-51-4; deuterium, 7782-39-0.

Organometallic Chemistry. 21. Silyl Trifluoromethanesulfonate (Triflate)-Boron Trichloride (Tribromide) Complexes¹

George A. Olah,* Khosrow Laali, and Omar Farooq

Donald P. and Katherine B. Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, University Park, Los Angeles, California 90089

Received March 19, 1984

Trimethyl- and triethylsilyl triflate form with BCl_3 and BBr_3 strongly polarized donor-acceptor complexes. The deshielding of the ^{29}Si NMR shifts [$\Delta\delta_{\text{Si}}$ (BCl_3) 28.81 and 32.26, respectively] and shielding of ^{11}B NMR shifts characteristic for tetracoordinated boron are consistent with the development of partial positive charge at silicon involving *O*-coordination of boron halide with the triflates. Data also show that no ionization to silicenium ions takes place. Similarly, dimethylisopropyl- and tri-*n*-butylsilyl triflate gave donor-acceptor complexes on reaction with BCl_3 [$\Delta\delta_{\text{Si}}$ 30.51 and 25.18, respectively]. In general BCl_3 was found to interact more strongly than BBr_3 . The reaction of a number of other alkylarylsilyl triflates or dialkylsilyl ditriflates with BX_3 was also investigated. The deshielding of ^{29}Si NMR shifts diminished when the steric bulkiness of the alkyl ligands on silicon was increased, such as in the case of (*i*-Pr) $_3\text{SiOTf}$ or $\text{Me}_2(t\text{-Bu})\text{SiOTf}$. Ligand exchange of the triflates with BX_3 is a competing process in the reactions, the rate of which increases significantly by increasing the temperature or reaction time. The reaction of tris(alkylthio)silyl triflate with BCl_3 at low temperature led only to ligand exchange. The reaction of Me_3SiOR ($\text{R} = \text{Me}, \text{Et}, \text{Ph}$) with BBr_3 (-30°C) or with BCl_3 (-75°C) gave Me_3SiBr , Me_3SiCl , and the corresponding ROSiX_2 indicative of initial complexation followed by rapid Si-O cleavage. The initial complexation, however, could not be observed by NMR. The reaction of AlCl_3 and AlBr_3 with silyl triflates resulted only in ligand exchange. Attempts to utilize the $\text{R}_3\text{SiOTf-BX}_3$ systems as electrophilic trialkylsilylating agents for aromatics were unsuccessful.

Introduction

We have previously shown that trimethylsilyl halides form donor-acceptor complexes with Lewis acid halides such as AlBr_3 , AlCl_3 , and BI_3 in solvents such as CS_2 , CH_2Cl_2 , and CH_2Br_2 .² Spectroscopic evidence for the formation of these polarized complexes is based on the observed deshielding of the ^{29}Si NMR shifts and a considerable sharpening of ^{27}Al NMR resonances, indicating a pseudotetrahedral environment around the aluminum atoms. However, the Si-halogen bonds are not ionized and no trivalent silicenium ions were formed in these systems (eq 1).



Recently, the generation of a stable silicenium ion was reported by Lambert by hydride abstraction from tris(isopropylthio)silane with trityl perchlorate in CH_2Cl_2 at room temperature. The poorly resolved ^{29}Si NMR signal was deshielded by only 17 ppm as compared to the silane precursor. This was attributed to "polymer formation", and it was suggested that the ^{29}Si NMR signal of the ion has a very slow relaxation time.³ We feel this evidence is inadequate, and further studies are needed to substantiate whether a truly electron-deficient silicenium ion was obtained. The formation of silicenium ylides and their

(1) For part 20 see: Olah, G. A.; Field, L. D.; Lammertsma, K.; Pacquin, D.; Suemmerman, K. *Nouv. J. Chim.* 1983, 7, 279.

(2) Olah, G. A.; Field, L. D. *Organometallics* 1982, 1, 1485.


(3) Lambert, J. B.; Schulz, W. J., Jr. *J. Am. Chem. Soc.* 1983, 105, 1671.

Table I. ^{29}Si and ^{11}B NMR Chemical Shifts of Silyl Triflate- BX_3 Complexes

| compound ^a | Lewis acid | $\delta^{29}\text{Si}$ | | | $\delta^{11}\text{B}$ complex |
|--|---|------------------------|---------|----------|----------------------------------|
| | | starting material | complex | exchange | |
| Me_3SiOTf | | 43.54 | | | |
| Me_3SiOTf | BCl_3 (-35°C) | | 72.35 | 29.02 | -2.62 |
| Me_3SiOTf | BBr_3 (rt) | | 61.52 | 25.74 | -2.45 |
| Me_3SiOTf | $\text{AlBr}_3/\text{CS}_2$ (rt) | | | 25.24 | |
| Me_3SiOTf | $\text{AlCl}_3/\text{CS}_2$ (rt) | 41.85 | | | |
| Me_3SiOTf | $\text{AlCl}_3/\text{CS}_2$ (-30°C) | 43.05 | | 29.07 | |
| Et_3SiOTf | | 44.46 | | | |
| Et_3SiOTf | BCl_3 (-35°C) | | 76.72 | 34.73 | -2.75 |
| $\text{Et}_2\text{Si}(\text{OTf})_2$ | | 13.35 | | | |
| $\text{Et}_2\text{Si}(\text{OTf})_2$ | BCl_3 (-35°C) | | 35.14 | 12.94 | -2.55 |
| $\text{Et}_2\text{Si}(\text{OTf})_2$ | $\text{BBr}_3/\text{CH}_2\text{Cl}_2$ (rt) | | 31.04 | | -2.45 |
| $i\text{-Pr}_3\text{SiOTf}$ | | 41.15 | | | |
| $i\text{-Pr}_3\text{SiOTf}$ | BCl_3 (-35°C) | | | 34.35 | |
| $n\text{-Bu}_3\text{SiOTf}$ | | 45.23 | | | |
| $n\text{-Bu}_3\text{SiOTf}$ | BCl_3 (-60°C) | | 70.41 | 31.81 | -3.09 |
| $\text{Me}_2(i\text{-Pr})\text{SiOTf}$ | | 45.05 | | | |
| $\text{Me}_2(i\text{-Pr})\text{SiOTf}$ | BCl_3 (-60°C) | | 75.56 | 33.27 | -2.82 |
| $\text{Me}_2(t\text{-Bu})\text{SiOTf}$ | | 43.50 | | | |
| $\text{Me}_2(t\text{-Bu})\text{SiOTf}$ | BCl_3 (-35°C) | 42.56 | | | |
| $\text{Me}_2(t\text{-Bu})\text{SiOTf}$ | BBr_3 (rt) | 43.79 | | 34.16 | |
| $\text{Ph}_2(\text{Me})\text{SiOTf}$ | | 11.54 | | | |
| $\text{Ph}_2(\text{Me})\text{SiOTf}$ | BBr_3 (rt) | | 29.79 | 8.42 | |
| $\text{Ph}_2(\text{Me})\text{SiOTf}$ | BCl_3 (-35°C) | 10.54 | 17.55 | | |

^a ^{29}Si NMR spectra of the starting triflates are recorded as neat solution at room temperature (rt).

stabilization with AlX_3 were also reported. The partial zwitterion character of these species is based on ^{29}Si and ^{27}Al NMR data.⁴




The silylating ability of trimethylsilyl triflate Me_3SiOTf is well-known. We have ourselves previously developed an improved synthetic method for the in situ generation of this reagent for silylating carboxylic acids, alcohols, phenols, mercaptans, and ketones.⁵ Our recent studies of $\text{R}_3\text{SO}_3\text{H}-\text{BX}_3$ systems and the finding that tricoordinated boron has a high tendency to coordinate triflate ligands⁶ led us to study the complexation of silyl trifluoromethanesulfonates (triflates) with BX_3 by NMR (^{29}Si , ^{11}B) and to search for suitable substituents on silicon to enhance the donor-acceptor complex formation. It was also attempted to use the trimethyl- and triethylsilyl triflate- BX_3 reagent for electrophilic silylation of alkylbenzenes.

Results and Discussion

When liquid BCl_3 was slowly added with efficient stirring to Me_3SiOTf at dry ice/acetone temperature, an oil layer was formed. The ^{29}Si NMR spectrum of the resulting oil recorded directly in BCl_3 at -35°C consisted of two absorptions, a deshielded signal at $\delta^{29}\text{Si}$ 72.35 for the complex ($\Delta\delta^{29}\text{Si}$ 28.81) and a shielded absorption at $\delta^{29}\text{Si}$ 29.02 for Me_3SiCl , formed by ligand exchange, in a 1:3 ratio. Upon standing at -75°C the complex slowly precipitated from BCl_3 solvent. After removal of excess BCl_3 , the complex was dissolved in SO_2ClF . The corresponding ^{11}B NMR spectrum of the complex was observed at -2.62 ppm,

viz., in the region of tetracoordinated boron compounds. Similarly, BBr_3 forms a complex that at room temperature shows two signals in the ^{29}Si NMR spectrum ($\delta^{29}\text{Si}$ 61.52, $\Delta\delta^{29}\text{Si}$ 18) and the ligand exchange product Me_3SiBr ($\delta^{29}\text{Si}$ 25.74). The corresponding ^{11}B NMR signal of the complex was observed at $\delta^{11}\text{B}$ -2.45. A variable-temperature ^{29}Si NMR study (-60 to -20°C) of $\text{Me}_3\text{SiOTf}-\text{BCl}_3$ system showed reversible exchange broadening of the ^{29}Si signal of the donor-acceptor complex with the solvent BCl_3 . The ligand exchange process is, however, irreversible, and Me_3SiCl increases slowly by increasing temperature at the expense of donor-acceptor complex. It is also noted that the expected $\text{B}(\text{OTf})\text{X}_2$ formed by ligand exchange does not show a separate signal from BCl_3 or BBr_3 solvent in the ^{11}B NMR spectra, suggesting that it undergoes exchange with the medium (eq 2). Whereas AlCl_3 and AlBr_3



were found to give donor-acceptor complexes with Me_3SiCl ,² with trimethylsilyl triflate only ligand exchange of chlorotrimethylsilane was observed. This is in agreement with the finding that AlX_3 readily reacts with triflic acid with the evolution of HX to form triflates.⁷

When triethylsilyl triflate was reacted with BCl_3 in a similar fashion in an NMR tube, the ^{29}Si NMR spectrum recorded at -35°C showed two absorptions at $\delta^{29}\text{Si}$ 76.72 and 34.73 for the donor-acceptor complex ($\Delta\delta^{29}\text{Si}$ 32.26) and Et_3SiCl , respectively, in a 1:2 ratio. Since the complex slowly precipitated out of solution at low temperature, in order to observe the ^{11}B NMR spectrum of the complex ($\delta^{11}\text{B}$ -2.75) it was necessary to remove excess BCl_3 and to dissolve the precipitate in SO_2ClF . Diethylsilyl ditriflate similarly formed a complex with BCl_3 at low temperature. The ^{29}Si NMR spectrum of the mixture again showed two absorptions at $\delta^{29}\text{Si}$ 35.14 and 12.94 for the donor-acceptor complex ($\Delta\delta^{29}\text{Si}$ 21.79) and the unchanged starting material, respectively, in a 1:4 ratio. Surprisingly, no ligand ex-

(4) Clegg, W.; Klingebell, U.; Neemann, J.; Sheldrick, G. M. *J. Organomet. Chem.* **1983**, *249*, 23.


(5) Olah, G. A.; Husain, A.; Gupta, B. G. B.; Sales, G. F.; Narang, S. C. *J. Org. Chem.* **1981**, *46*, 5212.

(6) Olah, G. A.; Laali, K.; Farooq, O., submitted for publication in *J. Org. Chem.*

(7) Olah, G. A.; Laali, K., unpublished results.

change product was formed. The corresponding ¹¹B NMR signal of the complex was at δ_{11B} -2.55. Treatment of Et₂Si(OTf)₂ with a dilute solution of BBr₃ in dry CH₂Cl₂ also resulted in a donor-acceptor complex with the absorption in the ²⁹Si NMR at δ_{29Si} 31.04 ($\Delta\delta_{29Si}$ 17.69) and the corresponding boron signal at δ_{11B} -2.45. *t*-Bu₂SiF₂ with triflic acid does not form *t*-Bu₂Si(OTf)₂. Similarly, *t*-Bu₂Si(OTf)₂ could not be obtained from its chloride and triflic acid.

i-Pr₃SiOTf on treatment with BCl₃ at low temperature did not give an observable complex. Only ligand exchange of triflate for Cl was observed, giving chlorotriisopropylsilane indicative of ionization followed by immediate quenching of the silicenium ion by chloride. *n*-Bu₃SiOTf formed a donor-acceptor complex with BCl₃ ($\Delta\delta_{29Si}$ 25.18) together with *n*-Bu₃SiCl in a 1:3 ratio at -60 °C. The corresponding ¹¹B NMR signal of the complex was at δ_{11B} -3.09 (SO₂ClF). Me₂(*i*-Pr)SiOTf similarly formed a donor-acceptor complex on addition of BCl₃ at dry ice/acetone temperature ($\Delta\delta_{29Si}$ 30.51) and Me₂(*i*-Pr)SiCl in a 0.8:1 ratio at -60 °C. ¹¹B NMR spectrum of the complex dissolved in SO₂ClF was observed at δ_{11B} -2.82. Me₂(*t*-Bu)SiOTf did not react with BCl₃. However, with BBr₃ ligand exchange was observed at ambient temperature. Bulkiness of the ligands on silicon therefore affects the stability of the donor-acceptor complexes. NMR data, however, cannot differentiate complexation on the sulfonyl or ether oxygen. Analogy with protonation (or complexation) of carboxylic acid esters, however, suggests that it is the ether oxygen complexation which results in Si-O bond cleavage with subsequent formation of the corresponding halosilane ligand exchange product.



In general, when the alkyl groups on silicon were replaced by phenyl, the extent of positive charge on silicon is reduced. This is in accord that diphenylmethylsilyl triflate gave an observable donor-acceptor complex on reaction with BBr₃ at room temperature, the ²⁹Si NMR absorption of which was δ_{29Si} 29.79 ($\Delta\delta_{29Si}$ 18.25). Subsequent ligand exchange was also observed, giving diphenylmethylsilyl bromide. On standing in BBr₃, side reactions occurred and the solution gradually darkened. With BCl₃ at low temperature the ²⁹Si NMR indicated only complex formation (δ_{29Si} 17.55, $\Delta\delta_{29Si}$ 6) and no ligand exchange was found. On the basis of the NMR data no apparent correlation between the magnitude of the positive charge on silicon, reflected in $\Delta\delta_{29Si}$ values, and the extent of cleavage vs. complex formation could be found.

The behavior of tris(isopropylthio)silyl triflate was also studied with BCl₃ in order to examine the influence of the alkylthio ligands on donor-acceptor complex formation. The product observed was chlorotris(isopropylthio)silane (δ_{29Si} 20.67) indicative of ligand exchange, and no intermediate donor-acceptor complex was observed.

Whereas the cleavage of trimethylsilyl ethers to trimethylsilyl halides by reaction with BBr₃ or BCl₃ is known,⁸ to our knowledge direct observation of the initial step, i.e., the formation of a donor-acceptor complex, has not been reported. We therefore examined the reaction

of Me₃SiOR (R = Me, Et, Ph) with BBr₃ at ambient temperature and at -30 °C as well as with BCl₃ at -75 °C. ²⁹Si NMR spectrum of a neat sample of Me₃SiOMe at room temperature gave a single absorption at δ_{29Si} 16.98 with the corresponding δ_{13C} shifts at -1.74 and 49.02 ppm at -20 °C. Slow addition of excess BBr₃ to this sample at -55 °C gave a colorless solution, the ²⁹Si spectrum of which showed only Me₃SiBr (δ_{29Si} 26.36). The ¹³C NMR spectrum of this sample showed two absorptions at δ_{13C} 4.12 (Me₃SiBr) and δ_{13C} 59.38. The latter is in close agreement with the reported values for methoxy attached to tricoordinated boron⁹ and is therefore assigned to MeOBBR₂ (δ_{11B} 26.27 at -30 °C). Similarly, low-temperature addition (-50 °C) of BBr₃ to Me₃SiOEt (δ_{29Si} 14.36) and Me₃SiOPh (δ_{29Si} 20.07) gave Me₃SiBr (δ_{29Si} 26.11 and 26.37). Excess BCl₃ reacted rather violently with Me₃SiOMe and Me₃SiOEt cooled to dry ice/acetone temperature. The ²⁹Si NMR spectra of both samples were consistent with Me₃SiCl (δ_{29Si} 30.91 and 30.70), and a donor-acceptor complex was not observed.

Attempted Silylation of Aromatics. Electrophilic silylation of aromatics under Friedel-Crafts condition has not so far been achieved. We felt of interest to study the reaction of the presently studied R₃SiOTf-BCl₃ (R = Me, Et) complexes with reactive aromatic compounds. When cumene was allowed to react with Me₃SiOTf-BCl₃ whereas GC analysis of the reaction mixture after removal of BCl₃ indicated the presence of higher boiling isomeric products, they were found to be disproportionation products (diisopropylbenzene isomers based on GC-MS analysis). The only silicon-containing compound identified was Me₃SiCl formed by ligand exchange with the Lewis acid, i.e., the same processes observed in our NMR studies of the complex. Control experiments showed that under the reaction condition, neither BCl₃ nor Me₃SiOTf itself were able to disproportionate cumene. The catalyst for disproportionation is therefore the conjugate acid formed by the reaction between BCl₃ and triflic acid impurity.⁶ Similarly, no silylated aromatics were found in reaction with toluene or anisole. The latter gave a 10% yield of isomeric methylphenols. In subsequent experiments 2,4,6-tri-*tert*-butylpyridine was used as a proton trap in order to suppress acid-catalyzed desilylation that would have taken place. Once again, however, silylated aromatics were not detected.

In summary as shown by ²⁹Si and ¹¹B NMR data we have obtained polarized donor-acceptor complexes from a series of silyl triflates with boron trihalides. No observable silicenium ions were formed in the systems, although triflate to halide ligand exchange indicates intermediate S_Ni-like ionization or S_N2-like exchange through five coordinated silicon species.

Experimental Section

Trimethylsilyl triflate (Petrach) and dimethyl-*tert*-butylsilyl triflate (Aldrich) were commercial samples of sufficient purity and were used as received. Diethylsilyl ditriflate was prepared from Et₂SiHCl and triflic acid using the method recently described by Corey¹⁰ for the preparation of *i*-Pr₂Si(OTf)₂.

Triethylsilyl triflate was prepared in ca. 90% yield (²⁹Si NMR) by an overnight reaction of Et₃SiCl and triflic acid (1 equiv) under dry nitrogen, followed by the removal of any dissolved HCl in solution under vacuum.

Tri-*n*-butylsilyl triflate and dimethylisopropylsilyl triflate were prepared from their corresponding chlorides and triflic acid (1 equiv) by room temperature mixing under argon for 18 h. ²⁹Si NMR spectrum of the reaction mixture showed a complete reaction, and no unreacted chloride could be detected.

(8) Wilberg, E.; Kruerke, U. Z. *Naturforsch., B: Anorg. Chem., Org. Chem., Biochem., Biophys., Biol.* 1953, 8B, 608.

(9) Wrachmeyer, B. *Prog. Nucl. Magn. Reson. Spectrosc.* 1979, 12, 227.

(10) Corey, E. J.; Hopkins, P. B. *Tetrahedron Lett.* 1982, 23, 4871.

Triisopropylsilyl triflate and diphenylmethylsilyl triflate were similarly prepared from their corresponding chlorides.

Tris(alkylthio)silyl chloride was prepared in ca. 95% yield (estimated by ^{29}Si NMR) from SiCl_4 and *i*-PrSH (3 equiv) using the method described in the literature for the preparation of tris(alkylthio)silane from trichlorosilane.¹¹ The reaction of the chloride with 1 equiv of triflic acid gave a ca. 65% yield of the triflate (estimated by ^{29}Si NMR). Unreacted triflic acid in the reaction mixture was removed by addition of the required amount of 2,4,6-tri-*tert*-butylpyridine. An aliquot of the solution was used for NMR studies (transferred to an NMR tube under nitrogen).

BBr_3 was doubly distilled, followed by distillation from aluminum powder, in an all glass distillation apparatus, until it was completely colorless. It was then stored in a drybox. BCl_3 (Matheson) was used as received. All aromatic compounds were commercial samples of highest purity and were used as received.

The ^{29}Si and ^{11}B NMR spectra were recorded on a Varian FT-80A instrument. The silicon chemical shifts were referenced to Me_4Si , whereas the boron shifts were referenced to $\text{BF}_3 \cdot \text{Et}_2\text{O}$. GC analyses were performed on a Varian Model 3700 gas chromatography using a capillary as well as a short packed column (OV 101).

Complexation of Silyl Triflates with BX_3 . The triflate (1 mL) was placed in an NMR tube and was cooled in dry ice/acetone under dry nitrogen with exclusion of moisture. After slow addition of BCl_3 (ca. 2 mL) and vortex mixing, the sample was first studied by ^{29}Si NMR directly in BCl_3 solvent, followed by ^{11}B NMR analysis usually after removal of excess BCl_3 and SO_2Cl_2 as solvent. With BBr_3 , whereas the addition was initially done at low temperature, the NMR tube was allowed to warm up slowly

to room temperature prior to NMR studies. Same procedure was followed for the reaction of Me_3SiOR with BCl_3 and BBr_3 .

For reactions with aromatics, the donor-acceptor complexes formed upon reaction of Me_3SiOTf or Et_3SiOTf with BCl_3 at low temperature were added to an excess of the aromatic compound precooled to ca. -40°C with efficient magnetic stirring under dry nitrogen. The temperature was then allowed to warm up slowly to room temperature, followed by an additional 30 min of mixing at room temperature. The remaining BCl_3 was then removed by passing a fast flow of nitrogen through the reaction mixture, and the mixture was analyzed by GLC. In some cases, the reaction mixtures were also subjected to aqueous workup and ether extraction prior to GLC analysis.

Acknowledgment. Support of our work by the National Science Foundation is gratefully acknowledged.

Registry No. $\text{Me}_3\text{SiOTf} \cdot \text{BCl}_3$, 91158-42-8; $\text{Me}_3\text{SiOTf} \cdot \text{BBr}_3$, 91158-43-9; $\text{Et}_3\text{SiOTf} \cdot \text{BCl}_3$, 91158-44-0; *n*- $\text{Bu}_3\text{SiOTf} \cdot \text{BCl}_3$, 91158-45-1; $\text{Me}_2(i\text{-Pr})\text{SiOTf} \cdot \text{BCl}_3$, 91158-46-2; $\text{Ph}_2(\text{Me})\text{SiOTf} \cdot \text{BBr}_3$, 91158-47-3; $\text{Ph}_2(\text{Me})\text{SiOTf} \cdot \text{BCl}_3$, 91158-48-4; Me_3SiCl , 75-77-4; Me_3SiBr , 2857-97-8; Et_3SiCl , 994-30-9; *i*- Pr_3SiCl , 13154-24-0; *n*- Bu_3SiCl , 995-45-9; $\text{Me}_2(i\text{-Pr})\text{SiCl}$, 3634-56-8; $\text{Me}_2(t\text{-Bu})\text{SiBr}$, 76358-45-7; $\text{Ph}_2(\text{Me})\text{SiBr}$, 17571-61-8; $\text{B}(\text{OTf})\text{Cl}_2$, 91158-32-6; $\text{B}(\text{OTf})\text{Br}_2$, 91158-33-7; Me_3SiOTf , 27607-77-8; Et_3SiOTf , 79271-56-0; $\text{Et}_2\text{Si}(\text{OTf})_2$, 91158-34-8; *i*- Pr_3SiOTf , 80522-42-5; *n*- Bu_3SiOTf , 91158-35-9; $\text{Me}_2(i\text{-Pr})\text{SiOTf}$, 91158-36-0; $\text{Me}_2(t\text{-Bu})\text{SiOTf}$, 69739-34-0; $\text{Ph}_2(\text{Me})\text{SiOTf}$, 91158-37-1; BCl_3 , 10294-34-5; BBr_3 , 10294-33-4; AlBr_3 , 7727-15-3; AlCl_3 , 7446-70-0; Me_3SiOMe , 1825-61-2; Me_3SiOEt , 1825-62-3; Me_3SiOPh , 1529-17-5; MeOBBR_2 , 29877-99-4; EtOBBR_2 , 91158-38-2; PhOBBR_2 , 91158-39-3; MeOBCl_2 , 867-46-9; EtOBCl_2 , 16339-28-9; PhOBCl_2 , 75088-70-9; cumene, 98-82-8; tris(isopropylthio)silyl triflate, 91158-40-6; chlorotris(isopropylthio)silane, 91158-41-7.

(11) Wolinski, L.; Tiechemann, H.; Post, H. W. *J. Org. Chem.* 1951, 16, 395.

Thermolysis of 6-Oxa-3-silabicyclo[3.1.0]hexanes: A New Convenient Route to Silicon-Oxygen π -Bonded Species

Georges Manuel* and Guy Bertrand

Laboratoire des Organométalliques, Université Paul-Sabatier, 31062 Toulouse Cedex, France

William P. Weber* and Samih A. Kazoura

Loker Hydrocarbon Research Institute, Department of Chemistry, University of Southern California, Los Angeles, California 90089-1661

Received December 21, 1983

The direct generation of silanones and germanones from easily prepared 6-oxa-3-metallabicyclo[3.1.0]hexanes is reported. Application of this method to the transient formation of an $[\text{O}=\text{Si}=\text{O}]$ species is given. This silicon analogue of carbon dioxide was trapped by hexamethylcyclotrisiloxane to yield 2,2,4,4,6,6,10,10,12,12,14,14-dodecamethyl-2,4,6,8,10,12,14-heptasila-1,3,5,7,9,11,13,15-octaexaspiro[7.7]-pentadecane, the expected spiro adduct, in 60% yield.

Silanones, compounds containing a silicon-oxygen double bond, have been postulated as transient intermediates in a variety of reactions.¹ Most of these reactions involve the conversion of other unsaturated silicon intermediates into silanones. For example, reaction of non-enolizable aldehydes or ketones with silenes,²⁻⁶ silanimines,^{7,8}

or disilenes⁹⁻¹¹ yield silanones. Likewise, reaction of silylenes with sulfoxides give silanones.¹²⁻¹⁷ On the other

(1) For a general review of π -bonded silicon see: L. E. Gusel'nikov and N. S. Nametkin, *Chem. Rev.*, 79, 529 (1979).

(2) T. J. Barton and E. A. Kline, Third International Symposium on Organosilicon Chemistry, Madison, WI, Aug 1972.

(3) W. Ando, A. Sekiguchi, and T. Migita, *J. Am. Chem. Soc.*, 97, 7159 (1975).

(4) D. N. Roark and L. H. Sommer, *J. Chem. Soc., Chem. Commun.*, 167 (1973).

(5) P. B. Valkovich and W. P. Weber, *J. Organomet. Chem.*, 99, 231 (1975).

(6) I. M. T. Davidson, C. E. Dean, and F. T. Lawrence, *J. Chem. Soc., Chem. Commun.*, 52 (1981).

(7) C. M. Golino, R. D. Bush, and L. H. Sommer, *J. Am. Chem. Soc.*, 96, 614 (1974).

(8) D. R. Parker and L. H. Sommer, *J. Organomet. Chem.*, 110, C1 (1976).

(9) T. J. Barton and J. A. Kilgour, *J. Am. Chem. Soc.*, 96, 2278 (1974).