

for $K = \pm 3$, while much narrower widths (≤ 5 kHz) are predicted for $K = \pm 1$. For K states other than 0, ± 3 , and ± 6 to be forbidden, the symmetric top must have a 3-fold axis associated with three zero-spin nuclei.¹¹ These criteria are met by the molecular cluster $^{40}\text{Ar}_3\text{-H/DF}$, which is an oblate symmetric top.

Confirmation of this composition is provided by analysis of B_0 . Neglecting any tilt of the Ar_3 plane and vibrations other than torsional oscillations of the H/DF , we have¹³

$$I_{\text{bb}}(\text{cluster}) = (1/2)(1 + \cos^2\alpha)I_{\text{bb}}(\text{H/DF}) + I_{\text{bb}}(\text{Ar}_3) + \mu_c R^2 \quad (1)$$

where R is the distance between the H/DF and Ar_3 centers of mass (c.m.) and $I_{\text{bb}}(\text{Ar}_3)$ is a function of r , the Ar-Ar distance. For argon dimer, a B_0 of 0.05778 cm^{-1} has been determined from a high-resolution Raman study.¹⁴ It gives an r_0 of 3.821 \AA , which in eq 1 leads to values of 2.785 and 2.479 \AA for R in $\text{Ar}_3\text{-HF}$ and -DF . The smaller R for the DF species show that the H/D end is pointed at the Ar_3 . The corresponding H/DF c.m. to Ar distances of 3.553 and 3.525 \AA are comparable with the 3.510 and 3.461 \AA found in Ar-H/DF .² Also, the H/DF c.m. to Ar line is 38.5° from the c axis. A full account will be presented later, including the results of a search for the isotopic species $^{36}\text{Ar}^{40}\text{Ar}_2\text{-H/DF}$ to determine accurate bond distances.

Registry No. Ar, 7440-37-1; HF, 7664-39-3; DF, 14333-26-7.

(13) Read, W. G.; Campbell, E. J.; Henderson, G. *J. Chem. Phys.* **1983**, *78*, 3501.

(14) Godfried, H. F.; Silvera, I. F. *Phys. Rev. Lett.* **1982**, *48*, 1337 and prior work cited therein.

Synthesis, Trapping, and Spectral Characterization of 1H-Cyclopropa[1]phenanthrene¹


Brian Halton,*^{2a} Barry R. Dent,^{2a} Stefan Böhm,^{2b}
David L. Officer,^{2a,b} Hans Schmickler,^{2b} Fritz Schopphoff,^{2b}
and Emanuel Vogel*^{2b}

Department of Chemistry
Victoria University of Wellington
Wellington, New Zealand
Institut für Organische Chemie der Universität Köln
5 Köln 41, West Germany


Received June 3, 1985

The cycloproparenes **1**,³ **2**,⁴ and **3**,⁵ and their derivatives are surprisingly stable, readily available materials.⁶ By comparison cyclopropa[*a*]naphthalene (**4**) explodes upon melting^{7a} and its 1,1-difluoro derivative^{7b} is stable in solution only at temperatures below -30°C . This decrease in stability perhaps reflects the formally enhanced olefinic character of the bridge bond. 1H-Cyclopropa[1]phenanthrene (**5**), the most important unknown structural variant within this family, should exemplify further this

Scheme I



Scheme II



treatment with potassium *tert*-butoxide, probably¹² via the ylide **10**. The primary hydrocarbon products **5** and **11** cannot be isolated despite many variations in the reaction conditions. However, with added furan they are trapped as cycloadducts **12** (20%),^{13,14} **13** (13%),^{13,15} and **14** (7%)¹⁶ (Scheme I). The symmetry planes in **12** and **13** are evident from the appearance of only 10 resonances in their ¹³C NMR spectra^{14,15} and the orientation of the three-membered ring with respect to the oxygen bridge is confidently assigned from the deshielding influence the oxygen atom has upon the ¹H NMR shift of H12a in **12**. Thus **12** displays a one-proton doublet ($J = 4.6$ Hz) at 2.55 ppm for H12a, whereas the equivalent resonance in **13** (1.97 ppm, $J = 5.2$ Hz) is more shielded. It is interesting that the major isolable products emanate from **5**, formed by abstraction of the benzylic H9b of **10**. The appearance of **14**¹⁶ as a minor product reflects the loss of H1 from **10** to give **11**, which is known¹⁶ to add stereospecifically to furan to give **14**.

A complementary approach to **5**, free of competing elimination modes, employs retrodiene cleavage and parallels the successful syntheses of **13a** and **4**.^{7a} Thus vacuum pyrolysis of **16**,¹⁷ the dicyanoacetylene adduct^{7a} of **15**,¹⁸ followed by low-temperature manipulation of the product mixture, has provided a dibromodifluoromethane solution of **5**¹⁹ free of *o*-dicyanobenzene (**17**) (Scheme II). The NMR spectral data recorded at -60 °C are in full accord with **5**. The C1 methylene protons appear as a singlet (3.54 ppm) in the expected⁶ range for the cycloproparenes and the aromatic proton multiplets typify the phenanthrene system. The single high-field carbon resonance (23.6 ppm, C1) has the highest chemical shift (by 4 ppm) yet recorded for C1 of a C1-unsubstituted cycloproparene, but the magnitude of the C1-H coupling (170.9 Hz) is normal.⁶ Seven distinct aromatic carbon resonances are noted but the shielding of the ortho carbons C1b(9a) is reduced from the usual⁶ 14-16 ppm to ~6 ppm. Prepared in this way, **5** reacts with furan to give **12** (28%) and **13** (17%) and with α -pyrone to yield tetraene **15** (41%) (after loss of carbon dioxide below 0 °C) (Scheme II). On prolonged standing the solution of **5** decomposes even at -60 °C. However, **5** has been isolated as a moderately stable solid which decomposes slowly over a period of days at -78 °C.

Acknowledgment. Financial assistance from the New Zealand Universities Grants Committee (to B.R.D.) and an Alexander von Humboldt fellowship (to D.L.O.; 1984-1985) are acknowledged gratefully.

Registry No. **5**, 278-91-1; **8**, 98420-55-4; **9**, 98420-57-6; **10**, 98420-58-7; **11**, 57741-39-6; **12**, 98420-59-8; **13**, 98523-81-0; **14**, 53137-63-6; **15**, 98420-60-1; **16**, 98420-61-2; **17**, 91-15-6; furan, 110-00-9; dicyanoacetylene, 1071-98-3; α -pyrone, 504-31-4.


Dewar Furan: Its Generation and Trapping with Isobenzofuran

Ian G. Pitt, Richard A. Russell,¹ and Ronald N. Warrener*

Chemistry Department, The Faculties
Australian National University
Canberra, A.C.T. 2601, Australia

Received June 25, 1985

It is now more than 20 years since van Tamelen and Pappas announced, in a landmark paper at that time, the preparation of the parent Dewar valence isomer **1** of benzene.² In contrast, the



heterocyclic Dewar valence isomers **2-4**³ remained unknown in unsubstituted form until 1985, when Strausz and his co-workers reported the trapping of Dewar thiophene (**3**), albeit in very low yield.⁴ Photoisomerization of the appropriate five-membered heterocycle has formed the major thrust for preparing such valence isomers and has been applied successfully to the generation of substituted Dewar thiophenes⁵ and Dewar pyrroles.⁶ However, irradiation of furans, while producing a variety of rearrangement products, offers no evidence for Dewar furans (5-oxabicyclo-[2.1.0]pent-2-enes).⁷ Indeed the few known examples,^{8,9} e.g., the perfluoromethyl derivative **5** reported by Lemal,⁸ have been prepared by indirect means.

Thus, in seeking a route to Dewar furan we designed the photochemical precursor **13** which is based on the established success of the 1,2-photoaromatization reaction to produce highly strained four-membered cyclic olefins.¹⁰ In this paper we discuss the results of this approach and announce the first example of the generation of Dewar furan **4** and its efficient trapping in adduct form with isobenzofuran (**15**).¹¹

The photosubstrate **13** was prepared as outlined in Scheme I.¹²

(12) Halazy, S.; Krief, A. *Tetrahedron Lett.* **1979**, 4233-4236.

(13) Compounds **12** and **13** may be named as *syn*- and *anti*-3,4:5,6-dibenzo-11-oxatetracyclo[6.2.1.1^{2,7}.0^{2,7}]dodeca-3,5,9-triene, respectively.

(14) **12**: mp 179.5-180.5 °C. Anal. (C₁₉H₁₄O) C, H. ¹H NMR δ 0.58 (d, $J = 4.6$ Hz, H12b), 2.55 (d, $J = 4.6$ Hz, H12a), 5.26 (s, H1,8), 6.34 (s, H9,10), 7.20-7.30 (m, H14,15,18,19), 7.45-7.60 (m, H13,20), 7.85-7.95 (m, H16,17); ¹³C NMR δ 25.6 (t, C12), 35.6 (d, C2,7), 79.4 (d, C1,8), 123.6, 126.7, 127.9, 128.9 (all d, C13-20), 130.5 (s, C3,6), 134.5 (s, C4,5), 139.4 (d, C9,10).

(15) **13**: mp 183-185 °C. Anal. (C₁₉H₁₄O) C, H. ¹H NMR δ 0.77 (d, $J = 5.2$ Hz, H12b), 1.97 (d, $J = 5.2$ Hz, H12a), 5.36 (s, H1,8), 6.45 (s, H9,10), 7.25-7.35 (m, H14,15,18,19), 7.50-7.60 (m, H13,20), 7.95-8.05 (m, H16,17); ¹³C NMR δ 31.9 (s, C2,7), 35.2 (t, C12), 84.8 (d, C1,8), 123.5, 126.6, 128.0, 129.0 (all d, C13-20), 130.5 (s, C3,6), 133.1 (d, C9,10), 134.8 (s, C4,5).

(16) Compound **14** provides analytical and spectroscopic data in accord with those previously published: Coburn, T. T.; Jones, W. M. *J. Am. Chem. Soc.* **1974**, 96, 5218-5227.

(17) Compound **16** may be named as 3,4:5,6-dibenzo-9,10-dicyano-tetracyclo[6.2.2.1^{2,7}.0^{2,7}]trideca-3,5,9,11-tetraene: 67%, mp 214-215 °C. Anal. (C₂₃H₁₄N₂) C, H, N. ¹H NMR δ 0.32 (d, $J = 5.7$ Hz, H13b), 1.99 (d, $J = 5.7$ Hz, H13a), 4.80 (t, $J = 3.9$ Hz, H1,8), 6.38 (t, $J = 3.9$ Hz, H1,12), 7.30-7.45 (m, H15,16,19,20), 7.70-7.80 (m, H14,21), 7.90-8.00 (m, H17,18); ¹³C NMR δ 26.6 (t, C13), 32.6 (s, C2,7), 47.0 (d, C1,8), 114.1 (s, CN), 123.3, 127.5, 128.1, 129.0, 131.1 (all d, C11,12,14-21), 127.0, 133.4, 137.4 (all s, C3-6,9,10).

(18) Compound **15** is available from triphenylene in seven steps: Tokita, S.; Schopphoff, F.; Vogel, E. manuscript in preparation.

(19) 1*H*-Cyclopropa[*l*]phenanthrene (**5**): ¹H NMR δ 3.54 (s, H1), 7.68-7.80 (m, H3,4,7,8), 7.95-8.00 (m, H2,9), 8.75-8.85 (m, H5,6); ¹³C NMR δ 23.6 (t, $J_{CH} = 170.9$ Hz, C1), 124.1, 124.6 (both s, C1a,9b and C1b,9a), 125.3, 125.8, 127.4, 128.6 (all d, C2-9), 133.5 (s, C5a,5b).

(1) Visiting Fellow ANU.

(2) van Tamelen, E. E.; Pappas, S. P. *J. Am. Chem. Soc.* **1963**, 85, 3297-3298.

(3) Kobayashi, Y.; Kumadaki, I. *Adv. Heterocycl. Chem.* **1982**, 31, 169-206. Kobayashi, Y.; Kumadaki, I. *Top. Curr. Chem.* **1984**, 123, 103-149.

(4) Rendall, W. A.; Torres, M.; Strausz, O. P. *J. Am. Chem. Soc.* **1985**, 107, 723-724.

(5) Weibe, H. A.; Braslavsky, S.; Heicklen, J. *Can. J. Chem.* **1972**, 50, 2721-2724. Kobayashi, Y.; Kumadaki, I.; Ohsawa, A.; Sekine, Y.; Mochizuki, H. *Chem. Pharm. Bull.* **1975**, 23, 2773-2778. Kobayashi, Y.; Kawada, K.; Ando, A.; Kumadaki, I. *Heterocycles* **1983**, 20, 174. Kobayashi, Y.; Kawada, K.; Ando, A.; Kumadaki, I. *Tetrahedron Lett.* **1984**, 25, 1917-1920. Barltrop, J. A.; Day, A. C.; Irving, E. *J. Chem. Soc., Chem. Commun.* **1979**, 881-883.

(6) Barltrop, J. A.; Day, A. C.; Ward, R. R. *J. Chem. Soc., Chem. Commun.* **1978**, 131-133. Hiraoka, H. *Ibid.* **1971**, 1610-1611.

(7) Padwa, A. "Rearrangements in Ground and Excited States"; De Mayo, P., Ed.; Academic Press: New York, 1980; Vol. 3. Boué, S.; Srinivasan, R. *J. Am. Chem. Soc.* **1970**, 92, 1824-1828. van Tamelen, E. E.; Whitesides, T. H. *Ibid.* **1971**, 93, 6129-6140. Barton, T. J.; Hussman, G. P. *Ibid.* **1983**, 105, 6316-6318.

(8) Wirth, D.; Lemal, D. M. *J. Am. Chem. Soc.* **1982**, 104, 847-848.

(9) Warrener, R. N.; Pitt, I. G.; Russell, R. A. *J. Chem. Soc., Chem. Commun.* **1984**, 1464-1466. Pitt, I. G.; Russell, R. A.; Warrener, R. N. *Ibid.* **1984**, 1466-1467.

(10) Warrener, R. N.; Nunn, E. E.; Paddon-Row, M. N. *Aust. J. Chem.* **1979**, 32, 2650-2674. Warrener, R. N.; Kretschmer, G.; Paddon-Row, M. N. *J. Chem. Soc., Chem. Commun.* **1977**, 806-807. Nunn, E. E.; Warrener, R. N. *Ibid.* **1972**, 818-819. Kretschmer, G.; Warrener, R. N. *Tetrahedron Lett.* **1975**, 1335-1338.

(11) Warrener, R. N. *J. Am. Chem. Soc.* **1971**, 93, 2346-2348.