

for $\text{Cr}(\text{CO})_3(\text{C}_6\text{H}_6)_3$, favoring the conclusions we arrived at in a previous paper.³

We thank Prof. G. Natta and Prof. R. Ercoli for their helpful suggestions.

(3) P. Corradini and G. Allegra, *THIS JOURNAL*, **81**, 5510 (1959).

DEPARTMENT OF INDUSTRIAL CHEMISTRY
POLYTECHNIC INSTITUTE OF MILAN

PAOLO CORRADINI
GIUSEPPE ALLEGRA


RECEIVED FEBRUARY 25, 1960

THE REACTION OF DICHLORODIPHENYLSILANE WITH SODIUM

Sir:

One of the reactions examined by Kipping and co-workers in their extensive studies with organosilicon compounds was that of dichlorodiphenylsilane with sodium.¹ When the reaction was carried out above the melting point of sodium without solvent, or with toluene or xylene as solvent, a complex mixture of chlorine-free organosilicon compounds resulted. This mixture was resolved into six components after a painstaking series of operations. One of these designated as Compound (A) first was reported not to melt at 300° and later was reported to melt at 335°. This sparingly soluble compound was crystallized from benzene and gave satisfactory analyses for a compound having four diphenylsilylene groups, $[(\text{C}_6\text{H}_5)_2\text{Si}]_4$. Several possible structures were considered, and of these the one given preference on the basis of apparent unsaturated characteristics of Compound (A) was that containing "two trivalent silicon atoms": $-\text{SiPh}_2\cdot\text{SiPh}_2\cdot\text{SiPh}_2\cdot\text{Ph}_2\text{Si}-$.

Incidental to a study concerned with cyclic organosilicon compounds, we have examined Compound (A), and have found that it does not contain any trivalent silicon atoms and that the high reactivity or "unsaturation" characteristics may be explained by a cyclic structure




This octaphenylcyclotetrasilane undergoes ready cleavage by lithium in tetrahydrofuran to give a novel organosilicon compound containing two silyllithium groups. The di-silyllithium compound reacts with trimethyl phosphate to give 1,4-dimethyloctaphenyltetrasilane which melts at 223° after crystallization from a 1:1 mixture of benzene and petroleum ether (b.p. 50–60°).

Anal. Calcd. for $\text{C}_{50}\text{H}_{46}\text{Si}_4$: C, 79.09; H, 6.11; Si, 14.8. Found: C, 79.16 and 79.11; H, 6.41 and 6.24; Si, 14.85 and 14.72. Hydrogen value (moist piperidine): calcd., 101; found, 103, 104.

The same dimethyl compound was obtained from either 1,4-dichlorooctaphenyltetrasilane² or 1,4-

dibromooctaphenyltetrasilane,² and methylolithium. Also, and most significantly, the dimethyl compound was obtained by the reaction of 1,2-dichlorotetraphenyldisilane and methyldiphenylsilyllithium, two compounds of unequivocal structure.



The exclusion of the biradical or trivalent silicon structure was confirmed by the measurements of J. H. Chaudet and Dr. W. D. Kennedy. The compound when examined by electro paramagnetic resonance both in the solid state and as a saturated solution in toluene failed to give a free radical resonance. The sensitivity for the detection of free radicals by the electro paramagnetic resonance is of the order of 10^{15} unpaired electrons per gram of compound. It should also be stated that the octaphenylcyclotetrasilane lacked sufficient solubility for a nuclear magnetic resonance examination.

Acknowledgments.—The authors are grateful to J. H. Chaudet, and Dr. W. D. Kennedy, of the Tennessee Eastman Co., for the electron paramagnetic resonance studies. This research was supported in part by the United States Air Force under contract AF 33(616)-3510 monitored by Materials Laboratory, Directorate of Laboratories, Wright Air Development Center, Wright-Patterson AFB, Ohio.

CHEMISTRY DEPARTMENT
IOWA STATE UNIVERSITY
AMES, IOWA

H. GILMAN
D. J. PETERSON
A. W. JARVIE
H. J. S. WINKLER

RECEIVED MARCH 2, 1960

PRIMARY PHOSPHINE OXIDES

Sir:

Tertiary phosphine oxides were prepared early in the history of organophosphorus chemistry and are now accessible by a number of methods.¹ Oxides of secondary phosphines were discovered much later, and general methods of preparation have come to light only recently.^{2–4} We wish to report here the first preparations of primary phosphine oxides.

These substances were observed initially as products of the reaction of phosphine with ketones. Our work indicates that this apparently general reaction takes the course indicated, which involves transfer of oxygen from carbon to phosphorus in the first step.

(1) G. M. Kosolapoff, "Organophosphorus Compounds," John Wiley and Sons, Inc., New York, N. Y., 1950, p. 98.

(2) R. H. Williams and L. A. Hamilton, *THIS JOURNAL*, **74**, 5418 (1952).

(3) R. H. Williams and L. A. Hamilton, *ibid.*, **77**, 3411 (1955).

(4) M. M. Rauhut, I. Hechenbleikner, H. A. Currier and V. P. Wystrach, *ibid.*, **80**, 6690 (1958).

(1) F. S. Kipping and J. E. Sands, *J. Chem. Soc.*, **119**, 830 and 848 (1921); F. S. Kipping, *ibid.*, **123**, 2590 and 2598 (1923); F. S. Kipping, *ibid.*, **125**, 2291 (1924); F. S. Kipping, 2719 and 2728 (1927); F. S. Kipping and H. E. Murray, *ibid.*, 360 (1929); A. R. Steele and F. S. Kipping, *ibid.*, 2545 (1929); and F. S. Kipping, "The Bakerian Lecture," *Proc. Roy. Soc.*, **159A**, 139 (1937).

(2) The dichloro and dibromo compounds were prepared from compound (A) and tetrachloroethane and ethylene bromide, respectively. A later report will describe these and a wide variety of other reactions of the octaphenylcyclotetrasilane as well as a corrected formula for "Compound B" formed from it.