

Eigenenergies of 3D-Coulomb and 3D-Harmonic Oscillator Potentials from WKB Quantization: Point Canonical Transformation

Hosung Sun

Department of Chemistry, Sungkyunkwan University, Suwon 440-746, Korea. E-mail: hsun@skku.edu

Received September 28, 2007

A direct application of the WKB quantization to the three-dimensional Coulomb potential does not yield the exact eigenenergies. The three-dimensional Coulomb potential is converted to a Morse potential by using the point canonical transformation. Then the WKB quantization is applied to the Morse potential to find a relationship between the eigenenergies of the Coulomb and those of the Morse potentials. From the relationship the exact eigenenergies of the Coulomb potential are determined. The same method is found to be also valid for the three-dimensional harmonic oscillator potential. And the Langer modified WKB quantization is algebraically derived.

Key Words : WKB quantization, Coulomb potential, Harmonic oscillator potential

Background

The Schrödinger equation for one-dimensional system is (in units of $2m = \hbar = 1$)

$$\left[-\frac{d^2}{dr^2} + V(r) \right] \Psi_n(r) = E_n \Psi_n(r). \quad (1)$$

Here we consider a potential $V(r)$ that varies smoothly, *i.e.*, there exist two classical turning points for all bound states. The lowest order Wentzel-Kramers-Brillouin approximation (or the WKB quantization condition) for the above system is^{1,2}

$$\int_{r_L}^{r_R} \sqrt{E_n - V(r)} dr = \left(n + \frac{1}{2} \right) \pi \quad n = 0, 1, 2, \dots \quad (2)$$

where r_L and r_R are inner (short distance) and outer (long distance) turning point, respectively, *i.e.*, $V(r_L) = V(r_R) = E_n$. The WKB quantization is a semiclassical method of obtaining approximate eigenenergies in the limiting case of large quantum numbers or high energy. But it is found to be exact for the one-dimensional harmonic oscillator potential and the one-dimensional Morse potential.²⁻⁴ For other potentials, for example, the 3D-Coulomb potential or the 3D-harmonic oscillator potential, the WKB quantization does not reproduce the exact eigenenergies.

Sukhatme *et al.* reported that all solvable potentials are inter-related to each other by a certain transformation.^{5,6} They classify the solvable potentials by type-I and type-II. The type-II category includes three potentials, *i.e.*, the one-dimensional Morse ($V^M(r)$), the 3D-Coulomb ($V^C(r)$) and the 3D-harmonic oscillator ($V^H(r)$) potentials. And the three potentials can be transformed to each other by the so-called point canonical transformation.

In this work we briefly introduce a new method of determining the exact eigenenergies of $V^C(r)$ from the point canonical transformation and the WKB quantization. As mentioned before, the exact eigenenergies of $V^M(r)$ can be

obtained from the WKB quantization but the eigenenergies of $V^C(r)$ are not. First, we transform the Schrödinger equation with $V^C(r)$ into another Schrödinger equation with $V^M(r)$. And then we apply the WKB quantization to the $V^M(r)$ to obtain a relationship between the eigenenergies of $V^C(r)$ and those of $V^M(r)$. Since the exact eigenenergies of $V^M(r)$ are known, the eigenenergies of $V^C(r)$ can be determined from the relationship. The above argument is also valid for the 3D-harmonic oscillator potential $V^H(r)$.

We define the 3D-Coulomb and the 3D-harmonic oscillator potentials as follows.⁴⁻⁶ For a given angular momentum quantum number l , the 3D-Coulomb potential ($q =$ charge parameter) is

$$V(r) = V^C(r) = -\frac{q^2}{r} + \frac{l(l+1)}{r^2} \quad (0 < r < \infty) \quad (3)$$

and the eigenenergies $E_n^C = -\frac{q^4}{4(n+l+1)^2}$ for $n = 0, 1, 2, \dots$ (4)

Since $V^C(r)$ goes to zero as $r \rightarrow \infty$, the bound state eigenenergies E_n^C are negative.

For a given l , the 3D-harmonic oscillator potential ($\omega =$ frequency parameter) is

$$V(r) = V^H(r) = \frac{1}{4} \omega^2 r^2 + \frac{l(l+1)}{r^2} \quad (0 < r < \infty) \quad (5)$$

and the eigenenergies $E_n^H = \left(2n + l + \frac{3}{2} \right) \omega$ for $n = 0, 1, 2, \dots$ (6)

For reference, the one-dimensional Morse potential ($A, B, \alpha =$ parameters) is

$$V^M(x) = B^2 e^{-2\alpha x} - 2B \left(A + \frac{\alpha}{2} \right) e^{-\alpha x} \quad (-\infty < x < \infty). \quad (7)$$

The phase-space integral in the WKB quantization can be analytically evaluated as⁴

$$\int_{x_L}^{x_R} \sqrt{E_n^M - V^M(x)} dx = \frac{\pi}{\alpha} \left(A + \frac{\alpha}{2} - \sqrt{-E_n^M} \right) \quad (8)$$

where $V^M(x_L) = V^M(x_R) = E_n^M$. Since $V^M(x)$ goes to zero as $x \rightarrow \infty$, the eigenenergies E_n^M are negative.

Canonical Transformation and WKB for 3D-Coulomb Potential

For the 3D-Coulomb potential, the Schrödinger equation is

$$\left[-\frac{d^2}{dr^2} - \frac{q^2}{r} + \frac{l(l+1)}{r^2} \right] \Psi_n^C(r) = E_n^C \Psi_n^C(r) \quad (0 < r < \infty) \quad (9)$$

with the boundary condition of $\Psi_n^C(r) = 0$ as $r \rightarrow 0$ and $r \rightarrow \infty$. Canonically transforming, *i.e.*, changing the variable r to $x = -\ln r$ and the wave function $\Psi_n^C(r)$ to $\Phi_n(x) = e^{x/2} \Psi_n^C(r)$, one obtains

$$\left[-\frac{d^2}{dx^2} - E_n^C e^{-2x} - q^2 e^{-x} + \left(l + \frac{1}{2} \right)^2 \right] \Phi_n(x) = 0 \quad (-\infty < x < \infty) \quad (10)$$

The new function $\Phi_n(x)$ satisfies the boundary condition of $\Phi_n(x) = 0$ as $x \rightarrow -\infty$ and $x \rightarrow \infty$. The transformation is sometimes called the Langer transformation.⁷⁻⁹ Note that the canonical transformation converts Eq. (9) on the half-line $(0, \infty)$ to Eq. (10) on the full-line $(-\infty, \infty)$.

Once E_n^C and $\Phi_n(x)$ are given, Eq. (10) can be considered as a Schrödinger equation for a Morse potential

$$V^M(x) = -E_n^C e^{-2x} - q^2 e^{-x} \quad \text{with} \quad E_n^M = -\left(l + \frac{1}{2} \right)^2, \quad \text{i.e.,}$$

$$\left[-\frac{d^2}{dx^2} + V^M(x) \right] \Phi_n(x) = E_n^M \Phi_n(x). \quad (11)$$

Eq. (11) looks awkward because the potential contains the n -dependent E_n^C term. But, in fact, the potential is not a function of n because the potential is defined for a fixed n , *i.e.*, Eq. (11) is defined for each n . Note that the eigenvalue E_n^M is the same for all n . Therefore Eq. (11), as a whole, is a Schrödinger equation with the eigenfunction $\Phi_n(x)$ and the eigenenergy E_n^M .¹⁰

One can evaluate E_n^M in Eq. (11) using the WKB quantization in Eq. (2) that is exact for Morse potentials,

$$\int_{x_L}^{x_R} \sqrt{E_n^M - (-E_n^C e^{-2x} - q^2 e^{-x})} dx = \left(n + \frac{1}{2} \right) \pi \quad (12)$$

where $V^M(x_L) = V^M(x_R) = E_n^M$. (In Appendix we explicitly show why the WKB quantization is exact for Morse potentials but not exact for 3D-Coulomb potentials or for 3D-harmonic oscillator potentials.) Comparing the potential in Eq. (12) with that in Eq. (7), one immediately finds that $\alpha = 1$, $B = \sqrt{-E_n^C}$, and $A = q^2/2\sqrt{-E_n^C} - 1/2$. From the phase-space integral expression in Eq. (8) and the WKB quantization in Eq. (12), one obtains

$$\pi \left(\frac{q^2}{2\sqrt{-E_n^C}} - \sqrt{-E_n^M} \right) = \left(n + \frac{1}{2} \right) \pi \quad \text{for } n = 0, 1, 2, \dots \quad (13)$$

And the Morse eigenenergy E_n^M is

$$E_n^M = -\left(\frac{q^2}{2\sqrt{-E_n^C}} - n - \frac{1}{2} \right)^2. \quad (14)$$

Eq. (14) shows the relationship between the eigenenergies of the original 3D-Coulomb potential $-q^2/r + l(l+1)/r^2$ and those of the transformed Morse potential $-E_n^C e^{-2x} - q^2 e^{-x}$. Since $E_n^M = -(l+1/2)^2$, Eq. (14) gives the eigenenergies of the 3D-Coulomb potential $E_n^C = -q^4/4(n+l+1)^2$ ($n = 0, 1, 2, \dots$) that is identical with that in Eq. (4). Indeed the new method that utilizes the point canonical transformation and the WKB quantization reproduces the exact eigenenergies of the 3D-Coulomb potential.

To make the WKB quantization exact for the 3D-coulomb potential, modified versions of WKB quantization have been suggested. The earliest work was the Langer modification where the centrifugal term $l(l+1)/r^2$ in Eq. (19) is replaced by $(l+1/2)^2/r^2$.⁷ Though various modifications or improvements of the WKB quantization, since the Langer's work, have been studied, they all include the term $(l+1/2)^2/r^2$ (or some operator related to it) in their modification.^{8,9,11}

The Langer modification has its own physical origin, but using our transformation technique we can justify the existence of the modification algebraically. Inserting $E_n^M = -(l+1/2)^2$ into the WKB quantization in Eq. (12), one obtains

$$\int_{x_L}^{x_R} \sqrt{-\left(l + \frac{1}{2} \right)^2 + E_n^C e^{-2x} + q^2 e^{-x}} dx = \left(n + \frac{1}{2} \right) \pi. \quad (15)$$

Changing the variable x back to r using the previous $x = -\ln r$, one obtains

$$\int_{r_L}^{r_R} \sqrt{E_n^C - \left\{ -\frac{q^2}{r} + \frac{(l+1/2)^2}{r^2} \right\}} dr = \left(n + \frac{1}{2} \right) \pi. \quad (16)$$

Eq. (16) is none other than the Langer modified WKB quantization that can exactly reproduce the eigenenergies of the 3D-Coulomb potential. We would like to stress that our transformation technique naturally introduces the Langer modification without imposing any semiclassical approximation.

Example of 3D-Harmonic Oscillator Potential

The method proposed in the previous section can be applied to any potential that can be transformed to a Morse potential. In this section, taking the 3D-harmonic oscillator potential as another example, we repeatedly show how the method works.

For the 3D-harmonic oscillator potential, the Schrödinger equation is

$$\left[-\frac{d^2}{dr^2} + \frac{1}{4}\omega^2 r^2 + \frac{l(l+1)}{r^2} \right] \Psi_n^H(r) = E_n^H \Psi_n^H(r) \quad (-\infty < x < \infty). \quad (17)$$

Changing the variable r to $x = -2\ln r$ and the wave function $\Psi_n^H(r)$ to $\Phi_n(x) = e^{x/4} \Psi_n^H(r)$, one obtains

$$\left[-\frac{d^2}{dx^2} + \frac{\omega^2}{16} e^{-2x} - \frac{E_n^H}{4} e^{-x} + \frac{1}{4} \left(l + \frac{1}{2} \right)^2 \right] \Phi_n(x) = 0. \quad (18)$$

Eq. (18) can be considered as a Schrödinger equation for a Morse potential $V^M(x) = (\omega^2/16)e^{-2x} - (E_n^H/4)e^{-x}$ with eigenenergy $E_n^M = -(1/4)(l+1/2)^2$.

The WKB quantization for Eq. (18) is

$$\int_{x_L}^{x_R} \sqrt{E_n^M - \left(\frac{\omega^2}{16} e^{-2x} - \frac{E_n^H}{4} e^{-x} \right)} dx = \left(n + \frac{1}{2} \right) \pi. \quad (19)$$

Comparing the Morse potential in Eq. (19) with that in Eq. (7), one finds that $\alpha = 1$, $B = \omega/4$, and $A = E_n^M/2\omega - 1/2$. From Eq. (8) and Eq. (19), one obtains

$$\pi \left(\frac{E_n^H}{2\omega} - \sqrt{-E_n^M} \right) = \left(n + \frac{1}{2} \right) \pi \quad \text{for } n = 0, 1, 2, \dots \quad (20)$$

Then the Morse eigenenergy E_n^M is

$$E_n^M = - \left(\frac{E_n^H}{2\omega} - n - \frac{1}{2} \right)^2. \quad (21)$$

Since $E_n^M = -(1/4)(l+1/2)^2$, the eigenenergies of the 3D-harmonic oscillator potential is $E_n^H = (2n+l+3/2)\omega$ ($n = 0, 1, 2, \dots$) that is identical with the exact eigenenergies in Eq. (6). Again we are able to derive the exact eigenenergies of the 3D-harmonic oscillator potential using the point canonical transformation and the WKB quantization.

Following the same way for the 3D-Coulomb potential, the Langer modified WKB quantization is derived as

$$\int_{r_L}^{r_R} \sqrt{E_n^H - \left\{ \frac{1}{4}\omega^2 r^2 + \frac{(l+1/2)^2}{r^2} \right\}} dr = \left(n + \frac{1}{2} \right) \pi. \quad (22)$$

The Langer modification for the 3D-harmonic oscillator potential turns out to be the same as that for the 3D-Coulomb potential. Actually it is the same for all solvable spherically symmetric potentials.^{9,12}

Conclusion

The point canonical transformation has been introduced to explain the inter-relation among the exactly solvable potentials.^{5,6} This transformation has been utilized to explain the supersymmetry^{6,13} of certain potentials.¹⁰ In this work we have found another interesting usage of the transformation.

Using the point canonical transformation, the Schrödinger equation for the 3D-Coulomb potential is converted to another Schrödinger equation with a Morse potential. Then

an application of the WKB quantization to the Morse potential yields a relationship between the eigenenergies of the Coulomb and those of the Morse potentials. From the relationship the exact eigenenergies of the Coulomb potential are derived. The Langer modified WKB quantization is also algebraically derived. The proposed method is valid for any potential that can be transformed to a Morse potential. As another example, the exact eigenenergies of the 3D-harmonic oscillator potential are determined by using the same method.

Other solvable potentials belonging to the type-I category (e.g., the Eckart potential, the Scarf potential, and the Rosen-Morse potential, etc.) are related to the type-II potentials (e.g., the Morse potential, etc.) through redefinition of parameters and the so-called limiting procedure.⁵ Therefore the type-I potentials are easily transformed to a Morse potential and, consequently, our method can be again used to determine the exact eigenenergies of the type-I potentials. As a matter of fact, all the type-I and type-II potentials have a common feature of Natanzon potentials that are exactly solvable.

One cannot say that the WKB quantization is exact for all solvable potentials. However it is not appropriate for one to assert that the WKB quantization is useless when one desires exact eigenenergies of solvable potentials other than the Morse potential (or one-dimensional harmonic oscillator potential.) As shown in this work, the WKB quantization may be useful even in low energy regime where semi-classical approximation is not valid.

Appendix

Another way of modifying the WKB quantization is to introduce the nonintegral Maslov index μ .^{2,14} In this scheme the exact quantization condition is written as

$$\int_{r_L}^{r_R} \sqrt{E_n - V(r)} dr = \left(n + \frac{\mu}{4} \right) \pi. \quad (A1)$$

When the Maslov index $\mu = 2$, Eq. (A1) is, of course, reduced to the WKB quantization. There is no general way of evaluating the Maslov index for an arbitrary potential. However it can be evaluated for exactly solvable potentials. The direct method of evaluating the Maslov index is to adopt an exact quantization, for example, the analytical transfer matrix method (ATMM).

The ATMM quantization condition can be summarized as¹⁵

$$\int_{r_L}^{r_R} \sqrt{E_n - V(r)} dr + \delta_n = (n+1)\pi. \quad (A2)$$

And the so-called scattering-led phase shift δ_n is

$$\delta_n = - \int_{r_L}^{r_R} k_n'(r) \frac{P_n(r)}{P_n'(r)} dr \quad (A3)$$

where $k_n'(r) = \frac{d[\sqrt{E_n - V(r)}]}{dr}$ and $P_n'(r) = \frac{dP_n(r)}{dr}$.

$P_n(r)$ is the minus log derivative of the n -th state wave function $\Psi_n(r)$, i.e., $P_n(r) = (d\Psi_n(r)/dr)/\Psi_n(r)$. Comparing Eq. (A1) with Eq. (A2), one immediately finds that the Maslov index $\mu = 4(1 - \delta_n/\pi)$.

For the general form of Morse potential given in Eq. (7) including the forms in Eqs. (12) and (19), we have found $\delta_n = 1/2\pi$ in our former work.¹⁶ Consequently the Maslov index μ is 2. It proves that

the WKB quantization should be valid for the Morse potential.

In our previous paper¹⁷ in this Journal we have presented $\delta_n = (\sqrt{l(l+1)} - l)\pi$ for the 3D-Coulomb potential in Eq. (3). Therefore the Maslov index μ is $4[l+1 - \sqrt{l(l+1)}]$. Using the same method suggested in the previous paper we algebraically evaluate the phase shift and the Maslov index for the 3D-harmonic oscillator potential Eq. (5) for the first time. And the result is that $\delta_n = [(\sqrt{l(l+1)} - l)/2 + 1/4]\pi$ and $\mu = 3 - 2\sqrt{l(l+1)} + 2l$. (In Ref. 14, based on qualitative arguments, the same Maslov index has been deduced.) It shows that the WKB quantization is not valid for the 3D-Coulomb potential nor for the 3D-harmonic oscillator potential since μ is not 2. However, this work extends the validity of the WKB quantization to the 3D-Coulomb and 3D-harmonic oscillator potentials through the point canonical transformation.

References

1. Fröman, N.; Fröman, P. O. *JWKB Approximation*; North Holland: Amsterdam, 1965.
 2. Friedrich, H.; Trost, J. *Phys. Rep.* **2004**, 397, 359.
 3. Ivanov, I. A. *J. Phys. A: Math. Gen.* **1997**, 30, 3977.
 4. Hruška, M.; Keung, W.-Y.; Sukhatme, U. *Phys. Rev. A* **1997**, 55, 3345.
 5. Gangopadhyaya, A.; Panigrahi, P. K.; Sukhatme, U. P. *Helv. Phys. Acta* **1994**, 67, 363.
 6. Cooper, F.; Khare, A.; Sukhatme, U. P. *Phys. Rep.* **1995**, 251, 267.
 7. Langer, R. E. *Phys. Rev.* **1937**, 51, 669.
 8. Krieger, J. B.; Rosenzweig, C. *Phys. Rev.* **1967**, 164, 171.
 9. Sergeenko, M. N. *Phys. Rev. A* **1996**, 53, 3798.
 10. Haymaker, R. W.; Rau, A. R. P. *Am. J. Phys.* **1986**, 54, 928.
 11. Bruev, A. S. *Phys. Lett. A* **1992**, 161, 407.
 12. Ou, Y.; Cao, Z.; Shen, Q. *Phys. Lett. A* **2003**, 318, 36.
 13. Sun, H. *Bull. Korean Chem. Soc.* **2006**, 27, 515.
 14. Friedrich, H.; Trost, J. *Phys. Rev. A* **1996**, 54, 1136.
 15. Ou, Y. C.; Cao, Z.; Shen, Q. *J. Chem. Phys.* **2004**, 121, 8175.
 16. Sun, H. *Phys. Lett. A* **2005**, 338, 309.
 17. Sun, H. *Bull. Korean Chem. Soc.* **2007**, 28, 408.
-