

trast with an increase observed, in the $|\rho_{YZ}|$ values. This can be rationalized with the enhanced contribution of resonance "shunt" by the α -CO as the charge transfer increases, which has a shortening effect on the C_α - C_β bond length due to the double bond character in structure (III); the greater degree of charge transfer (the larger ρ_Y) will bring the greater contribution of resonance shunt, which in turn will result in a shorter C_α - C_β bond. The shorter C_α - C_β bond will naturally mean a larger $|\rho_{YZ}|$ value as observed; hence the parallel increase in the $|\rho_{YZ}|$ with ρ_Y and ρ_Z can be accommodated.

In structure (IV), there is an unpaired electron on C_α so that the substituent Y can interact with the C_β atom, which in turn means the interaction between X and Y; there will be no decrease in the interaction (and hence in ρ_{XY}) between X and Y. Thus our results clearly support involvement of structure III, but not (IV), in the TS of the nucleophilic substitution reaction of α -carbonyl compounds. We can conclude that the involvement of the bridged TS structure is a tenuous one in the light of our results of cross interaction constants.

Acknowledgement. We thank the Korea Science and Engineering Foundation and the Ministry of Education for support of this work. One of us (C. S. Shim) also thanks Dae Woo foundation for a postgraduate research grant.

References

- (a) S. Wolfe and D. Kost, *Nouv. J. Chim.*, **2**, 441 (1978); (b) D. Kost and K. Aviram, *Tetrahedron Lett.*, **23**, 4157 (1982); (c) S. Wolfe, D. J. Mitchell and H. B. Schlegel, *Can. J. Chem.*, **60**, 1291 (1982); (d) S. S. Shaik, *J. Am. Chem. Soc.*, **105**, 4359 (1983).
- (a) J. B. Conant and W. R. Kirner, *J. Am. Chem. Soc.*, **46**, 233 (1924); (b) J. B. Conant, W. R. Kirner and R. E. Hussey, *ibid.*, **47**, 488 (1925); (c) A. Halvorsen and J. Songstad, *J. Chem. Soc. Chem. Commun.*, 327 (1978); (d) J. W. Thorpe and J. Warkentin, *Can. J. Chem.*, **51**, 927 (1973); (e) F. G. Bordwell and W. T. Brannen, *J. Am. Chem. Soc.*, **86**, 4645 (1964).
- D. J. McLennan and A. Pross, *J. Chem. Soc., Perkin 2*, 981 (1984).
- M. J. S. Dewar, "The Electronic Theory of Organic Chemistry", Oxford Univ. Press, Oxford, p.73 (1949).
- (a) S. Winstein, E. Grunwald and H. W. Jones, *J. Am. Chem. Soc.*, **73**, 2700 (1951); (b) A. Streitwieser, "Solvolytic Displacement Reactions", McGraw-Hill, New York, p.28 (1962).
- (a) I. Lee and S. C. Sohn, *J. Chem. Soc. Chem. Commun.*, 1055 (1986); (b) I. Lee, submitted for publication; (c) I. Lee, H. Y. Kim and H. K. Kang, Submitted for publication; (d) I. Lee and H. K. Kang, *Tetrahedron Lett.*, In press.
- J. Shorter, "Correlation Analysis of Organic Reactivity", Research studies Press, Chichester, Chap. 2 (1982).
- I. Lee and K. R. Han, Unpublished results.
- Calculated from; J. Banger, A. E. Cockrill and G. L. O. Davies, *J. Chem. Soc. (B)*, 498 (1971).
- The ρ_{XZ} value for reactions of substituted (X) N,N-dimethylanilines with Benzylarenesulfonates (Z) in acetone at 35°C was calculated to be $\rho_{XZ} = -0.15$ from H. Yamataka and T. Ando, *tetrahedron Lett.*, **23**, 4805 (1982).
- Substituted (X) anilines reacting with various substrate (with Y) gave the following ρ_{XY} values:

$$Y\phi CH_2SO_2Cl \xrightarrow[35.0^\circ C]{MeOH} \rho_{XY} = -0.69^{12}; \quad Y\phi SO_2Cl$$

$$\xrightarrow[35.0^\circ C]{MeOH} \rho_{XY} = -0.70^{13}; \quad Y\phi CH_2Cl \xrightarrow[50^\circ C]{EtOH} \rho_{XY} = -0.77^{14};$$

$$Y\phi CH_2OS\phi_2 \xrightarrow[35.0^\circ C]{MeOH} \rho_{XY} = -0.62^{6a}; \quad Y\phi CH_2SO_2F$$

$$\xrightarrow[45.0^\circ C]{MeOH} \rho_{XY} = -1.24^{6e}.$$
- H. K. Kang, Ph. D. thesis, Inha University, (1987).
- Calculated from: I. Lee and I. S. Koo, *Tetrahedron*, **39**, 1803 (1983).
- Calculated from: F. P. Ballistreri, E. Maccarone and A. Mamo, *J. Org. Chem.*, **41**, 3364 (1976).
- (a) G. M. Loudin and C. Berke, *J. Am. Chem. Soc.*, **96**, 4508 (1974). (b) J. P. Richard, M. E. Rothenberg and W. P. Jencks, *ibid.*, 106, 1361 (1984).

A Convenient Synthesis of (E)- β -Farnesene

Suk-Ku Kang*, Gyung-Yun Chung, and Dong-Ha Lee

Department of Chemistry, Sung Kyun Kwan University Natural Science Campus, Suwon 170.

Received April 21, 1987

(E)- β -Farnesene(**1**), (6E)-7,11-dimethyl-3-methylene-1,6,10-dodecatriene (Fig. 1) was known as a constituent of various essential oils¹ and recently used for enhancing the aroma of perfumed materials, e.g. soaps, detergents, and magnolia fragrance². In 1972, W.S. Bowers identified (E)- β -farnesene as an alarm pheromone of several economically important species of aphids³. The damage to crops by aphids either by feeding or by transmitting virus disease can be prevented to some extent by insecticides, but aphids are

developing increased resistance, particularly to systemic organophosphates⁴. Also, such insecticides do not prevent damage to crops by non-persistent viruses, such as potato virus Y, since these are rapidly acquired and transmitted by aphids. The fast-acting pyrethroids may control infection with non-persistent viruses⁵ but are less effective with the more resistant biotypes *Myzus persicae*, so alternative novel approaches appear desirable. Of the chemicals that influence aphid behavior, the most active is the alarm pheromone⁶

which, for most important pest species, is the sesquiterpene hydrocarbon, (E)- β -Farnesene (**1**). Synthetic (E)- β -Farnesene substantially increases the effectiveness of contact pesticides such as permethrin against aphids in the laboratory and this approach is being investigated in field crops⁷. (E)- β -Farnesene was used as an insecticide for aphid control. It was reported that (E)- β -Farnesene (10-20 μ g/nm²) on radish leaves killed *Myzus persicae* and *Neomyzus circumflexus*⁸.





Figure 1

Several methods were reported for the dehydration⁹ of farnesol and nerolidol with both acidic² and basic catalyst to synthesize (E)- β -farnesene. By dehydration, mixtures including α - and β -farnesenes were formed. Even if some improved dehydration reactions^{9c,9f} were reported, none are universally applicable and in fact, often give poor yields or unmanageable mixtures. The multistep specific conversion of (E,E)-farnesol into β -farnesene was known¹⁰. Recently Otera *et al*¹¹ reported the conversion of farnesol THP ether into the conjugated diene **1**. Several other syntheses^{12,13,14,15} have been reported in the literature. Three groups¹² have reported the synthesis of (E)- β -farnesene by the establishment of the (E)-configuration at C-6 position and the β -substituted dienic moiety from C-1 to C-3 position from readily available starting materials. Two syntheses¹³ involved the construction of the butadienyl moiety. Stille's synthesis¹⁴ depended on the cross-coupling of geranyl bromide which has (E)-configuration at C-6 position with (2-methylene-3-butenyl) trimethyltin in the presence of zinc chloride. Otsuka *et al*¹⁵ described the synthesis of β -farnesene by isoprene oligomerization.

Here we wish to report a convenient and practical method to synthesize (E)- β -farnesene by regioselective 1,4-elimination reaction of farnesyl chloride, which in turn was readily prepared from farnesol.

Farnesyl chloride (**3**) was easily prepared¹⁶ from farnesol (**2**) by treatment with thionyl chloride and pyridine. Exposure of farnesyl chloride (**3**) to BuOK in DMSO (Method A) or in THF in the presence of 18-crown-6 (Method B) at 40°C afforded diene **1** in β -form exclusively. No other isomers were detectable. t-BuOK attacks the methylhydrogen preferentially rather than the methylene hydrogen. This explains why the β -isomers are formed exclusively from allylic chloride **3**. The addition of 18-crown-6 accelerated the rate.



^a (a) SOCl₂, pyr., 0°C, 88%; (b) KOtBu, DMSO, 40°C, 62%; (c) KOtBu, THF, 18-crown-6, 40°C, 87%

Scheme 1

Typical procedure is as follows. Method A: To a stirred solution of farnesyl chloride (0.25g, 1.0 mmol) was added

KOtBu (0.30g, 2.7 mmol) in DMSO (5.0 ml). The mixture was stirred at 40°C for 14 hr. To the reaction mixture ether (30 ml) was added and washed with water and saturated NaCl solution. The organic layer was dried over MgSO₄ and then concentrated in vacuo. The crude product was separated by preparative thin layer chromatography using n-hexane: ethyl acetate (9:1) (R_f = 0.72) as eluents to afford (E)- β -farnesene¹⁷ (0.13g, 62%). The (E)- β -farnesene thus obtained was identical in all respects (TLC, IR, ¹H- and ¹³C-NMR) with the material reported in the literature.^{9-15,18} Method B: To a stirred solution of farnesyl chloride (0.25g, 1.0 mmol) was added 18-crown-6-ether (0.53g, 2.0 mmol) followed by KOtBu (1.1g, 10 mmol) in dry THF (5.0 ml). The mixture was stirred at 40°C for 12hr. To the reaction mixture ether (30 ml) was added and washed with water and saturated NaCl solution. The organic layer was dried over MgSO₄ and then concentrated in vacuo. The crude product was separated by preparative thin layer chromatography using n-hexane: ethyl acetate (9:1) (R_f = 0.72) as eluents to afford β -farnesene (0.18g, 87%).

The above synthetic method described is considered to be convenient and economic to prepare a fair amount of β -farnesene, the alarm pheromone of aphids to conduct field test experiments¹⁹.

Acknowledgement. This research was supported by a grant (1985-1987) from the Korea Science and Engineering Foundation.

References

- (a) "The Merck Index 10th Edition, M. Windholz, Ed., Merck & Co. Inc., Rahway, N. J., 1983, p 568." (b) P. E. Shaw, "Citrus Science and Technology", Vol 1; S. Nagy, P. E. Shaw, M. K. Veldhuis, Ari Publishing Co., Westport, CT 1977, Chapter 11; (c) F. Sorm and J. Arient, *Coll. Czech. Chem. Soc.*, **15**, 1975 (1950); (d) F. Sorm, *et al.*, *ibid*, **14**, 274 (1949).
- B. D. Mookherjee, R. W. Trenkel, B. J. Chant, O. A. Van, V. Kamath, and C. J. Mussinan, *Fr. Demande FR 2,511, 244* (1983) (CA: 98, 221627K).
- (a) W. S. Bowers, L. R. Nault, R. E. Webb, and S. R. Dutky, *Science*, **177**, 1121 (1972); (b) L. J. Edwards, J. B. Siddall, L. L. Dunham, P. Uden, and C. J. Kislow, *Nature* (London), **241**, 126 (1973); (c) W. H. Wientjens, A. C. Lakwijk, and T. Van der Marel, *Experientia*, **29**, 658 (1973) (CA: **79**, 40112a); (d) W. S. Bowers, C. Nishino, M. E. Montgomery, and L. R. Nault, *J. Insect Physiol.*, **33**, 697 (1977) (CA: **87**, 181142n); (e) J. A. Pickett and D. C. Griffiths, *J. Chem. Ecol.*, **6**, 349 (1980); (f) C. Nishino, W. S. Bowers, M. E. Montgomery, and L. R. Nault, *Appl. Ent. Zool.*, **11**, 340 (1976).
- R. M. Sawick, A. L. Devonshire, A. D. Rice, G. D. Moores, S. M. Petzing, and A. Cameron, *Pest. Sci.*, **9**, 189 (1978).
- R. W. Gibson, A. D. Rice, and R. M. Sawick, *Ann. Appl. Biol.*, **100**, 49 (1982) (CA: **97**, 122033m).
- J. Miyamoto and P. C. Kearney "Pesticide Chemistry: Human Welfare and Environment", Vol 2; N. Takahashi, H. Yoshioka, T. Misato, and S. Matsunaka, Pergamon press, 1983, pp 117.
- (a) D. C. Griffiths and J. A. Pickett, *J. Chem. Ecol.*, **6**, 349 (1980) (CA: **93**, 90177 K); (b) J. A. Pickett, G. W.

- Dawson, R. W. Gibson, D. C. Griffiths, A. D. Rice, R. M. Sawacki, M. C. Smith, and C. M. Woodcock, *Les Colloques del' I.N.R.A.*, **7**, 243 (1982) (CA: **98**, 102622V).
8. J. Gut, O. Van, and M. Adriaan, *Eur. Pat. App. EP* 73,080 (1983) (CA: 193390g).
9. (a) Y. Naves, *Helv. Chim. Acta.*, **49**, 1029 (1966); (b) G. Brieger, *J. Org. Chem.*, **32**, 3270 (1967); (c) G. W. Dawson, D. C. Griffiths, J. A. Pickett, M. C. Smith, and C. M. Woodcock, *J. Chem. Ecol.*, **8**, 1111 (1982); (d) G. Brieger, S. W. Watson, D. G. Barar, and A. L. Shene, *J. Org. Chem.*, **44**, 1340 (1979); (e) A. Bhati, *Perfumery Essent. Oil Record*, **53**, 376 (1962); (f) G. Brieger, T. J. Nestrick, and C. Mc Kenna, *J. Org. Chem.*, **34**, 3789 (1969).
10. S. Tanaka, A. Yasuda, H. Yamamoto, and H. Nozaki, *J. Am. Chem. Soc.*, **37**, 3252 (1975).
11. J. Otera, Y. Niibo, and K. Okuda, *Chemistry Letters*, 1829 (1986).
12. (a) O. P. Vig, A. K. Vig, and S. D. Kumar, *Indian J. Chem.*, **13**, 1003 (1975); (b) T. Mandai, K. Kawada, and J. Otera, *J. Org. Chem.*, **48**, 5183 (1983); (c) S-K Kang, and S-G Kim, *Bull. Korean Chem. Soc.*, **7**, 157 (1986).
13. (a) B. Cazes, E. Guittet, S. Julia, and O. Ruel, *J. Organomet. Chem.*, **177**, 67 (1979); (b) A. Carpita, F. Bonaccorsi, and R. Rossi, *Tetrahedron Letters*, 5193 (1984).
14. J. P. Godschalx and J. K. Stille, *Tetrahedron Letters*, 1905 (1983).
15. S. Akutagawa, T. Taketomi, and S. Otsuka, *Chemistry Letters*, 485 (1976).
16. (a) W. G. Young, *J. Org. Chem.*, 6164 (1960); (b) S. Akutagawa, T. Taketomi, H. Kumobayashi, K. Takayama, T. Someya, and S. Otsuka, *Bull. Chem. Soc. Japan*, **51**, 1158 (1978).
17. Satisfactory physical properties and spectroscopic data ($^1\text{H-NMR}$, $^{13}\text{C-NMR}$, IR) were obtained for the synthetic compound **1**: bp99°C/3 mmHg; IR (NaCl, neat) 3095, 2970, 2930, 1592, 1440, 1372, 990, 900, 890, 830 cm^{-1} ; $^1\text{H-NMR}$ (80 MHz, CDCl_3) 1.60 (s, 6H), 1.69 (s, 3H), 1.90-2.22 (m, 8H), 4.45-5.38 (m, 6H), 6.36 (dd, 1H, $J=17\text{Hz}$ and 11Hz); $^{13}\text{C-NMR}$ (80 MHz, CDCl_3) 115.67, 139.10, 146.26, 31.56, 26.71, 124.13, 135.46, 39.80, 26.82, 124.46, 131.39, 25.73, 17.69, 16.06, 113.05 (from $\text{C}_1\text{-C}_{15}$).
18. B. V. Burger, M. le Roux, H. S. C. Spies, V. Truter, and R. C. Bigalke, *Tetrahedron Letters*, 5221 (1978).
19. In the literature, escape response of pea aphids *A. pisum* to β -farnesene has been reported: P. Wohlers, *Entomol. Exp. Appl.*, **27**, 156 (1980) (CA: **93**, 20710e). The escape response test of the synthetic alarm pheromone for aphids *Lipaphis erysimi* K. and *Myzus persicae* S. was conducted by Department of Entomology, Institute of Agricultural Science, O. R. D., Suwon 170, Korea: S-K. Kang, G-Y. Chung, J-O. Lee, and H. G. Goh, *Sung Kyun Kwan Univ. Journal* (Natural Science), **37**, 141 (1986).