

## Stoichiometric Solvation Effects. Solvolysis of Methanesulfonyl Chloride

In Sun Koo,\* Kiyull Yang, Sun Kyoung An, Chong-Kwang Lee,<sup>†</sup> and Ikchoon Lee<sup>‡</sup>

Department of Chemical Education, Gyeongsang National University, Chinju 660-701, Korea

<sup>†</sup>Department of Chemistry, Gyeongsang National University, Chinju 660-701, Korea

<sup>‡</sup>Department of Chemistry, Inha University, Incheon 402-751, Korea

Received July 21, 2000

Solvolyses of methanesulfonyl chloride in water, D<sub>2</sub>O, CH<sub>3</sub>OD, and in aqueous binary mixtures of acetone, ethanol and methanol are investigated at 25, 35 and 45 °C. The Grunwald-Winstein plot of first-order rate constants for the solvolytic reaction of methanesulfonyl chloride with Y<sub>Cl</sub> (based on 2-adamantyl chloride) shows marked dispersions into three separate lines for three aqueous mixtures with a small *m* value (*m* < 0.30), and shows a rate maximum for aqueous alcoholic solvents. Stoichiometric third-order rate constants, *k*<sub>ww</sub> and *k*<sub>aa</sub> were calculated from the observed first-order rate constants and (*k*<sub>aw</sub> + *k*<sub>wa</sub>) was calculated from the *k*<sub>ww</sub> and *k*<sub>aa</sub> values. The kinetic solvent isotope effects determined in water and methanol are consistent with the proposed mechanism of the general base catalyzed and/or S<sub>A</sub>N/S<sub>N</sub>2 reaction mechanism for methanesulfonyl chloride solvolyses based on mass law and stoichiometric solvation effect studies.

### Introduction

Sulfonyl halides are known to solvolyze by a borderline mechanism<sup>1</sup> as in the solvolysis of benzyl chloride.<sup>2</sup> There have been, however, much dispute over the mechanism as to whether it is an S<sub>N</sub>2<sup>3</sup> or an S<sub>A</sub>N<sup>4</sup> process, the former being preferred lately. A particularly important system which contains tetracoordinate sulfur is RSO<sub>2</sub>Cl; sulfonyl chlorides are important reagents in organic synthesis and substitution reactions of these compounds bridge inorganic and organic chemistry.

Solvent effects and linear free energy relationships in the solvolyses of sulfonyl halides, especially substituted benzenesulfonyl chlorides, have received much experimental attention,<sup>5</sup> but little work has been done on the solvent stoichiometric effect on alcohol-water mixed solvents, especially for the solvolyses of aliphatic sulfonyl chlorides.

In order to examine the quantitative solvent effects, it is necessary to take into account the stoichiometric solvation effects based on a third order reaction mechanism. For solvolyses in alcohol-water mixtures, interpretations based on a third order reaction mechanism are more complex, but significant new information is available because two products are formed. Thus there are four possible third order rate constants in the solvolysis reaction, *k*<sub>aa</sub>, *k*<sub>aw</sub>, *k*<sub>wa</sub>, and *k*<sub>ww</sub>.<sup>6,9</sup>

In this study, we determined the pseudo first order rate constants for solvolyses of methanesulfonyl chloride (MSC) in alcohol-water mixtures, and estimated third order constants, *k*<sub>ww</sub>, (*k*<sub>wa</sub> + *k*<sub>aw</sub>) and *k*<sub>aa</sub>. We will discuss the mechanism of solvolytic reaction of MSC using third order rate constants and kinetic solvent isotope effects. We will also compare results of solvolytic reaction to that of molecular orbital calculation for the solvolytic reaction of MSC.<sup>10</sup>

### Result and Discussion

The rate constants for solvolyses of MSC in methanol-

water, ethanol-water, and acetone-water at 25, 35 and 45 °C are summarized in Table 1. Reference to the table reveals that the rate increases in the order acetone-water < ethanol-water < methanol-water at three temperature, and it shows maximum rate constants at near 40% methanol-water (40M) mixtures and at near 30% ethanol-water (30E) mixtures. The rate increases slowly as the water content of the mixtures increases; this means that the rate is slightly accelerated by the solvent with higher ionizing power, *Y*, suggesting that bond breaking in the transition state is of little importance. First-order rate constants for solvolysis of MSC vary over only a fourteen-fold range in alcohol-water mixtures, whereas the observed first-order rate constants for benzyl chloride, para-methoxybenzoyl chloride and thenoyl chloride which are known to react *via* an S<sub>N</sub>2 or S<sub>N</sub>1 reaction mechanism varying over several thousand-fold range.<sup>11</sup>

These results indicate that the rate determining step is not the bond breaking step, which was found in the reaction of

**Table 1.** Solvolyses of methanesulfonyl chloride in aqueous alcohol and acetone binary solvent mixtures at 25, 35 and 45 °C

v/v %	MeOH			EtOH			Acetone		
	<i>k</i> × 10 <sup>4</sup>								
	25 °C	35 °C	45 °C	25 °C	35 °C	45 °C	25 °C	35 °C	45 °C
100	0.267	0.737	1.74	0.149	0.281	0.642			
90	0.695	1.78	4.57	0.403	0.997	2.45			
80	1.14	2.92	7.08	0.673	1.68	3.92	0.199	0.406	0.847
70	1.66	4.04	9.73	0.922	2.29	5.53	0.418	1.00	1.77
60	2.09	5.31	12.5	1.23	3.06	7.29	0.630	1.44	3.34
50	2.33	6.20	14.7	1.61	3.88	9.34	1.05	2.25	5.36
40	2.65	6.74	16.8	2.03	4.87	12.0	1.31	3.33	7.94
30	2.52	6.99	17.7	2.33	5.88	14.5	1.76	4.36	10.6
20	2.36	6.84	17.4	2.37	6.21	15.8	2.16	5.29	13.0
10	2.23	6.32	16.8	2.18	6.12	16.4	2.17	5.79	15.1
0	2.04	5.96	16.4	2.04	5.96	16.4	2.04	5.96	16.4


$S_N1$  or  $S_N2$  substrate, but the addition step for the reaction of  $S_N$  substrate where transition state is not sensitive to the solvent ionizing power. These results are very similar to those of solvolytic reactions of *p*-nitrobenzoyl chloride,<sup>12</sup> *p*-nitrobenzenesulfonyl chloride,<sup>13</sup> furoyl chloride,<sup>11a</sup> and phenyl chloroformates,<sup>14</sup> but they are different from the results of solvolytic reactions of benzyl chloride, para-methoxybenzoyl chloride and thenoyl chloride.<sup>11a,11c</sup>

The Grunwald-Winstein plots (Eq. 1) of the rates in Table 1 are presented in Figure 1 using the solvent ionizing power scale  $Y_{Cl}$ , based on 1-adamantyl chloride.<sup>15</sup> Investigation of Figure 1 shows the plots for the three separate curves with maximum rate constants at near 40M and near 30E. The slopes ( $m$ ) for the range of relatively good linearity are very small values of  $m = 0.26$  ( $r = 0.995$ ),  $m = 0.28$  ( $r = 0.996$ ), and  $m = 0.27$  ( $r = 0.990$ ) for the ranges of 100M ~60M, 100E ~50E, and 80A ~20A, respectively, implying that the solvolysis of MSC in the binary mixtures proceeds by addition-elimination ( $S_N$ ) or associative  $S_N2$  channel.


$$\log(k/k_0)_{RX} = mY_{Cl} \quad (1)$$

Logarithm of third-order rate constants for solvolysis of MSC in acetone-water mixtures ( $\log k_{ww} = \log k_{obs}/[H_2O]^2$ ) versus  $Y_{Cl}$  plots are presented in Figure 2. The slope of Figure 2 show a very small  $m$  value of near zero (-0.036), implying that the  $k_{ww}$  terms of solvolysis of MSC in the binary mixtures is relatively independent of medium. These results indicate that the reaction proceeds through the transition state which is not sensitive to the solvent ionizing power again.<sup>12a</sup>

In order to examine the non-linear Grunwald-Winstein plot, it is necessary to take into account the stoichiometric solvation effects based on third order rate constants. For corresponding solvolyses in alcohol-water mixtures, interpretations based on a third order mechanism are more complex, but significant new information is available because two



**Figure 1.** Logarithms of first-order rate constants for solvolyses of methanesulfonyl chloride at 25 °C  $\log(k/k_0)$  vs.  $Y_{Cl}$  (solvent codes: ●; methanol, ○; ethanol, ▲; acetone).



**Figure 2.** Logarithms of calculated third-order rate constants for hydrolysis of methanesulfonyl chloride in acetone/water at 25 °C vs.  $Y_{Cl}$  (slope  $m = -0.036$ , intercept = -6.824, correlation coefficient = 0.963).

products are formed. Thus there are four possible third order rate constants in the solvolysis reaction: (i)  $k_{aa}$  for a mechanism in which one molecule of alcohol acts as a nucleophile and second molecule acts as a general base; (ii)  $k_{aw}$  in which alcohol acts as a nucleophile and water acts as a general base; (iii)  $k_{wa}$  in which water acts as nucleophile and alcohol acts as general base; (iv)  $k_{ww}$  in which water acts as both nucleophile and general base.<sup>6-9</sup> Therefore, observed first-order rate constants in alcohol-water mixtures are given by Eq. (2).


$$k_{obs} = k_{aa}[\text{alcohol}]^2 + (k_{aw} + k_{wa})[\text{alcohol}][\text{water}] + k_{ww}[\text{water}]^2 \quad (2)$$

The  $k_{aa}$  term can be calculated from observed first order rate constants in pure alcohol ( $k_{aa} = k_{obs}/[ROH]^2$ ); similarly  $k_{ww}$  can be obtained from the observed first order rate constant in water ( $k_{ww} = k_{obs}/[H_2O]^2$ ).<sup>7,9</sup>

It is impossible to separate  $k_{aw}$  and  $k_{wa}$  from  $k_{obs}$ , but  $(k_{aw} + k_{wa})$  term can be calculated from  $k_{obs} - (k_{ww}[H_2O]^2 + k_{aa}[ROH]^2)$ .

The third-order rate constants are summarized in Table 2 for solvolysis of MSC in aqueous methanol and ethanol. Reference to Table 2 reveals that the  $k_{ww}$  terms increase with increasing of water content of the mixtures, but the  $k_{aa}$  terms decrease. The contributions of third-order terms,  $(k_{wa} + k_{aw})$  are larger than  $k_{ww}$  and  $k_{aa}$  terms, and show a maximum at near 40M and 30E in aqueous alcohol mixtures. Such a maximum rate behaviour often observed<sup>13b,14a,b,c</sup> in the solvolysis of alcohol mixtures, and this due to exceptionally greater contribution of  $(k_{wa} + k_{aw})$  term, i.e., alcohol is acting as a nucleophile and water is acting as general base catalysis or *vice versa*.

Recently, we reported *ab initio* calculation on the hydrolysis and solvolysis of MSC by water methanol, and methanol-water mixtures. In that work, the solvent-catalyzed hydroly-



Scheme 1


sis and methanolysis of MSC with a trigonal bipyramidal transition state in solvent was suggested (Scheme 1).<sup>10</sup>

These results are in good agreement with the general base catalysis  $S_{AN}/S_N2^{17}$  reaction mechanism proceeding through a tight transition state, where bond formation is much more progressed than bond breaking in the transition state.

A plot of logarithm of rate constants *versus* logarithm of water concentration, which is often referred to as Kivinen plot,<sup>18</sup> is shown in Figure 3 for three aqueous mixtures. The slopes ( $n$ ) of the linear parts were 0.77 ( $r = 0.997$ ) for 90M ~40M, 0.91 ( $r = 0.986$ ) for 90E ~30E, and 1.8 ( $r = 0.994$ ) for 80A ~30A. Kivinen  $n$  values increase in the order methanol-water < ethanol-water < acetone-water. The larger  $n$  value of acetone-water mixture indicates that the ionizing power increases with increasing water contents, whereas there is no

**Table 2.** Separated rate constants for methanesulfonyl chloride solvolysis in aqueous methanol (M) and ethanol (E)

Alcohol	/ $10^{-5}s^{-1}$		
% (v/v)	$k_{aa}$	$k_{aw} + k_{wa}$	$k_{ww}$
90M	2.16	6.05	0.204
80M	1.71	10.2	0.817
70M	1.31	14.8	1.84
60M	0.962	19.9	3.27
50M	0.668	24.9	5.10
40M	0.428	26.5	7.36
30M	0.240	25.5	10.1
20M	0.107	21.6	13.1
10M	0.0267	17.0	16.5
90E	1.21	2.62	0.204
80E	0.953	4.96	0.817
70E	0.729	6.65	1.84
60E	0.536	8.49	3.27
50E	0.372	10.6	5.11
40E	0.238	12.7	7.36
30E	0.134	13.2	10.0
20E	0.0596	10.5	13.1
10E	0.0148	5.29	16.5



**Figure 3.** Correlation of logarithms of rate constants for solvolyses of methanesulfonyl chloride at 25 °C  $\log(k/k_0)$  vs.  $\log[H_2O]$  (40M-90M: slope = 0.77 ( $r = 0.994$ ); 30E-90E: slope = 0.918 ( $r = 0.986$ ); 30A-80A: slope = 1.8 ( $r = 0.997$ ) (solvent codes: ●; methanol, ○; ethanol, ▲; acetone).

general base catalysis in acetone-water mixtures, because acetone acts as cosolvent. The increase of high ionizing power ( $Y_{Cl} = 4.57$ ) with water contents in acetone-water mixtures, it mean that bond breaking increases in the transition state. These results are in good agreement with  $S_{AN}/S_N2^{17}$  reaction mechanism with a bond breaking slightly more progressed in the transition state which increases with increasing water contents in acetone-water mixtures.

The activation parameters for solvolyses of MSC in aqueous solvents at 25 °C are summarized in Table 3. The large negative  $\Delta S^\ddagger$  and large positive  $\Delta H^\ddagger$  reveals that the solvolytic reaction proceeds *via* a typical bimolecular reaction.<sup>19</sup> The magnitude of  $\Delta S^\ddagger$  and  $\Delta H^\ddagger$  increase as the water content of the mixtures increases, suggesting that bond formation is very important and bond breaking is of little importance in the transition state. The variation of activation

**Table 3.** Activation parameters of the methanesulfonyl chloride in aqueous alcohol and acetone binary solvent mixtures at 25 °C

v/v%	MeOH		EtOH		Acetone	
	$\Delta H^\ddagger$	$-\Delta S^\ddagger$ (e.u.)	$\Delta H^\ddagger$	$-\Delta S^\ddagger$ (e.u.)	$\Delta H^\ddagger$	$-\Delta S^\ddagger$ (e.u.)
100	17.1	22	13.1	37		
90	17.1	20	16.4	24		
80	16.6	21	16.0	24	13.0	36
70	16.1	22	16.3	22	13.0	35
60	16.3	21	16.2	22	15.1	27
50	16.8	19	16.0	22	14.7	27
40	16.8	19	16.1	21	16.4	21
30	17.8	15	16.6	19	16.3	21
20	18.2	14	17.3	17	16.3	21
10	18.4	13	18.4	14	17.7	16
0 <sup>(H2O)</sup>	19.0	12				

**Table 4.** Kinetic solvent isotope effect of methanesulfonyl chloride at 25 °C and 35 °C

Temp.		$k_{\text{SOH}}$	$k_{\text{SOD}}$	KSIE
25 °C	methanol	$2.67 \times 10^{-5}$	$1.67 \times 10^{-5}$	1.62
	water	$2.04 \times 10^{-4}$	$1.13 \times 10^{-4}$	1.81
35 °C	methanol	$7.36 \times 10^{-5}$	$4.87 \times 10^{-5}$	1.51
	water	$5.96 \times 10^{-4}$	$3.89 \times 10^{-4}$	1.53

parameters with the change of component of mixed solvents indicates that bond formation is important in the transition state, but bond breaking is negligible at the transition state.

The solvent isotope effects for reaction of MSC in methanol and water at 25 °C and 35 °C are summarized in Table 4. The solvent isotope effects are relatively large, which decreases a little at elevated temperature suggesting that the OH bond breaks partially for nucleophilic methanol in the transition state.<sup>14c</sup> This is another piece of evidence in support of the  $S_{\text{A}}\text{N}/S_{\text{N}}2$ <sup>17</sup> mechanism in methanol and in water, in which methanol and water molecules act as a general base catalyst, as shown in Scheme 1.

### Experiment Section

Methanesulfonyl chloride were commercial grade (Wako Gr-grade > 99%). Merk Gr-grade (< 0.1% H<sub>2</sub>O) acetone, ethanol and methanol were used without further purification. D<sub>2</sub>O and CH<sub>3</sub>OD were from Aldrich (99.9% D). Distilled water was redistilled with Buchi Fontavapor 210 and treated using ELGA UHQ PS to obtain specific conductivity of less than  $1.0 \times 10^{-6}$  mhos/cm. Rates were measured conductometrically at least in duplicate as in previous work.<sup>15</sup> Activation parameters were determined using Arrhenius equation and Eyring equation.

### References

- (a) Barker, J. W.; Nathan, W. W. *J. Chem. Soc.* **1936**, 236. (b) Swain, C. G.; Langsdorf, W. P. *J. Am. Chem. Soc.* **1951**, 73, 2813. (c) Rossel, J. B. *J. Chem. Soc.* **1963**, 5183. (d) Yoh, S. D.; Tsuno, Y.; Yukawa, Y. *J. Korean Chem. Soc.* **1984**, 28, 433. (e) Yoh, S. D. *J. Korean Chem. Soc.* **1975**, 19, 240.
- Lee, I.; Rhyu, K. B.; Lee, B. C. *J. Korean Chem. Soc.* **1979**, 23, 277.
- (a) Rogne, O. *J. Chem. Soc.(B)* **1968**, 1294. (b) Kim, W. K.; Lee, I. *J. Korean Chem. Soc.* **1974**, 18, 8.
- (a) Ciuffarin, E.; Senatore, L.; Isola, M. *J. Chem. Soc., Perkin Trans.* **2** **1972**, 468. (b) Stangeland, L. J.; Senatore, L.; Ciuffarin, E. *J. Chem. Soc. Perkin Trans.* **2** **1972**, 852.
- (a) Hall, H. K. Jr. *J. Am. Chem. Soc.* **1956**, 78, 1450. (b) Robertson, R. E.; Laughton, P. M. *Can. J. Chem.* **1957**, 35, 1319. (c) Rossall, B.; Robertson, R. E. *Can. J. Chem.* **1971**, 49, 1451. (d) Ballistreri, F. P.; Cantona, A.; Maccaroni, E.; Tomaselli, G. A.; Tripolone, M. *J. Chem. Soc. Perkin Trans.* **2** **1981**, 438.
- (a) Bentley, T. W.; Carter, G. E.; Harris, H. C. *J. Chem. Soc., Perkin Trans.* **2** **1985**, 938. (b) Bentley, T. W.; Harris, H. C. *J. Chem. Soc., Perkin Trans.* **2** **1986**, 619. (c) Bentley, T. W.; Koo, I. S. *J. Chem. Soc., Chem. Commun.* **1988**, 41.
- Bentley, T. W.; Jones, R. O.; Koo, I. S. *J. Chem. Soc., Perkin Trans.* **2** **1994**, 753.
- Bentley, T. W.; Jones, R. O.; Koo, I. S. *J. Chem. Soc., Perkin Trans.* **2** **1994**, 753.
- Bentley, T. W.; Jones, R. O. *J. Chem. Soc., Perkin Trans.* **2** **1993**, 2351.
- (a) Yang, K.; Koo, I. S.; Lee, I. *J. Phys. Chem.* **1995**, 99, 15035. (b) Yang, K.; Koo, I. S.; Kang, D. H.; Lee, I. *Bull. Korean Chem. Soc.* **1994**, 15, 419.
- (a) Oh, J.; Yang, K.; Koo, I. S.; Lee, I. *J. Chem. Res.* **1993**, 310. (b) Bentley, T. W.; Harris, H. C.; Koo, I. S. *J. Chem. Soc., Perkin Trans.* **2** **1988**, 783. (c) Bentley, T. W.; Koo, I. S.; Norman, S. J. *J. Org. Chem.* **1991**, 56, 1604.
- (a) Bentley, T. W.; Harris, H. C. *J. Org. Chem.* **1988**, 53, 724. (b) Lee, I.; Koo, I. S.; Shon, S. C.; L, H. H. *Bull. Korean Chem. Soc.* **1982**, 3, 92. (c) Bentley, T. W.; Jones, R. O. *J. Chem. Soc., Perkin Trans.* **2** **1993**, 2351.
- (a) Lee, I.; Koo, I. S.; Kang, H. K. *Bull. Korean Chem. Soc.* **1981**, 2, 41. (b) Bentley, T. W.; Jones, R. O.; Koo, I. S. *J. Chem. Soc., Perkin Trans.* **2** **1994**, 753.
- (a) Koo, I. S.; Yang, K.; Kang, K.; Lee, I.; Bentley, T. W. *J. Chem. Soc., Perkin Trans.* **2** **1998**, 1179. (b) Koo, I. S.; Yang, K.; Kang, K.; Lee, I. *Bull. Korean Chem. Soc.* **1998**, 19, 968. (c) Koo, I. S.; Lee, J. S.; Yang, K.; Kang, K.; Lee, I. *Bull. Korean Chem. Soc.* **1999**, 20, 573.
- Bentley, T. W.; Llewellyn, G. *Prog. Phys. Org. Chem.* **1990**, 17, 121.
- Bender, M. L.; Chen, M. C. *J. Am. Chem. Soc.* **1963**, 85, 30.
- We used  $S_{\text{A}}\text{N}/S_{\text{N}}2$  terminology because nucleophilic substitution reaction with product like transition state is termed as an  $S_{\text{A}}\text{N}$  or  $S_{\text{N}}2$ , in view of addition of nucleophilic or leaving of leaving group, respectively.
- Kivinen, A. *Acta Chem. Scand.* **1965**, 19, 835.
- Frost, A.; Pearson, R. G. *Kinetic and Mechanism*, 2nd Ed.; Wiley: New York, 1961; Chap. 7.