

References and Notes

1. A. J. Weinheimer and R. L. Spraggins, *Tetrahedron Lett.*, 5185 (1969).
2. N. Fusetani and K. Hashimoto, *Nippon Suisan Gakkaishi*, **50**, 465 (1984).
3. The ^{13}C NMR (22.5 MHz, CD_3OD , δ) data for fatty acid moiety of GV-5(1) are as follows: 174.9(s), 174.7(s), 174(s), 131.4(d), 131.0(d), 130.9(d), 130.1(d), 129.6(d), 129.3(d), 129.1(d), 128.9(d), 35.3(t), 35.1(t), 34.7(t), 34.5(t), 33.2(t), 32.8(t), 30.9(t), 30.8(t), 30.6(t), 30.4(t), 28.4(t), 27.7(t), 26.8(t), 26.2(t), 26.1(t), 23.9(t), 23.8(t), 14.8(q).
4. T. Taniyama, Ph. D. Dissertation, Graduate School, Osaka University, Osaka 565, Japan, 1988. The structure of the glyceroglycolipid is 1-O- β -D-galactopyranosyl-2',3'-di-O-acyl-D-glycerol[acyl: linolenate (90%), linolate(8%), palmitate(2%)].
5. H. Kikuchi, Y. Tsukitani, T. Manda, T. Fujii, H. Nakaniishi, M. Kobayashi, and I. Kitagawa, *Chem. Pharm. Bull.*, **30**, 3544 (1982), and the structure of the M-5 is 1-O- β -D-galactopyranosyl-2',3'-di-O-acyl-D-glycerol(acyl; myristate: 8-hexadecenoate: palmitate = 1:10:10).
6. TMS-glycerol and methyl TMS-galactoside were identified by direct GLC comparisons (2% SE-30 on Chromosorb WAW DMCS, 80-100 mesh, 3 mm $\times$ 2 m) with authentic samples: TMS-glycerol [column temp. 120°C, carrier gas N_2 at flow rate 25 ml/min., $t_R(\text{min.}) = 5'18''$] and methyl TMS-galactoside [column temp. 170°C, N_2 at flow rate 30 ml/min., $t_R(\text{min.}) = 8'23''$, $9'39''$, $11'01''$].
7. ^{13}C NMR data (22.5 MHz, $\text{D}_2\text{O}-\text{CD}_3\text{OD} = 2:1$, δ) of **2**: 104.4(C-1), 72.1(C-2), 74.0(C-3), 69.9(C-4), 76.4(C-5), 62.2(C-6), 63.6(C-1'), 71.7(C-2'), 71.9(C-3').
8. B. Wickberg, *Acta Chem. Scand.*, **12**, 1187 (1958).
9. A mixture of fatty acid methyl esters was identified by direct GLC comparison (15% polyethylene glycol succinate on Chromosorb WAW DMCS, 80-100 mesh, 3 mm $\times$ 2 m, column temp. 210°C, carrier gas N_2 at flow rate 35 ml/min.) with authentic samples and its ratio was obtained by comparison of peak area for each of the fatty acid methyl esters: $t_R(\text{min.}) = \mathbf{a} \ 2'23''$, $\mathbf{b} \ 4'14''$, $\mathbf{c} \ 11'01''$; $\mathbf{a}:\mathbf{b}:\mathbf{c} = 4:1:9$.
10. S. R. Johns, D. R. Leslie, R. I. Willing, and D. G. Bishop, *Aust. J. Chem.*, **30**, 823 (1977).
11. GV-4; ^1H NMR (90 MHz, CDCl_3 , δ): 5.37(triplet like, olefinic protons) 1.26(methylene protons), 0.89(a deformed triplet, terminal methyl protons), ^{13}C NMR (22.5 MHz, CDCl_3 , δ): 180.1(br.s), 130.5(d), 129.2(d), 128.8(d), 128.7(d), 128.3(d), 128.2(d), 127.9(d), 127.6(d), 34.1(t), 33.5(t), 32.0(t), 31.6(t), 29.8(t), 29.4(t), 29.2(t), 27.3(t), 26.6(t), 25.7(t), 24.8(t), 24.6(t), 22.6(t), 14.1(q), and methyl ester of GV-4; ^1H NMR (90 MHz, CDCl_3 , δ): 5.35(triplet like, olefinic protons), 3.65(s), 1.25(methylene protons), 0.88 (a deformed triplet, terminal methyl protons).
12. Analytical conditions for GLC were the same as described in that of a mixture of fatty acid methyl esters of GV-5⁹: $t_R(\text{min.}) = \mathbf{a} \ 2'24''$, $\mathbf{b} \ 4'19''$, $\mathbf{c} \ 11'03''$; $\mathbf{a}:\mathbf{b}:\mathbf{c} = 2:1:6$.
13. T. Yasumoto, N. Seino, Y. Murakami, and M. Murata, *Biol. Bull.*, **172**, 128 (1987).
14. N. Fusetani and Y. Hashimoto, *Agr. Biol. Chem.*, **39**, 2021 (1975).
15. P. Baruah, N. C. Baruah, R. P. Sharma, J. N. Baruah, P. Kulanthaivel, and W. Herz, *Phytochemistry*, **22** 1741 (1983).
16. I. Kitagawa, Y. Hamamoto, and M. Kobayashi, *Chem. Pharm. Bull.*, **27**, 1934 (1979).
17. Y. Nagai, Y. Isono, and M. Hoshi, *Shishitsu Seikagaku Kenkyu*, **1968**, 151.


Theoretical Studies on [1,3] Sigmatropic Group Rearrangements[†]Ikchoon Lee*, Jeoung Ki Cho, and Hyuck Keun Oh[†]

Department of Chemistry, Inha University, Incheon 402-751

[†]Department of Chemistry, Chonbuk National University, Chonju 560-756

Received April 8, 1988

Sigmatropic rearrangements are not confined to hydrogen atoms.^{1,2} Groups of nearly every type have been found to migrate. We report here semiempirical MO theoretical results on the course of such reactions involving migrating groups with lone pairs(F, OH, NH_2 , SH, Cl) and those which can migrate by either σ - or p-type interactions (CH_3 , NO) between 1,3-2p orbitals of propenyl system. Four types of orbital interactions can be envisaged, since both σ and p type can interact either suprafacially(s) or antarafacially(a) (Scheme 1).



Scheme 1

[†] Determination of Reactivity by MO Theory (part 53).

Table 1. Energy Barriers (ΔH_f ; kcal/mol) and Types of 1,3-Sigmatropic Group Rearrangements Investigated in This Work

X	Method		Type ^a
	AMI	MNDO	
CH ₃	106.51	123.37	sp
NH ₂	75.57	59.01	sp
OH	78.51	82.42	s σ
CN	102.25	91.93	a σ
NO	68.32	53.26	sp,s σ
F	---	102.79	a σ
Cl	---	56.15	s σ
SH	---	57.53	S σ

^a s, a, σ , and p denote suprafacial, antarafacial, sigma, and p-type interactions respectively.

Obviously the σ type should result in retention whereas the p type should give inversion of configuration at the migrating group. For neutral, four electron systems, only [1,3] a σ and [1,3] sp shifts are allowed according to the orbital symmetry rules,³ since normally the interactions are between propenyl-HOMO and migrating group-LUMO in terms of frontier orbital (FMO) theory.⁴

Our MNDO^{5,7} and AMI^{6,7} computations on the transition states for 1,3-shifts of eight groups have shown that (i) antarafacial processes are sterically unfavorable so that suprafacial processes are favored for relatively large migrating groups ([1,3]sp for CH₃ group), (ii) if a lone pair (nonbonding) level is high compared with the nonbonding level (HOMO) of the propenyl system, the lone pair participate in the migration so that six electrons are involved in the [1,3] shift, for which propenyl-LUMO and migrating group-HOMO interactions become dominant and allowed processes change to [1,3]s σ and [1,3]ap ([1,3]s σ for OH, Cl, and SH groups), and (iii) the activation energy barriers were found to decrease in the order CH₃>F>CN>OH>NH₂>SH $\approx$ Cl>NO (Table 1).

Normal four electron [1,3]a σ process was followed only by CN group, while NO exhibited both aspects of σ - and p-type interactions. Symmetry forbidden processes were found to take place for the two 6 electron systems involving NH₂ and F groups; the former has a p-type lone pair so that it should proceed by [1,3]ap, but due to the large steric repulsion of two H atoms it was forced to go through [1,3]sp, whereas the latter, having a rather low lying lone pair, gave [1,3]a σ i.e., behaved like a four electron system.

We conclude that in [1,3]group rearrangements antarafacial

process is always sterically unfavorable and the lone pairs can participate in σ and p type shifts providing two extra electrons to the system so that selection rules can also change.

Acknowledgements. We thank the Ministry of Education and the Korea Center for Theoretical Physics and Chemistry for support of this work.

References

- (a) R. C. Bingham and M. J. S. Dewar, *J. Am. Chem. Soc.*, **94**, 9107 (1972); (b) S. Kato and K. Fukui, *ibid.*, **99**, 684 (1977); (c) W. J. Bouma, D. Poppinger, and L. Radom, *ibid.*, **99**, 6443 (1977); (d) W. J. Bouma and L. Radom, *ibid.*, **101**, 3487 (1979); (e) L. Carlsen and F. Duus, *ibid.*, **100**, 281 (1978); (f) A. D. Isaacson and K. Morokuma, *ibid.*, **97**, 4453 (1975); (g) J. E. Del Bene and W. L. Kochenour, *ibid.*, **98**, 2041 (1976); (h) P. D. Adeney, W. J. Bouma, L. Radom, and W. R. Rodwell, *ibid.*, **102**, 4069 (1980); (i) W. J. Bouma, M. A. Vincent, and L. Radom, *Int. J. Quantum Chem.*, **14**, 767 (1978); (j) W. R. Rodwell, W. J. Bouma, and L. Radom, *ibid.*, **18**, 107 (1980); (k) K. Yamashita, H. Kaminoyama, T. Yamabe, and K. Fukui, *Theor. Chim. Acta*, **60**, 302 (1981); (l) J. K. Cho, I. Lee, and H. K. Oh, and I. H. Cho, *J. Korean Chem. Soc.*, **28**, 217 (1984).
- J. K. Cho, I. Lee, H. K. Oh, and I. H. Cho, *Bull. Korean Chem. Soc.*, **5**, 179 (1984).
- (a) R. B. Woodward and R. Hoffmann, "The Conservation of Orbital Symmetry", Berlin, Verlag Chemie GmbH, 1971; (b) R. G. Pearson, "Symmetry Rules for Chemical Reactions", New York, John Wiley and Sons, 1976; (c) T. L. Gilchrist and R. C. Storr, "Organic Reactions and Orbital Symmetry", 2nd ed., Cambridge University Press, Cambridge, 1979.
- (a) R. B. Woodward and R. Hoffmann, *Angew. Chem. Int. Ed. Engl.*, **8**, 781 (1969); (b) M. J. S. Dewar, *ibid.*, **10**, 761 (1971); (c) K. Fukui, "Theory of Orientation and Stereoselection", Berlin and New York, Springer-Verlag, 1975; (d) K. Fukui, *Acc. Chem. Res.*, **4**, 57 (1971); (e) I. Fleming, "Frontier Orbitals and Organic Chemical Reactions", New York, John Wiley and Sons, 1976.
- M. J. S. Dewar and W. J. Thiel, *J. Am. Chem. Soc.*, **99**, 4899 (1977).
- M. J. S. Dewar, E. G. Zoebisch, E. F. Healy, and J. J. P. Stewart, *J. Am. Chem. Soc.*, **107**, 3902 (1985).
- Available from Quantum Chemistry Program Exchange (QCPE), No. 506.