

# BULLETIN OF THE KOREAN CHEMICAL SOCIETY

VOLUME 6, NUMBER 1, FEBRUARY 20, 1985

## Calculation of NMR Chemical Shift for $5d^n$ Systems (I). Application of the Expansion Method for Spherical Harmonics for Computation of Overlap and Dipole Moment Matrix Elements

Sangwoon Ahn and Euisuh Park

*Department of Chemistry, Jeonbug National University, Jeonju 520, Korea*

Se Woung Oh

*Department of Chemistry, Mogpo National College, Mogpo 580, Korea (Received February 14, 1984)*

The general formulas for SCF atomic orbitals  $|5s\rangle$  and  $|6p\rangle$  have been derived separately by expressing the spherical harmonics part in terms of the coordinate  $(r_1, r_2)$  of the reference point, and by translating the exponential part,  $r^4 \exp(-\beta r)$ , in terms of  $r_1$  and  $r_2$  and the modified Bessel functions. Master formulas for overlap and dipole moment matrix elements are derived. The computed values of overlap and dipole moment matrix elements for hypothetical NO molecule are exactly in agreement with those for the previous methods.

### 1. Introduction

The first systematic and comprehensive study of two center overlap integrals was carried out, adopting Slater type atomic orbitals<sup>1</sup>. The evaluation of two center integrals has been performed using the spheroidal coordinate system<sup>2</sup>. In this spheroidal coordinate system, it is required to transform the spherical polar coordinate to the spheroidal coordinate for evaluation of two center integrals. Since two atoms must lie in one coordinate axis to evaluate two center integral in the spheroidal coordinate system<sup>3</sup>, namely the  $z$  axis, this required, in general, a different coordinate transformation of each matrix element. To overcome the inconvenience of different coordinate transformations, the expansion method for spherical harmonics has been developed by Golding and Stubbs<sup>4</sup>. This method was applied to evaluate specific physical molecular properties such as dipole moments<sup>5,6</sup>, quadrupole moments<sup>7,8</sup>, polarizabilities<sup>9,10</sup> and NMR shifts in paramagnetic systems<sup>11,12</sup>. In the previous works<sup>4,13</sup>, the expansion formulas for Slater type orbitals,  $|1s\rangle$ ,  $|2p\rangle$ ,  $|3d\rangle$ ,  $|2s\rangle$ ,  $|3p\rangle$ ,  $|4d\rangle$ ,  $|3s\rangle$ ,  $|4p\rangle$ ,  $|5d\rangle$ ,  $|4s\rangle$ , and  $|5p\rangle$  have been derived to evaluate two center integrals for any pairs of those Slater type orbitals. However, it is required

to derive the expansion formulas for SCF atomic orbitals<sup>14</sup>  $|5s\rangle$  and  $|6p\rangle$  to calculate the NMR chemical shift for SCF atomic orbitals  $|5d\rangle$ .

The purpose of this work is to derive the general expansion formulas for SCF atomic orbitals  $|5s\rangle$  and  $|6p\rangle$  and to apply these formulas to obtain the analytical formulas for overlap and dipole moment matrix elements for any pairs of SCF atomic orbitals. We also compare the computed results obtained from the analytical formulas with those of Mulliken's method.

### 2. Derivation of the Expansion Formulas for SCF Atomic Orbital $|5s\rangle$ and $|6p\rangle$

If a SCF atomic orbital is assumed to be located at the point B of Figure 1, the spherical harmonic part may be expressed in terms of the reference point A, using the following identities<sup>15</sup>,

$$r^l Y_{lm}(\theta, \phi) = \sum_{l_1=0}^l \sum_{l_2=0}^l \sum_{m_1=-l_1}^{l_1} \sum_{m_2=-l_2}^{l_2} (-1)^{l_1} \delta(l_1+l_2, l) \left\{ \frac{4\pi(2l+1)!}{(2l_1+1)!(2l_2+1)!} \right\}^{\frac{1}{2}} \langle l_1 l_2 m_1 m_2 | l_1 l_2 l m \rangle r_1^{l_1} r_2^{l_2} Y_{l_1 m_1}(\theta_1, \phi_1) Y_{l_2 m_2}(\theta_2, \phi_2) \quad (1)$$

Where  $\langle l_1 l_2 m_1 m_2 | l_1 l_2 l m \rangle$  is Clebsch-Gordan coefficient.

The exponential part may also be translated into the following form

$$r^{K-l} \exp(-\beta r) = 4\pi \sum_{n=0}^{\infty} k_n(r_1, r_2) \sum_{h=-n}^n Y_{nh}^*(\theta_1, \phi_1) Y_{nh}(\theta_2, \phi_2) \quad (2)$$

$$\text{because } |\phi_B\rangle = N r^K \exp(-\beta r) Y_{lm}(\theta, \phi) \quad (3)$$

In equation (3), if  $K=l$ ,  $k_n(r_1, r_2) = b_n(r_1, r_2)^3$ ,

if  $K=l+1$ ,  $k_n(r_1, r_2) = z_n(r_1, r_2)^3$ ,

if  $K=l+2$ ,  $k_n(r_1, r_2) = h_n(r_1, r_2)^{13}$  and

if  $K=l+3$ ,  $k_n(r_1, r_2) = a_n(r_1, r_2)^{13}$

Where

$$a_n(r_1, r_2) = (r_{<} r_{>})^{-\frac{1}{2}} \sum_{i=0}^4 A_i I_{n-\frac{3}{2}+i}(\beta r_{<}) K_{n-\frac{3}{2}+i}(\beta r_{>}) \quad (4)$$

In this expression  $I_v$  and  $K_v$  are the modified Bessel functions and the coefficients,  $A_i$ , are as follows.

$$\begin{aligned} A_0 &= \frac{4n(n-1)}{(2n+1)(2n-1)} r_{<}^2 r_{>}^2 \\ A_1 &= -\frac{4n}{(2n+1)} (r_{<}^3 r_{>} + r_{<} r_{>}^3) \\ A_2 &= \left( r_{<}^4 + r_{>}^4 + \frac{2(8n^2+8n-5)}{(2n+3)(2n-1)} r_{<}^2 r_{>}^2 \right) \\ A_3 &= -\frac{4(n+1)}{(2n+1)} (r_{<}^3 r_{>} + r_{<} r_{>}^3) \\ A_4 &= \frac{4(n+1)(n+2)}{(2n+1)(2n+3)} r_{<}^2 r_{>}^2 \end{aligned}$$

If equation (4) is differentiated with respect to  $\beta$ , we may obtain the expansion formula for  $r^4 \exp(-\beta r)$  as given by

$$\begin{aligned} -\frac{\partial}{\partial \beta} [r^4 \exp(-\beta r)] &= -4\pi \sum_{n=0}^{\infty} \frac{\partial}{\partial \beta} [a_n(r_1, r_2)] \\ &= \sum_{h=-n}^n Y_{nh}^*(\theta_1, \phi_1) Y_{nh}(\theta_2, \phi_2) \\ &= 4\pi \sum_{n=0}^{\infty} d_n(r_1, r_2) \sum_{h=-n}^n Y_{nh}^*(\theta_1, \phi_1) Y_{nh}(\theta_2, \phi_2) \quad (5) \end{aligned}$$

Where

$$d_n(r_1, r_2) = (r_{<} r_{>})^{-\frac{1}{2}} \sum_{i=0}^5 D_i I_{n-\frac{5}{2}+i}(\beta r_{<}) K_{n-\frac{5}{2}+i}(\beta r_{>}) \quad (6)$$

Where




Figure 1. The coordinate system for two center integral of the expansion method.

$$\begin{aligned} D_0 &= \frac{4n(n-1)}{(2n+1)(2n-1)} r_{<}^2 r_{>}^3 \\ D_1 &= -\frac{4n}{(2n+1)} \left( \frac{(3n-2)}{(2n-1)} r_{<}^3 r_{>}^2 + r_{<} r_{>}^4 \right) \\ D_2 &= \left( \frac{(6n+1)}{(2n+1)} r_{<}^4 r_{>} + \frac{2(24n^3+32n^2-8n-5)}{(2n+3)(2n+1)(2n-1)} r_{<}^2 r_{>}^3 + r_{>}^5 \right) \\ D_3 &= -\left( r_{<}^5 + \frac{2(24n^3+40n^2-11)}{(2n+3)(2n+1)(2n-1)} r_{<}^3 r_{>}^2 + \frac{(6n+5)}{(2n+1)} r_{<} r_{>}^4 \right) \\ D_4 &= \frac{4(n+1)}{(2n+1)} \left( r_{<}^4 r_{>} + \frac{(3n+5)}{(2n+3)} r_{<}^2 r_{>}^3 \right) \\ D_5 &= -\frac{4(n+1)(n+2)}{(2n+1)(2n+3)} r_{<}^3 r_{>}^2 \end{aligned}$$

Therefore

$$\begin{aligned} |\phi_B\rangle &= N r^K \exp(-\beta r) Y_{lm}(\theta, \phi) \\ &= 4\pi N \sum_{l_1=0}^l \sum_{l_2=0}^l \sum_{m_1=-l_1}^{l_1} \sum_{m_2=-l_2}^{l_2} (-1)^{l_1} \delta(l_1+l_2, l) \\ &\quad \left\{ \frac{4\pi(2l+1)!}{(2l_1+1)!(2l_2+1)!} \right\}^{\frac{1}{2}} \langle l_1 l_2 m_1 m_2 | l_1 l_2 l m \rangle \\ &\quad \sum_{n=0}^{\infty} d_n(r_1, r_2) r_1^{l_1} r_2^{l_2} \sum_{h=-n}^n Y_{nh}^*(\theta_1, \phi_1) Y_{nh}(\theta_2, \phi_2) \\ &\quad Y_{l_1 m_1}(\theta_1, \phi_1) Y_{l_2 m_2}(\theta_2, \phi_2) \end{aligned}$$

### 3. Overlap Matrix Elements

Combining the SCF atomic orbital located at the point A with the SCF atomic orbital which is expressed in a common coordinate system represented by equation (7) gives the general formula for overlap matrix element.

$$\begin{aligned} \langle \phi_A | \phi_B \rangle &= 4\pi N M \sum_{l_1=0}^l \sum_{l_2=0}^l \sum_{m_1=-l_1}^{l_1} \sum_{m_2=-l_2}^{l_2} \sum_{n=0}^{\infty} \sum_{h=-n}^n \sum_{l_3=|n-l_1|}^{n+l_1} \sum_{m_4=-l_4}^{l_4} \\ &\quad (-1)^{l_2+h+m_3-m} \delta(l_1+l_2, l) \delta(l_3+l_4, l) (2l+1)(2n+1) \\ &\quad \left\{ \frac{(2l_3+1)(2l_4+1)(2l)!}{4\pi(2l_1)!(2l_2)!} \right\}^{\frac{1}{2}} r_1^{l_1} Y_{l_3 m_3}^*(\theta_1, \phi_1) \\ &\quad \int_0^{\infty} d_n(r_1, r_2) r_2^{2+l_2+U} \exp(-\beta r_2) dr_2 \\ &\quad \begin{pmatrix} l_1 & l_2 & l \\ m_1 & m_2 & -m \end{pmatrix} \begin{pmatrix} n & l_1 & l_4 \\ -h & m_1 & m_4 \end{pmatrix} \begin{pmatrix} l_3 & n & l_2 \\ -m_3 & h & m_2 \end{pmatrix} \\ &\quad \begin{pmatrix} l_3 & n & l_2 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} n & l_1 & l_4 \\ 0 & 0 & 0 \end{pmatrix} \quad (8) \end{aligned}$$

Where  $|\phi_A\rangle = M r^U \exp(-\beta r_2) Y_{l_3 m_3}(\theta_2, \phi_2)$

Substitution of the numerical values of 3-j symbols<sup>16</sup> for the specified  $n, l, m, h, l_i$  and  $m_i$  and the radial integrals into equation (8) leads to master formulas of two center overlap matrix elements listed in Table 2.

The radial integrals involving the modified Bessel functions are evaluated in elsewhere<sup>17</sup>. We denote the radial integrals, depending on  $n$  and  $K$  as represented in Table 1.

### 4. Dipole Moment Matrix Elements

If a LCAO MO is used, the dipole moment matrix element for molecular orbitals is given by

$$\langle \Psi_i | \Sigma_i r_i | \Psi_j \rangle = \Sigma_i \Sigma_u \Sigma_v C_{ui} C_{vj} \langle \phi_u | r_i | \phi_v \rangle \quad (9)$$

Where  $\langle \phi_u | r_i | \phi_v \rangle$  is called the dipole moment matrix

TABLE 1: Definition of the Radial Part Integrals

K	$\int_0^\infty d_n(r_1, r_2) r_2 \exp(-\alpha r_2) dr_2$	n			
		0	1	2	3
2	$E_n$	$E_0$			
3	$F_n$	$F_0$	$F_1$		
4	$G_n$	$G_0$	$G_1$	$G_2$	
5	$H_n$	$H_0$	$H_1$	$H_2$	$H_3$
6	$I_n$	$I_0$	$I_1$	$I_2$	$I_3$
7	$J_n$	$J_0$	$J_1$	$J_2$	$J_3$

TABLE 2: Master Formulas for Two Center Overlap Matrix Elements

$$\begin{aligned}
\langle 1s | 5s \rangle &= (32/14175)^{1/2} (\alpha/\beta)^{3/2} E_0 \\
\langle 2s | 5s \rangle &= (32/42525)^{1/2} (\alpha/\beta)^{5/2} F_0 \\
\langle 2p_z | 5s \rangle &= (32/14175)^{1/2} (\alpha/\beta)^{5/2} \cos \theta F_1 \\
\langle 2p_x | 5s \rangle &= (32/14175)^{1/2} (\alpha/\beta)^{5/2} \sin \theta \cos \phi F_1 \\
\langle 2p_y | 5s \rangle &= (32/14175)^{1/2} (\alpha/\beta)^{5/2} \sin \theta \sin \phi F_1 \\
\langle 3s | 5s \rangle &= (8/135)(1/35)^{1/2} (\alpha/\beta)^{7/2} G_0 \\
\langle 3p_z | 5s \rangle &= (8/45)(1/105)^{1/2} (\alpha/\beta)^{7/2} \cos \theta G_1 \\
\langle 3p_x | 5s \rangle &= (8/45)(1/105)^{1/2} (\alpha/\beta)^{7/2} \sin \theta \cos \phi G_1 \\
\langle 3p_y | 5s \rangle &= (8/45)(1/105)^{1/2} (\alpha/\beta)^{7/2} \sin \theta \sin \phi G_1 \\
\langle 3d_{z^2} | 5s \rangle &= (4/135)(1/7)^{1/2} (\alpha/\beta)^{7/2} (2\cos^2\theta - \sin^2\theta) G_2 \\
\langle 3d_{xx} | 5s \rangle &= (8/45)(1/21)^{1/2} (\alpha/\beta)^{7/2} \cos \theta \sin \theta \cos \phi G_2 \\
\langle 3d_{yy} | 5s \rangle &= (8/45)(1/21)^{1/2} (\alpha/\beta)^{7/2} \cos \theta \sin \theta \sin \phi G_2 \\
\langle 3d_{xy} | 5s \rangle &= (8/45)(1/21)^{1/2} (\alpha/\beta)^{7/2} \sin^2\theta \sin \phi \cos \phi G_2 \\
\langle 3d_{x^2-y^2} | 5s \rangle &= (2/45)(1/21)^{1/2} (\alpha/\beta)^{7/2} \sin^2\theta (\cos^2\phi - \sin^2\phi) G_2 \\
\langle 4s | 5s \rangle &= (4/945)(2/5)^{1/2} (\alpha/\beta)^{9/2} H_0 \\
\langle 4p_z | 5s \rangle &= (4/315)(2/15)^{1/2} (\alpha/\beta)^{9/2} \cos \theta H_1 \\
\langle 4p_x | 5s \rangle &= (4/315)(2/15)^{1/2} (\alpha/\beta)^{9/2} \sin \theta \cos \phi H_1 \\
\langle 4p_y | 5s \rangle &= (4/315)(2/15)^{1/2} (\alpha/\beta)^{9/2} \sin \theta \sin \phi H_1 \\
\langle 5s | 5s \rangle &= (8/14175)(\alpha/\beta)^{11/2} I_0 \\
\langle 5p_z | 5s \rangle &= (8/14175)(3)^{1/2} (\alpha/\beta)^{11/2} \cos \theta I_1 \\
\langle 5p_x | 5s \rangle &= (8/14175)(3)^{1/2} (\alpha/\beta)^{11/2} \sin \theta \cos \phi I_1 \\
\langle 5p_y | 5s \rangle &= (8/14175)(3)^{1/2} (\alpha/\beta)^{11/2} \sin \theta \sin \phi I_1
\end{aligned}$$

element in terms of the atomic orbitals. The dipole moment matrix elements along the  $x$ ,  $y$  and  $z$  axes can be obtained by replacing the dipole moment operator  $r$  by  $x$ ,  $y$  and  $z$ .

In this paper, the dipole moment operators are expressed in spherical harmonics form,<sup>18</sup> for convenience, namely

$$\begin{aligned}
x &= (2/3)^{1/2} r \{ Y_{1-1}(\theta, \phi) - Y_{11}(\theta, \phi) \} \\
y &= (2/3)^{1/2} r \{ Y_{1-1}(\theta, \phi) + Y_{11}(\theta, \phi) \} \\
z &= (4/3)^{1/2} r Y_{10}(\theta, \phi)
\end{aligned} \quad (10)$$

To represent the dipole moment matrix elements in analytical forms, the dipole moment operators must be expressed in terms of the coordinate system of the reference point. If an arbitrary point is taken as the reference point, the dipole moment operator may be represented in the general form,

$$r_3 Y_{1P}(\theta_3, \phi_3) = r_2 Y_{1P}(\theta_2, \phi_2) + R Y_{1P}(\theta_2, \phi_2) \quad (11)$$

Where  $P=0$  or  $\pm 1$ .

Combining the general form of the dipole moment operator with the expansion formulas for SCF orbitals gives the general

formula for the imaginary form of the dipole moment matrix element.

$$\begin{aligned}
\langle \phi_A | r_3 Y_{1P}(\theta_3, \phi_3) | \phi_B \rangle &= \langle \phi_A | r_2 Y_{1P}(\theta_2, \phi_2) \\
| \phi_B \rangle &+ R Y_{1P}(\theta_2, \phi_2) \langle \phi_A | \phi_B \rangle
\end{aligned} \quad (12)$$

Where the first part is the imaginary form of the dipole moment matrix element when point A in Figure 1 is taken as a reference point. The general formula is given by

$$\begin{aligned}
\langle \phi_A | r_2 Y_{1P}(\theta_2, \phi_2) | \phi_B \rangle &= NM \sum_{l_1=0}^l \sum_{l_2=0}^l \sum_{m_1=-l_1}^{l_1} \sum_{m_2=-l_2}^{l_2} \sum_{l_4=|n-l_1|}^{n+l_1} \\
&\sum_{l_5=|n-l_2|}^{n+l_2} \sum_{m_4=-l_4}^{l_4} \sum_{m_5=-l_5}^{l_5} \sum_{n=0}^{\infty} \sum_{h=-n}^n \\
&\delta(l_1+l_2, l) (-1)^{l_2-m+h+m_3+m_5} r_1^{l_1} Y_{lm_4}^*(\theta_1, \phi_1) \\
&\int_0^\infty d_n(r_1, r_2) r_2^{l_2+U+3} \exp(-\beta r) dr_2 (2l+1) \\
&(2n+1) (2l_5+1) \left\{ \frac{3(2l_3+1)(2l_4+1)(2l)!}{(2l_1)!(2l_2)!} \right\}^{1/2} \\
&\begin{pmatrix} l_1 & l_2 & l \\ m_1 & m_2 & -m \end{pmatrix} \begin{pmatrix} n & l_1 & l_4 \\ -h & m_1 & m_4 \end{pmatrix} \begin{pmatrix} n & l_2 & l_5 \\ h & m_2 & m_5 \end{pmatrix} \begin{pmatrix} l_3 & 1 & l_5 \\ -m_3 & P & -m_5 \end{pmatrix} \\
&\begin{pmatrix} n & l_1 & l_4 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} n & l_2 & l_5 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} l_3 & 1 & l_5 \\ 0 & 0 & 0 \end{pmatrix}
\end{aligned} \quad (13)$$

Substituting the numerical values of the 3-j symbols for the specified  $n, l, m, h, P, l_i$  and  $m_i$  and the radial integrals gives the imaginary form of the dipole moment matrix elements. The dipole moment matrix elements may be obtained by transforming the imaginary form of the dipole moment matrix elements into the real form. The dipole moment matrix elements between pairs of SCF orbitals centered at two different atoms are listed in appendix.

## 5. Results and Discussion

For a hypothetical NO molecule, the calculated overlap integrals using the analytical formulas listed in Table 2 are represented in Table 3 with Mulliken's results. As shown in Table 3, the numerical values of two center overlap integrals is exactly in agreement with those for Mulliken's method.

As shown in Table 4, the numerical values of the dipole moment matrix elements evaluated using the analytical formulas listed in Appendix are also exactly in agreement with those of the transformation method of the dipole moment matrix element into overlap matrix elements<sup>5</sup>.

TABLE 3: The Numerical Values of Two Center Overlap Integrals for a Hypothetical NO Molecule ( $\alpha=1.950$ ,  $\beta=2.275$  and  $r=1.50\text{\AA}$ )

Overlap integral	Numerical value	
	This method	Mulliken's method
$\langle 1s   5s \rangle$	0.264763	0.264763
$\langle 2s   5s \rangle$	0.402010	0.402010
$\langle 2p_z   5s \rangle$	0.259706	0.259706
$\langle 3s   5s \rangle$	0.504653	0.504653
$\langle 3p_z   5s \rangle$	0.377752	0.377752
$\langle 4s   5s \rangle$	0.570577	0.570577
$\langle 4p_z   5s \rangle$	0.480267	0.480267
$\langle 5s   5s \rangle$	0.601650	0.601650
$\langle 5p_z   5s \rangle$	0.557729	0.557729

**TABLE 4: The Numerical Results for the Dipole Moment Matrix Elements and the Corresponding Transformed Overlap Integrals ( $\alpha=1.950$ ,  $\beta=2.275$  and  $r=1.50$  Å)**

Dipole moment matrix elements	Numerical value	Transformed overlap integrals	Numerical value
$\langle 1s   z   5s \rangle$	0.070480	$(1/\alpha)\langle 2p_z   5s \rangle$	0.070480
$\langle 2s   z   5s \rangle$	0.162092	$(2/5)^{1/2}(1/\alpha)\langle 3p_z   5s \rangle$	0.162092
$\langle 3s   z   5s \rangle$	0.281560	$(14/3)^{1/2}(1/\alpha)\langle 4p_z   5s \rangle$	0.281560
$\langle 4s   z   5s \rangle$	0.414514	$(15/2)^{1/2}(1/\alpha)\langle 5p_z   5s \rangle$	0.414514
$\langle 5s   z   5s \rangle$	0.543160	$(11)^{1/2}(1/\alpha)\langle 6p_z   5s \rangle$	0.543160
$\langle 2p_x   z   5s \rangle$	0.255091	$(5/2)^{1/2}(1/\alpha)\langle 3s   5s \rangle$ $+ (2)^{1/2}(1/\alpha)\langle 3d_{x^2}   5s \rangle$	0.255091
$\langle 3p_x   z   5s \rangle$	0.417419	$(14/3)^{1/2}(1/\alpha)\langle 4s   5s \rangle$ $+ (56/15)^{1/2}(1/\alpha)\langle 4d_{x^2}   5s \rangle$	0.417419
$\langle 4p_x   z   5s \rangle$	0.594937	$(15/2)^{1/2}(1/\alpha)\langle 5s   5s \rangle$ $+ (6)^{1/2}(1/\alpha)\langle 5d_{x^2}   5s \rangle$	0.594937
$\langle 5p_x   z   5s \rangle$	0.768339	$(11)^{1/2}(1/\alpha)\langle 6s   5s \rangle$ $+ (44/5)^{1/2}(1/\alpha)\langle 6d_{x^2}   5s \rangle$	0.768339

Such results indicate that we may evaluate two center integrals such as the overlap and the dipole moment matrix elements for any pairs of SCF orbital  $|5s\rangle$ ,  $|6p\rangle$  without coordinate transformations into spheroidal coordinate. We have evaluated the hyperfine integrals for  $4d^n$  systems<sup>19</sup> and calculated the NMR chemical shift for a  $4d^1$  system in a strong crystal field environment of octahedral and tetragonal symmetries<sup>20,21</sup>.

To calculate the NMR chemical shift and the hyperfine interaction tensor components for  $5d^1$  systems, the expansion formula for  $r^4 \exp(-\beta r)$  is required.

Therefore, this work may be applied to calculate the two center overlap and dipole moment matrix elements for second and third series of transition metal complexes. As mentioned in the previous paragraph, the expansion formula for SCF orbitals  $|5s\rangle$  and  $|6p\rangle$  can be used to calculate the NMR chemical shift and the hyperfine interaction tensor components for  $5d^n$  system.

#### APPENDIX: Master Formulas for Dipole Moment Matrix Elements

$$\begin{aligned}
 \langle 1s | z | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{3/2}(1/\beta) \cos \theta F_1 \\
 \langle 2s | z | 5s \rangle &= (32/42525)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \cos \theta G_1 \\
 \langle 2p_x | z | 5s \rangle &= (4/135)(2/7)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \{G_0 + (2\cos^2\theta - \sin^2\theta) G_2\} \\
 \langle 2p_x | z | 5s \rangle &= (4/45)(2/7)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \cos \theta \sin \theta \cos \phi G_2 \\
 \langle 2p_y | z | 5s \rangle &= (4/45)(2/7)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \cos \theta \sin \theta \sin \phi G_2 \\
 \langle 3s | z | 5s \rangle &= (8/135)(1/35)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta H_1 \\
 \langle 3p_x | z | 5s \rangle &= (8/135)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \{H_0 + (2\cos^2\theta - \sin^2\theta) H_2\} \\
 \langle 3p_x | z | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin \theta \cos \phi H_2 \\
 \langle 3p_y | z | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin \theta \sin \phi H_2 \\
 \langle 3d_{x^2} | z | 5s \rangle &= (4/675)(1/7)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \{4H_1 + 3(2\cos^2\theta - 3\sin^2\theta) H_3\} \\
 \langle 3d_{xx} | z | 5s \rangle &= (8/225)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \cos \phi \{H_1 + (4\cos^2\theta - \sin^2\theta) H_3\}
 \end{aligned}$$

$$\begin{aligned}
 \langle 3d_{yz} | z | 5s \rangle &= (8/225)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \sin \phi \{H_1 + (4\cos^2\theta - \sin^2\theta) H_3\} \\
 \langle 3d_{xy} | z | 5s \rangle &= (8/45)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin^2\theta \sin \phi \cos \phi H_3 \\
 \langle 3d_{x^2-y^2} | z | 5s \rangle &= (4/45)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin^2\theta (\cos^2\phi - \sin^2\phi) H_3 \\
 \langle 4s | z | 5s \rangle &= (4/945)(2/5)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \cos \theta I_1 \\
 \langle 4p_x | z | 5s \rangle &= (4/945)(2/15)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \{I_0 + (2\cos^2\theta - \sin^2\theta) I_2\} \\
 \langle 4p_x | z | 5s \rangle &= (4/315)(2/15)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \cos \theta \sin \theta \cos \phi I_2 \\
 \langle 4p_y | z | 5s \rangle &= (4/315)(2/15)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \cos \theta \sin \theta \sin \phi I_2 \\
 \langle 5s | z | 5s \rangle &= (8/14175)(\alpha/\beta)^{11/2}(1/\beta) \cos \theta J_1 \\
 \langle 5p_x | z | 5s \rangle &= (8/14175)(1/3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \{J_0 + (2\cos^2\theta - \sin^2\theta) J_2\} \\
 \langle 5p_x | z | 5s \rangle &= (8/14175)(3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \cos \theta \sin \theta \cos \phi J_2 \\
 \langle 5p_y | z | 5s \rangle &= (8/14175)(3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \cos \theta \sin \theta \sin \phi J_2 \\
 \langle 1s | x | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{3/2}(1/\beta) \sin \theta \cos \phi F_1 \\
 \langle 2s | z | 5s \rangle &= (32/42525)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \sin \theta \cos \phi G_1 \\
 \langle 2p_x | x | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \cos \theta \sin \theta \cos \phi G_2 \\
 \langle 2p_x | x | 5s \rangle &= (2/135)(2/7)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \{G_0 - [(2\cos^2\theta - \sin^2\theta) - 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] G_2\} \\
 \langle 2p_y | x | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \sin^2\theta \sin \phi \cos \phi G_2 \\
 \langle 3s | x | 5s \rangle &= (8/135)(1/35)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \cos \phi H_1 \\
 \langle 3p_x | x | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin \theta \cos \phi H_2 \\
 \langle 3p_x | x | 5s \rangle &= (4/135)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \{H_0 - [(2\cos^2\theta - \sin^2\theta) - 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] H_2\} \\
 \langle 3p_y | x | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin^2\theta \sin \phi \cos \phi H_2 \\
 \langle 3d_{x^2} | x | 5s \rangle &= (4/1575)(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \cos \phi \{3(4\cos^2\theta - \sin^2\theta) H_3 - 2H_1\} \\
 \langle 3d_{xx} | x | 5s \rangle &= (4/675)(3/7)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \{2H_1 - [(2\cos^2\theta - 3\sin^2\theta) - 5\sin^2\theta(\cos^2\phi - \sin^2\phi)] H_3\} \\
 \langle 3d_{yz} | x | 5s \rangle &= (8/45)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin^2\theta \sin \phi \cos \phi H_3 \\
 \langle 3d_{xy} | x | 5s \rangle &= (2/225)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \sin \phi \{4H_1 + [5\sin \theta(3 - 2\sin^2\phi) - (4\cos^2\theta - \sin^2\theta)] H_3\} \\
 \langle 3d_{x^2-y^2} | x | 5s \rangle &= (2/225)(1/21)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \cos \phi \{4H_1 + [5\sin \theta(3 - 4\sin^2\phi) - (4\cos^2\theta - \sin^2\theta)] H_3\} \\
 \langle 4s | x | 5s \rangle &= (4/945)(2/5)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \sin \theta \cos \phi I_1 \\
 \langle 4p_x | x | 5s \rangle &= (4/315)(2/15)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \cos \theta \sin \theta \cos \phi I_2 \\
 \langle 4p_x | x | 5s \rangle &= (2/945)(2/15)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \{I_0 - [(2\cos^2\theta - \sin^2\theta) - 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] I_2\} \\
 \langle 4p_y | x | 5s \rangle &= (4/315)(2/14)^{1/2}(\alpha/\beta)^{9/2}(1/\beta) \sin^2\theta \sin \phi \cos \phi I_2 \\
 \langle 5s | x | 5s \rangle &= (8/14175)(\alpha/\beta)^{11/2}(1/\beta) \sin \theta \cos \phi J_1 \\
 \langle 5p_x | x | 5s \rangle &= (8/14175)(3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \cos \theta \sin \theta \cos \phi J_2 \\
 \langle 5p_x | x | 5s \rangle &= (4/14175)(1/3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \{J_0 - [(2\cos^2\theta - \sin^2\theta) - 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] J_2\} \\
 \langle 5p_y | x | 5s \rangle &= (8/14175)(3)^{1/2}(\alpha/\beta)^{11/2}(1/\beta) \sin^2\theta \sin \phi \cos \phi J_2 \\
 \langle 1s | y | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{3/2}(1/\beta) \sin \theta \sin \phi F_1 \\
 \langle 2s | y | 5s \rangle &= (32/42525)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \sin \theta \sin \phi G_1 \\
 \langle 2p_x | y | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \cos \theta \sin \theta \sin \phi G_2 \\
 \langle 2p_x | y | 5s \rangle &= (32/14175)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \sin^2\theta \sin \theta \cos \phi G_2 \\
 \langle 2p_y | y | 5s \rangle &= (2/135)(2/7)^{1/2}(\alpha/\beta)^{5/2}(1/\beta) \{G_0 - [(2\cos^2\theta - \sin^2\theta) + 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] G_2\} \\
 \langle 3s | y | 5s \rangle &= (8/135)(1/35)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin \theta \sin \phi H_1 \\
 \langle 3p_x | y | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \cos \theta \sin \theta \sin \phi H_2 \\
 \langle 3p_x | y | 5s \rangle &= (8/45)(1/105)^{1/2}(\alpha/\beta)^{7/2}(1/\beta) \sin^2\theta \sin \phi \cos \phi H_2
 \end{aligned}$$

$$\begin{aligned}
\langle 3p_y | y | 5s \rangle &= (4/135)(1/21)^{1/2} (\alpha/\beta)^{7/2} (1/\beta) \{H_0 - [(2\cos^2\theta - \sin^2\theta) + 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] H_2\} \\
\langle 3d_{xz} | y | 5s \rangle &= (4/1575)(\alpha/\beta)^{7/2} (1/\beta) \sin\theta \sin\phi \{3(4\cos^2\theta - \sin^2\theta) H_3 - 2H_1\} \\
\langle 3d_{zx} | y | 5s \rangle &= (8/45)(1/21)^{1/2} (\alpha/\beta)^{7/2} (1/\beta) \cos\theta \sin^2\theta \sin\phi \cos\phi H_3 \\
\langle 3d_{yz} | y | 5s \rangle &= (4/675)(3/7)^{1/2} (\alpha/\beta)^{7/2} (1/\beta) \cos\theta \{2H_1 + [(2\cos^2\theta - 3\sin^2\theta) - 5\sin^2\theta(\cos^2\phi - \sin^2\phi)] H_3\} \\
\langle 3d_{xy} | y | 5s \rangle &= (2/225)(1/21)^{1/2} (\alpha/\beta)^{7/2} (1/\beta) \sin\phi \cos\phi \{[5\sin\theta(4\cos^2\phi - 3) + (4\cos^2\theta - \sin^2\theta)] H_3 - 4H_1\} \\
\langle 3d_{x^2-y^2} | y | 5s \rangle &= (2/225)(1/21)^{1/2} (\alpha/\beta)^{7/2} (1/\beta) \sin\theta \sin\phi \{[5\sin\theta(3 - 4\sin^2\phi) + (4\cos^2\theta - \sin^2\theta)] H_3 - 4H_1\} \\
\langle 4s | y | 5s \rangle &= (4/945)(2/5)^{1/2} (\alpha/\beta)^{9/2} (1/\beta) \sin\theta \sin\phi I_1 \\
\langle 4p_x | y | 5s \rangle &= (4/315)(2/15)^{1/2} (\alpha/\beta)^{9/2} (1/\beta) \cos\theta \sin\theta \sin\phi I_2 \\
\langle 4p_y | y | 5s \rangle &= (4/315)(2/15)^{1/2} (\alpha/\beta)^{9/2} (1/\beta) \sin^2\theta \sin\phi \cos\phi I_2 \\
\langle 4p_z | y | 5s \rangle &= (2/945)(2/15)^{1/2} (\alpha/\beta)^{9/2} (1/\beta) \{I_0 - [(2\cos^2\theta - \sin^2\theta) + 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] I_2\} \\
\langle 5s | y | 5s \rangle &= (8/14175)(\alpha/\beta)^{11/2} (1/\beta) \sin\theta \sin\phi J_1 \\
\langle 5p_x | y | 5s \rangle &= (8/14175)(3)^{1/2} (\alpha/\beta)^{11/2} (1/\beta) \cos\theta \sin\theta \cos\phi J_2 \\
\langle 5p_y | y | 5s \rangle &= (8/14175)(3)^{1/2} (\alpha/\beta)^{11/2} (1/\beta) \sin^2\theta \sin\phi \cos\phi J_2 \\
\langle 5p_z | y | 5s \rangle &= (4/14175)(1/3)^{1/2} (\alpha/\beta)^{11/2} (1/\beta) \{J_0 - [(2\cos^2\theta - \sin^2\theta) + 3\sin^2\theta(\cos^2\phi - \sin^2\phi)] J_2\}
\end{aligned}$$

**Acknowledgment.** This work is supported, partly, by the Ministry of Education.

## References

- (1) R. S. Mulliken, C. A. Rieke, D. Orloff and H. Orloff, *J. Chem. Phys.*, **17**, 248 (1949).
- (2) S. P. McGlynn, L. C. Vanquickenborne, M. Kinoshita and D. G. Carroll, "Introduction to Applied Quantum Chemistry," p. 44, Holt, Rinehart and Winston, Inc., New York, U.S.A., 1972.
- (3) S. Ahn, *J. Korean Chem. Soc.*, **22**, 117 (1978).
- (4) L. C. Stubbs, Unpublished Thesis (Honor Degree, University of New South Wales (1973).
- (5) S. Ahn, *J. Korean Chem. Soc.*, **22**, 229 (1978).
- (6) S. Ahn and B. B. Park, *J. Korean Chem. Soc.*, **22**, 357 (1978).
- (7) S. Ahn and J. S. Ko, *J. Korean Chem. Soc.*, **23**, 296 (1979).
- (8) S. Ahn, Unpublished Work.
- (9) S. Ahn, *J. Korean Chem. Soc.*, **22**, 52 (1978).
- (10) S. Ahn, *J. Korean Chem. Soc.*, **22**, 62 (1978).
- (11) R. M. Golding and L. C. Stubbs, *Proc. Roy. Soc.*, **A354**, 223 (1977).
- (12) R. M. Golding, R. O. Pascual and S. Ahn, *J. Magn. Reson.*, **46**, 406 (1982).
- (13) S. Ahn and S. W. Oh, *J. Korean Chem. Soc.*, **23**, 125 (1979).
- (14) E. Clementi, D. L. Raimondi and W. P. Reinhart, *J. Chem. Phys.*, **47**, 1300 (1967); E. Clementi and D. L. Raimondi, *J. Chem. Phys.*, **38**, 2686 (1963).
- (15) M. Moshinsky, *Nuclear Phys.*, **13**, 104 (1959).
- (16) M. Rotenberg, "The 3n-j Symbols," Cambridge University Press, London, U.K., 1959.
- (17) S. Ahn and S. W. Oh Unpublished Work.
- (18) A. R. Edmonds, "Angular Momentum in Quantum Mechanics", Princeton University Press, New Jersey, U.S.A., 1974.
- (19) S. Ahn, H. C. Suh and J. S. Ko, *Bull. Korean Chem. Soc.*, **4**, 17 (1983).
- (20) S. Ahn, S. W. Oh and E. Park, *Bull. Korean Chem. Soc.*, **4**, 64 (1983).
- (21) S. Ahn, S. W. Oh and E. Park, *Bull. Korean Chem. Soc.*, **5**(2) (1984) in press.

## The Crystal and Molecular Structure of Phlorizin Dihydrate

Whanchul Shin

Department of Chemistry, College of Natural Sciences, Seoul National University, Seoul 151, Korea

(Received June 21, 1984)

The crystal structure of phlorizin, a  $\beta$ -D-glucopyranoside of a flavonoid dihydrochalcone phloretin, has been determined by single crystal diffraction methods using diffractometer data obtained by the  $\omega$ - $2\theta$  scan technique with Cu K $\alpha$  radiation from a crystal with space group symmetry  $P2_12_12_1$  and unit cell parameters  $a=4.9094$  (2),  $b=19.109$  (1),  $c=23.275$  (4) Å. The structure was solved by direct methods and refined by full-matrix least-squares to a final  $R=0.047$  for the 1697 observed reflections. The dihydrochalcone moiety is flat and fully extended. The glucose ring has the  ${}^4C_1$  chair conformation and the conformation of the primary alcohol group is *gauche-gauche*. The crystal packing is dominated by an extensive hydrogen bonding pattern. There are one strong and two weak intramolecular hydrogen bonds in the phlorizin molecule.

## Introduction

Some of the flavonoid dihydrochalcones and their saccharide derivatives display an intensely sweet taste and have potential use for the nonnutritive sweeteners, while the cor-

responding flavanones are usually bitter compounds.<sup>1-3</sup>

Little is known about the conformational characteristics of the flavonoid dihydrochalcones themselves and their modes of linkage to the saccharides. Phlorizin, 2- $\beta$ -D-glucopyranoside of phloretin, is a sweet compound with bitter aftertaste.