

# Notes

## Syntheses of 3-Pyrimidyl- and 3-Pyranyl-5,6-benzocoumarin Derivatives

Ibrahim M. El-Deen,\* El-Sayed I. Al-Wakeel, and Ahmed G. El-Mawla

\*Faculty of Education, Suez Canal University, Port-said, Egypt

Faculty of Science, Suez Canal University, Ismailia, Egypt

Received October 3, 2001


**Keywords :** Syntheses, Heterocyclyl-5,6-benzocoumarins.

Sulpha drugs are well recognized for their various physiological activities,<sup>1,2</sup> likewise, many pyrimidine derivatives are used as therapeutic agents,<sup>3-8</sup> 5,6-Benzocoumarin derivatives show antimicrobial,<sup>9</sup> antiinflammatory<sup>10</sup> and anticancer<sup>11</sup> activities. The present work describes the syntheses of some new heterocyclyl-benzocoumarins, starting from 3-acetyl-5,6-benzocoumarin (**1**), which are depicted in Scheme 1 and 2.


3-(2'-Formyl-1'-chlorovinyl)-5,6-benzocoumarin (**2**) was prepared from 3-acetyl-5,6-benzocoumarin (**1**) and DMF-POCl<sub>3</sub>, according to literature procedure.<sup>12</sup> Treatment of

compound **2** with thiourea, 3-amino-1,2,4-triazole, 2-aminobenzimidazole and 3-amino-5-phenylpyrazole in dimethyl formamide gave the corresponding 3-(2'-mercapto or 1'',2'',4-triazolo[1',2'-b] or benzimidazole[1',2'-b] or 5''-phenylpyrazolo[1',2'-b]pyrimidin-6'-yl)-5,6-benzocoumarins (**3-6**).

It has been reported recently<sup>13-16</sup> that 3-(2'-formyl-1'-chlorovinyl)-5,6-benzocoumarin (**2**) reacts with chloroacetic acid in the presence of Ac<sub>2</sub>O-AcONa to afford the corresponding 3-(3'-chloro-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarin (**7**).



Scheme 1



Scheme 2

Treatment of compound **7** with sodium azide in acetic acid gave 3-(1'',2'',3''-triazolo[3',4'-b]pyran-6'-yl)-5,6-benzocoumarin (**8**). Also, compound **7** reacted with thiourea, hydrazine monosulphate and aromatic amines (namely aniline and *p*-toluidine) in ethanol to give 3-(2''-thioxo-2''H-imidazolo[3',4'-b]pyran-6'-yl)-5,6-benzocoumarin (**9**) and 3-(substituent-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarins (**10a-c**).

### Experimental Section

Melting points were determined on a Boetium Hostage apparatus and uncorrected. IR spectra were recorded on a Perkin-Elmer FTIR 1725 spectrometer. The H-NMR spectra were recorded on a General Electric QE 300, and chemical shifts were given with respect to TMS. Mass spectra were obtained on a VG Autospec CEI and FAB<sup>+</sup> and a Hewlett Packard MS-Engine thermospray. Microanalyses were conducted using an elemental analyzer 116.

**General procedure for synthesis of 3-(substituentpyrimidyl)-5,6-benzocoumarins (3-6).** A mixture of **2** (0.01 mol) and aminoheterocycles such as 3-aminotriazole, 2-aminobenzimidazole and 3-amino-5-phenylpyrazole (0.01 mol) or thiourea (0.01 mol), and potassium carbonate (0.03 mol) in DMF (60 mL) was heated under reflux for 6 hr. The solid formed after cooling was filtered off, dried and recrystallized from ethanol to give corresponding product **4-6**. After the reaction with thiourea, the reaction mixture was

cooled and acidified with diluted hydrochloric acid (2%). The product obtained was filtered, washed with water, dried and recrystallized from ethanol to give **3**.

3-(2'-Thioxo-2H-pyrimidin-6'-yl)-5,6-benzocoumarins (**3**), yield 64%; mp 205 °C; IR (cm<sup>-1</sup>) 3253(NH), 1721 (lactone of coumarin); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.19-8.20 (m, 8H, ArH and H-5 of pyrimidine), 8.79 (d, 1H, H-3 of pyrimidine), 10.20 (s, 1H, NH); Mass (m/z) 306 (51) M<sup>+</sup>; Found: C, 66.31; H, 3.13; N, 8.89; S, 10.21. C<sub>17</sub>H<sub>10</sub>N<sub>2</sub>O<sub>2</sub>S requires: C, 66.66; H, 3.27; N, 9.15; S, 10.45.

3-(1'',2'',4-Triazolo[1,2'-b]pyrimidin-6'-yl)-5,6-benzocoumarin (**4**), yield 93%; mp 175 °C; IR (cm<sup>-1</sup>) 1723 (lactone of coumarin), 1630 (C=N); Mass (m/z) 314 (38) M<sup>+</sup>; Found: C, 68.62; H, 2.97; N, 17.47. C<sub>18</sub>H<sub>10</sub>N<sub>4</sub>O<sub>2</sub> requires: C, 68.79; H, 3.18; N, 17.83.

3-(Benzimidazole[1',2'-b]pyrimidin-6'-yl)-5,6-benzocoumarin (**5**), yield 91%; mp 147 °C; IR (cm<sup>-1</sup>) 1719 (lactone of coumarin), 1628 (C=N); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.01-8.21 (m, 12H, ArH and H-5 of pyrimidine), 8.78 (d, 1H, H-4 of pyrimidine); Mass (m/z) 363 (42) M<sup>+</sup>; Found: C, 75.86; H, 3.37; N, 11.39. C<sub>23</sub>H<sub>13</sub>N<sub>2</sub>O<sub>2</sub> requires: C, 76.05; H, 3.58; N, 11.57.

3-(5''-Phenylpyrazolo[1',2'-b]pyrimidin-6'-yl)-5,6-benzocoumarin (**6**), yield 64%; mp 184 °C; IR (cm<sup>-1</sup>) 1721 (lactone of coumarin), 1629 (C=N); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 7.13-8.21 (m, 14H, ArH; pyrazol and H-5 of pyrimidine), 8.77 (d, H-4 of pyrimidine); Mass (m/z) 389 (82) M<sup>+</sup>;

Found: C, 77.00; H, 3.49; N, 10.52.  $C_{25}H_{15}N_3O_2$  requires: C, 77.12; H, 3.85; N, 10.79.

**3-(3'-Chloro-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarin (7).** A mixture of **2** (0.01 mol), chloroacetic acid (0.01 mol), acetic anhydride (0.01 mol) and fused sodium acetate (0.02 mol), was fused on a hot plate for 5-10 min. The reaction mixture was added to acetic acid (50 mL) and heated under reflux for 6 hr., then cooled and poured onto water. The resulting product was filtered off, washed with water, dried and recrystallized from ethanol to give **7**, yield 61%; mp 155-156 °C; IR ( $cm^{-1}$ ) 1729-1719 (br. lactones of coumarin and pyrane);  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  7.19-8.27 (m, 9H, ArH and pyrane ring); Mass (m/z) 325 (63)  $M^+$ ; Found: C, 66.32; H, 2.49; Cl, 10.72.  $C_{18}H_9ClO_4$  requires: C, 66.56; H, 2.77; Cl, 10.94.

**3-(1'',2'',3''-Triazolo[3',4'-b]pyran-6'-yl)-5,6-benzocoumarin (8).** A solution of **7** (0.01 mol) and sodium azide (0.01 mol) in acetic acid (50 mL) was heated in water-bath for 6 hr., then cooled and poured onto water. The resulting product was filtered off, washed with water, dried and recrystallized from ethanol to give **8**, yield 80%; mp 302 °C; IR ( $cm^{-1}$ ) 1729-1721 (lactones of coumarin and pyrane ring); Mass (m/z) 331 (36)  $M^+$ ; Found: C, 65.01; H, 2.48; N, 12.36.  $C_{18}H_9N_3O_4$  requires: C, 65.25; H, 2.72; N, 12.69.

**3-(2'-Thioxo-2H-imidazo[3',4'-b]pyran-6'-yl)-5,6-benzocoumarin (9).** A mixture of **7** (0.01 mol), thiourea (0.01 mol) and potassium carbonate (0.02 mol) in ethanol (50 mL) was heated under reflux for 6 hr. The reaction mixture was cooled and acidified with diluted HCl (2 mol/L). The deposited solid was filtered off, washed with water, dried and recrystallized from ethanol to give **9**, yield 55%; mp 210 °C; IR ( $cm^{-1}$ ) 3185 (NH), 1732-1717 (lactones of coumarin and pyrane ring);  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  7.20-9.19 (m, 8H, ArH and pyrane ring), 10.51-10.53 (br.s, 2H, NH); Mass (m/z) 362(73)  $M^+$ ; Found: C, 62.63; H, 2.49; N, 7.51; S, 8.48.  $C_{19}H_{10}N_2O_4S$  requires: C, 62.98; H, 2.76; N, 7.73; S, 8.84.

**3-(3'-Substituent-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarins (10a-c).** A solution of **7** (0.01 mol) and aromatic amines namely, aniline and p-toluidine (0.01 mol) or hydrazine monosulphate (0.01 mol), and sodium acetate (0.02 mol) in ethanol (70 mL) was heated under reflux for 4 hr. The product formed after cooling was filtered off, washed with water and recrystallized from ethanol, to give **10a-c**.

**3-(3'-Phenylamino-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarins (10a),** yield 77%; mp 190 °C; IR ( $cm^{-1}$ ) 3189

(NH), 1728-1717 (lactones of coumarin and pyrane ring);  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  7.02-8.20 (m, 14H, ArH, and pyrane ring), 10.31 (s, 1H, NH); Mass (m/z) 381 (36)  $M^+$ ; Found: C, 75.27; H, 3.66; N, 3.32.  $C_{24}H_{15}NO_4$  requires: C, 75.59; H, 3.94; N, 3.67.

**3-(3'-p-Methylphenylamino-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarins (10b),** yield 61%; mp 183 °C; IR ( $cm^{-1}$ ) 3186 (NH), 1726-1716 (lactones of coumarin and pyrane ring);  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  2.32 (s, 3H, CH<sub>3</sub>), 7.10-8.19 (m, 13H, ArH and pyrane ring), 10.33 (s, 1H, NH); Mass (m/z) 395 (39)  $M^+$ ; Found: C, 75.64; H, 4.03; N, 3.25.  $C_{25}H_{17}NO_4$  requires: C, 75.95; H, 4.30; N, 3.54.

**3-(3'-Hydrazino-2'-oxo-2'H-pyran-6'-yl)-5,6-benzocoumarins (10c),** yield 53%; mp 240 °C; IR ( $cm^{-1}$ ) 3340, 3251, 3180 (NH, NH<sub>2</sub>), 1729-1719 (lactones of coumarin and pyrane ring);  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  9.23 (s, 1H, NH); Mass (m/z) 320 (42)  $M^+$ ; Found: C, 67.29; H, 3.51; N, 8.44.  $C_{18}H_{12}N_2O_4$  requires: C, 67.50; H, 3.75; N, 8.75.

## References

- Guarneri, M. *Bull. Chem. Form.* **1990**, 99, 259.
- Albert, O.; Bergonzio, G. *Farmaco (Pavia) Ed. Sci.* **1961**, 16, 557.
- Wierzxhowsk, K. L.; Lifonskao, E.; Surger, D. *J. Am. Chem. Soc.* **1965**, 87, 4621.
- Howard, J. B.; Cevik, N.; Murphy, L. M. *Cancer Chemother.* **1966**, 50, 287.
- Flco, A. F.; Goodwier, G. W.; Hatchings, H. G.; Rollo, R. I.; Russell, B. P. *Brit. J. Pharmacol.* **1951**, 6, 186.
- Jain, R.; Pandey, P. *J. Indian Chem. Soc.* **1988**, 65, 354.
- Hampshier, J.; Hebbom, P.; Trigggle, M. M.; Trigggle, J. D.; Fickers, S. *J. Med. Chem.* **1965**, 8, 745.
- Gerg, G. H.; Prakash, C. *J. Med. Chem.* **1971**, 4, 175.
- Hammad, A.; El-Sayed, A. S.; Islam, I. E.; Shafik, N. *J. Chem. Soc. Pax.* **1990**, 12, 292 (Chem. Abstr. 1990, 115, 71279s).
- Kulkarni, G. M.; Kulkarni, H. V.; Patil, V. D.; Shridhar, D. B.; Laxmana, M. *Rev. Roum. Chim.* **1990**, 35, 549.
- Harvey, R. G.; Cortex, C.; Ananthanarayan, T. P.; Schmolka, S. *J. Org. Chem.* **1988**, 53, 3936.
- Gzemey, P.; Hartmann, H. *J. Prakt. Chem.* **1982**, 324, 225.
- El-Deen, I. M. *Chinese J. Chem.* **1998**, 16, 533 (Chem. Abstr. 1999, 130, 223197y).
- El-Deen, I. M. *Chinese J. Chem.* **1999**, 17, 391 (Chem. Abstr. 1999, 131, 228703r).
- El-Deen, I. M.; Abd El-Fattah, M. E. *Indian J. Heterocycl. Chem.* **1999**, 8, 319 (Chem. Abstr. 1999, 131, 170327c).
- El-Deen, I. M.; Ibrahim, H. K. *Phosphorus Sulfur Silicon* **2000**, 160, 241.