

## Synthesis of Cyclic Compounds Having *exo*-Methylene Groups through the Diels-Alder Reactions of Vinyl Allenes Obtained from Propargyl Bromide and Indium

Kooyeon Lee and Phil Ho Lee\*

Department of Chemistry and Institute for Basic Science, Kangwon National University, Chunchon 200-701, Korea

\*E-mail: phlee@kangwon.ac.kr

Received August 10, 2007

**Key Words :** Diels-Alder reaction, Indium, *exo*-Methylene compound, Propargyl bromide, Vinyl allene

Because allene is a very interesting compound having a hybrid character of C-C double and triple bond, vinyl allenenes have been recognized as versatile building blocks in organic synthesis.<sup>1</sup> In particular, vinyl allenenes take part in not only the Diels-Alder reaction<sup>2</sup> as the 1,3-diene moiety but also transition metal-catalyzed organic reactions,<sup>3</sup> affording efficient synthetic methods for complex ring compounds. However, because it is not easy to effectively prepare a variety of vinyl allenenes, its application to organic reactions has been limited despite the potential of vinyl allenenes in organic synthesis. Although vinyl allenenes were used in the Diels-Alder reactions, development of synthetic method of cyclic compounds having *exo*-methylene group is still required due to its utility in synthesis of natural products with biological activities.<sup>4</sup> Recently, we have demonstrated that allenylindiums generated in situ from indium and propargyl bromides are effective cross-coupling partners in palladium-catalyzed cross-coupling reactions to produce substituted allenenes in excellent yields.<sup>5</sup> In continuation of our studies directed toward preparative method of vinyl allenenes with allenylindium, we describe herein the Diels-Alder reaction of vinyl allenenes possessing 3,4- and 4,5-disubstituents and ketone group with a variety of dienophiles to give cyclic compounds having *exo*-methylene group (Scheme 1).

First, 3-methyl-4-phenyl-1,2,4-pentatriene (**1a**) as vinyl allene was prepared from the reaction of  $\alpha$ -bromostyrene with allenylindium obtained from indium and 1-bromo-2-butyne and then, the Diels-Alder reactions with dienophiles were examined to obtain cyclic compounds having *exo*-methylene group. The results are summarized in Table 1. Reaction of **1a** with maleic anhydride (**2a**) produced the Diels-Alder adduct **3a** in 96% yield in toluene at 100 °C for

3 h (entry 1). Also, 3-ethyl-4-phenyl-1,2,4-pentatriene (**1b**) reacted with a variety of dienophiles such as **2a**, *N*-phenyl maleimide (**2b**), dimethyl maleate (**2c**), and 1,4-naphthoquinone (**2d**), producing the desired products (**3b**, **3c**, **3d**, and **3e**) in good to excellent yields in toluene (entries 2-5). Stimulated by these results, tandem cross-coupling reaction of  $\alpha$ -bromostyrene with 1-bromo-2-pentyne and indium (1 equiv.) in the presence of Pd(PPh<sub>3</sub>)<sub>4</sub> (4 mol%) in DMF followed by Diels-Alder reaction with **2a** was attempted to obtain **3b** in one-pot procedure. However, the desired product **3b** was produced in 7% yield in DMF. Although the cross-coupling product **1b** was produced smoothly, the following Diels-Alder reaction did not proceed effectively. Treatment of **1b** with **2a** did not proceed in THF and DMF. In addition, hetero Diels-Alder reactions were tested. Subjecting **1b** to ethyl glyoxylate (**2e**) provided exclusively 3,6-dihydro-2*H*-pyran (**3f**) having *exo*-methylene group in 80% yield (entry 6). No constitutional isomeric product is formed in this reaction, indicating that the electron-rich central carbon of vinyl allene preferentially adds to the more electron-deficient carbon of dienophile. Although compound **1b** reacted with ethyl acrylate (**2f**) to afford mixture of





Figure 1. Vinyl allenenes.



Scheme 1. Synthesis of cyclic compounds having *exo*-methylene group from vinyl allenenes and dienophiles.





Figure 2. Dienophiles.

**Table 1.** [4+2] Cycloaddition reaction of vinyl allenes with dienophiles<sup>a</sup>

Entry	Reactants	Time (h)	Products	Yield (%) <sup>b</sup>
1	1a / 2a	3		96
2	1b / 2a	3		80
3	1b / 2b	4		85
4	1b / 2c	18		30 70 <sup>c</sup>
5	1b / 2d	5		43 87 <sup>c</sup>
6	1b / 2e	20		80 <sup>d</sup>
7	1b / 2f	20		76 <sup>e</sup> (1:2) <sup>f</sup>
8	1c / 2a	13		20 <sup>g</sup> 77
9	1c / 2b	13		40 <sup>g</sup> 83
10	1c / 2e	16		30 <sup>h</sup> 87 <sup>i</sup>
11	1d / 2b	12		92(1:1.5) <sup>g,j</sup>
12	1e / 2b	16		74

<sup>a</sup>Reactions were carried out with vinyl allene (1 equiv.) and dienophile (1 equiv.) unless otherwise noted. Reaction conditions: Toluene (100 °C) for entries 1-7 and 11-12. CH<sub>2</sub>Cl<sub>2</sub> (25 °C) for entries 8 and 9. CH<sub>2</sub>Cl<sub>2</sub> (65 °C) for entry 10. <sup>b</sup>Isolated yields. <sup>c</sup>Dienophile (3 equiv.) was used. <sup>d</sup>Dienophile (30 equiv.) was used. <sup>e</sup>Dienophile (5 equiv.) was used. <sup>f</sup>Regioisomeric ratio. Ethyl 4-ethyl-5-methylene-3-phenylcyclohex-3-enecarboxylate was produced in major. <sup>g</sup>Toluene (25 °C) was used. <sup>h</sup>CH<sub>2</sub>Cl<sub>2</sub> (25 °C) was used. <sup>i</sup>Dienophile (2 equiv.) was used. <sup>j</sup>Diastereomeric ratio.

**Figure 3.** Benzene derivatives from Diels-Alder reaction and aromatization.

regioisomeric products in 76% yield, ethyl 4-ethyl-5-methylene-3-phenylcyclohex-3-enecarboxylate was produced in major (entry 7). Although exposure of vinyl allene **1c** to **2a** gave *exo*-methylenecyclohexene derivative **3h** in 20% yield in toluene, adduct **3h** was produced in 77% yield in CH<sub>2</sub>Cl<sub>2</sub> at room temperature for 13 h (entry 8). In the case of **2b**, [4+2] cycloaddition adduct **3i** was obtained in 83% yield in CH<sub>2</sub>Cl<sub>2</sub> at room temperature for 13 h (entry 9). Reaction of **1c** with **2e** furnished pyran **3j** having *exo*-methylene group in 87% yield with complete regioselectivity (entry 10). Vinyl allenes (**1d** and **1e**) obtained from vinyl triflates and allenylindium were treated with **2b** to produce the desired products (**3k** and **3l**) in 92% (dr = 1:1.5) and 74% yields, respectively (entries 11 and 12). Dienophiles having triple bond such as ethyl propiolate (**2g**) and DMAD (**2h**) were used to Diels-Alder reaction to give rise to multi-substituted benzene derivatives. Reaction of **1b** with **2g** produced selectively ethyl 4-ethyl-3-methyl-5-phenyl benzoate (**4a**) in 65% yield in toluene at 100 °C for 20 h. Compound **1a** and **1c** were treated with DMAD to afford **4b** and **4c** in 82% (toluene, 100 °C, 4 h) and 95% (CH<sub>2</sub>Cl<sub>2</sub>, 65 °C, 15 h) yields, respectively, through Diels-Alder reaction followed by aromatization.

In summary, we have shown that cyclic compounds having *exo*-methylene group were selectively produced through the Diels-Alder reaction of vinyl allenes obtained from propargyl bromide and indium with a variety of symmetric and unsymmetric dienophiles in good to excellent yields. Also, dienophiles having triple bond gave multi-substituted benzene derivatives through Diels-Alder reaction followed by aromatization.

## Experimental Section

**5-Methyl-4-methylene-6-phenyl-3a,4,7,7a-tetrahydroisobenzofuran-1,3-dione (3a):** A mixture of 3-methyl-4-phenyl-1,2,4-pentatriene (**1a**) (46.8 mg, 0.3 mmol) and maleic anhydride (29.4 mg, 0.3 mmol) in toluene (1.0 mL) was heated at 100 °C for 4 h. The reaction mixture was cooling to room temperature and then quenched with NaHCO<sub>3</sub> (sat. aq.). The aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (20 mL × 3) and the combined organics were washed with brine, dried with MgSO<sub>4</sub> and concentrated. Recrystallization using methylene chloride and *n*-hexane gave **3a** (74.0 mg, 96%) as a white solid. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>, 25 °C, TMS) δ 7.40-7.28 (m, 3H), 7.18-7.15 (m, 2H), 5.52 (s, 1H), 5.47 (s, 1H), 4.06 (d, *J* = 9.34 Hz, 1H), 3.56 (ddd, *J* = 9.58, 6.66, 3.19 Hz, 1H), 2.94 (dd, *J* = 16.92,

2.80 Hz, 1H), 2.78-2.66 (m, 1H), 1.82 (d,  $J = 1.54$  Hz, 3H);  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  173.3, 171.3, 141.4, 135.5, 135.4, 129.6, 128.4, 128.1, 127.4, 116.2, 46.2, 40.0, 29.4, 17.0; IR (film) 2951, 1731, 1434, 1241, 1060, 912, 818, 759  $\text{cm}^{-1}$ ; m.p. 108-109 °C; HRMS (EI) calcd for  $\text{C}_{16}\text{H}_{14}\text{O}_3$   $\text{M}^+$  254.0943, found 254.0945.

**5-Ethyl-4-methylene-6-phenyl-3a,4,7,7a-tetrahydroisobenzofuran-1,3-dione (3b):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.38-7.34 (m, 2H), 7.31-7.28 (m, 1H), 7.14-7.11 (m, 1H), 5.50 (d,  $J = 0.7$  Hz, 1H), 5.45 (s, 1H), 4.04 (d,  $J = 9.51$  Hz, 1H), 3.56 (ddd,  $J = 9.39, 6.56, 2.70$  Hz, 1H), 2.87 (dd,  $J = 16.24, 2.81$  Hz, 1H), 2.67 (ddd,  $J = 16.25, 6.56, 2.23$  Hz, 1H), 2.40-2.31 (m, 1H), 2.23-2.14 (m, 1H), 0.88 (t,  $J = 7.40$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  173.5, 171.3, 141.3, 137.0, 135.0, 134.4, 128.5, 127.6, 127.3, 116.0, 47.7, 40.3, 29.8, 22.9, 13.3; IR (film) 1844, 1781, 1700, 904  $\text{cm}^{-1}$ ; m.p. 135-136 °C; MS (EI)  $m/z$  268 ( $\text{M}^+$ ).

**5-Ethyl-4-methylene-2,6-diphenyl-3a,4,7,7a-tetrahydroisindole-1,3-dione (3c):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.48-7.45 (m, 2H), 7.40-7.32 (m, 3H), 7.27-7.24 (m, 3H), 7.17-7.14 (m, 2H), 5.50 (s, 1H), 5.40 (s, 1H), 3.93 (d,  $J = 8.87$  Hz, 1H), 3.45-3.41 (m, 1H), 3.00 (dd,  $J = 15.73, 2.27$  Hz, 1H), 2.67 (ddd,  $J = 15.70, 6.39, 2.37$  Hz, 1H), 2.48-2.39 (m, 1H), 2.26-2.17 (m, 1H), 0.85 (t,  $J = 7.25$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  178.5, 176.3, 141.7, 137.2, 137.1, 134.7, 132.1, 129.2, 128.6, 128.3, 127.9, 127.0, 126.3, 115.1, 48.4, 39.9, 30.4, 23.0, 13.6; IR (film) 1781, 1715, 1381, 1175  $\text{cm}^{-1}$ ; m.p. 141-142 °C; MS (EI)  $m/z$  343 ( $\text{M}^+$ ).

**Dimethyl 4-ethyl-3-methylene-5-phenyl-cyclohex-4-ene-1,2-dicarboxylate (3d):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.36-7.31 (m, 2H), 7.28-7.24 (m, 1H), 7.15-7.12 (m, 2H), 5.30 (s, 1H), 5.23 (s, 1H), 3.93 (d,  $J = 3.56$  Hz, 1H), 3.72 (s, 3H), 3.69 (s, 3H), 3.05-2.94 (m, 2H), 2.69-2.58 (m, 1H), 2.21-2.05 (m, 2H), 0.91 (t,  $J = 7.43$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.5, 172.1, 142.9, 139.5, 137.2, 133.0, 128.3, 128.2, 127.7, 126.7, 113.9, 52.0, 48.6, 41.1, 32.3, 22.1, 13.7; IR (film) 2952, 1739, 1205  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  314 ( $\text{M}^+$ ).

**2-Ethyl-1-methylene-3-phenyl-1,4,4a,9a-tetrahydroanthraquinone (3e):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  8.10-8.07 (m, 1H), 8.05-8.03 (m, 1H), 7.78-7.72 (m, 2H), 7.38-7.33 (m, 2H), 7.29-7.25 (m, 3H), 5.31 (s, 1H), 4.88 (s, 1H), 4.11 (dt,  $J = 3.61, 1.41$  Hz, 1H), 3.58 (q,  $J = 5.56$  Hz, 1H), 2.95 (dd,  $J = 18.08, 5.97$  Hz, 1H), 2.61 (dd,  $J = 18.06, 5.45$  Hz, 1H), 2.17-2.11 (m, 2H), 0.92 (t,  $J = 7.45$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  197.0, 196.9, 142.9, 137.1, 136.4, 135.1, 134.4, 134.3, 134.1, 133.8, 128.3, 127.7, 126.9, 126.8, 127.7, 113.7, 54.8, 47.8, 32.8, 22.1, 14.0; IR (film) 3060, 1695, 1594, 703  $\text{cm}^{-1}$ ; m.p. 101-102 °C; MS (EI)  $m/z$  328 ( $\text{M}^+$ ).

**Ethyl 4-ethyl-3-methylene-5-phenyl-3,6-dihydro-2H-pyran-2-carboxylate (3f):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.38-7.28 (m, 3H), 7.18-7.15 (m, 2H), 5.32 (s, 1H), 5.15 (s, 1H), 4.88 (s, 1H), 4.71 (d,  $J = 17.13$  Hz, 1H), 4.37 (d,  $J = 17.13$  Hz, 1H), 4.28 (q,  $J = 7.11$  Hz, 2H), 2.19 (q,  $J = 7.29$  Hz, 2H), 1.32 (t,  $J = 7.17$  Hz, 3H), 0.97 (t,  $J =$

7.47 Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  170.5, 138.1, 136.0, 135.6, 131.3, 128.4, 128.3, 127.5, 111.1, 68.0, 61.2, 21.2, 14.3, 13.8; IR (film) 2975, 1739, 1609, 1465, 1130  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  199 ( $\text{M}^+ - \text{CO}_2\text{Et}$ ).

**Ethyl 4-ethyl-5-methylene-3-phenylcyclohex-3-ene-carboxylate (3g, major compound):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.35 (m, 2H), 7.27 (m, 1H), 7.16 (d,  $J = 6.83$  Hz, 2H), 5.09 (s, 1H), 4.97 (s, 1H), 4.14 (q,  $J = 7.11$  Hz, 2H), 2.83-2.77 (m, 1H), 2.71 (dd,  $J = 14.1, 3.46$  Hz, 1H), 2.65-2.54 (m, 3H), 2.18-2.07 (m, 2H), 1.25 (t,  $J = 7.16$  Hz, 3H), 0.93 (t,  $J = 7.4$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  174.8, 143.4, 141.0, 136.6, 134.6, 128.2, 127.7, 126.6, 110.1, 60.4, 40.2, 35.8, 35.1, 21.9, 14.3, 14.2; IR (film) 2932, 1732, 1442, 1178, 1038, 884, 761, 701  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  270 ( $\text{M}^+$ ).

**Ethyl 3-ethyl-2-methylene-4-phenylcyclohex-3-ene-carboxylate (3g, minor compound):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.33 (t,  $J = 7.38$  Hz, 2H), 7.24 (m, 1H), 7.12 (m, 2H), 5.22 (s, 1H), 4.98 (s, 1H), 4.19 (m, 2H), 3.40 (t,  $J = 5.18$  Hz, 1H), 2.49 (m, 1H), 2.33 (dt,  $J = 18.4, 5.20$  Hz, 1H), 2.25-2.12 (m, 3H), 1.98 (m, 1H), 1.28 (t,  $J = 7.10$  Hz, 3H), 0.95 (t,  $J = 7.44$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.9, 143.8, 140.5, 137.9, 133.7, 128.1, 127.6, 126.4, 111.1, 60.4, 47.5, 31.1, 25.7, 22.2, 14.3, 13.9; IR (film) 2931, 1732, 1605, 1443, 1373, 1308, 1258, 1155, 1040, 759  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  270 ( $\text{M}^+$ ).

**5-Ethyl-4-methylene-3a,7,8,9,9a,9b-hexahydro-4H-naphtho[1,2-c]furan-1,3,6-trione (3h):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  5.67 (s, 1H), 5.63 (s, 1H), 4.02 (d,  $J = 9.52$  Hz, 1H), 3.46 (dd,  $J = 9.50, 5.15$  Hz, 1H), 3.15 (octet,  $J = 7.27, 12.02, 14.68$  Hz, 1H), 2.68-2.62 (m, 1H), 2.57-2.48 (m, 1H), 2.46-2.35 (m, 1H), 2.33-2.03 (m, 2H), 2.09-1.97 (m, 2H), 1.76-1.63 (m, 1H), 1.00 (t,  $J = 7.39$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  199.2, 171.0, 170.4, 153.3, 137.6, 131.2, 121.0, 49.7, 44.7, 40.0, 37.2, 25.8, 25.3, 21.2, 13.1; IR (film) 1885, 1780, 1662, 1549, 1200  $\text{cm}^{-1}$ ; m.p. 156 °C; MS (EI)  $m/z$  260 ( $\text{M}^+$ ).

**5-Ethyl-4-methylene-2-phenyl-3a,7,8,9,9a,9b-hexahydro-4H-benzo[e]isindole-1,3,6-trione (3i):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  7.65-7.32 (m, 3H), 7.15-7.13 (m, 2H), 5.63 (s, 1H), 5.62 (s, 1H), 3.90 (d,  $J = 9.04$  Hz, 1H), 3.46 (dd,  $J = 4.92, 8.87$  Hz, 1H), 3.19 (octet,  $J = 7.29, 11.72, 14.63$  Hz, 1H), 2.72-2.67 (m, 1H), 2.54-2.33 (m, 3H), 2.28-2.17 (m, 1H), 2.08-1.99 (m, 2H), 1.76-1.62 (m, 1H), 1.00 (t,  $J = 7.33$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ):  $\delta$  199.9, 176.4, 175.0, 153.5, 139.7, 131.7, 131.1, 129.2, 128.7, 126.4, 120.0, 49.9, 43.7, 40.0, 38.2, 25.8, 25.5, 21.4, 13.4; IR (film) 1713, 1668, 1498  $\text{cm}^{-1}$ ; m.p. 160-161 °C; MS (EI)  $m/z$  335 ( $\text{M}^+$ ).

**Ethyl 4-ethyl-3-methylene-5-oxo-3,5,6,7,8,8a-hexahydro-2H-chromene-2-carboxylate (3j):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25 °C, TMS)  $\delta$  5.56 (s, 1H), 5.27 (s, 1H), 4.74 (t,  $J = 1.53$  Hz, 1H), 4.42-4.30 (m, 3H), 2.59-2.28 (m, 5H), 2.07-2.00 (m, 1H), 1.92-1.82 (m, 1H), 1.75-1.63 (m, 1H), 1.35 (t,  $J = 7.10$  Hz, 3H), 1.13 (t,  $J = 7.45$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  201.8, 169.2, 142.6, 137.3, 135.4, 114.1, 76.8, 61.5, 42.7, 31.5, 21.8, 19.2, 14.4, 14.2; IR (film) 2940,

1735, 1685, 1457, 1192  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  264 ( $\text{M}^+$ ).

**8-tert-Butyl-4-methylene-2-phenyl-3a,4,6,7,8,9,9a,9b-octahydrobenzo[e]isoindole-1,3-dione (3k, major compound):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25  $^\circ\text{C}$ , TMS)  $\delta$  7.44 (t,  $J = 7.70$  Hz, 2H), 7.36 (t,  $J = 7.47$  Hz, 1H), 7.23 (d,  $J = 7.56$  Hz, 2H), 6.04 (s, 1H), 5.30 (s, 1H), 5.17 (s, 1H), 3.80 (d,  $J = 8.30$  Hz, 1H), 3.36 (dd,  $J = 8.01, 6.86$  Hz, 1H), 2.77 (m, 1H), 2.37 (m, 1H), 2.25 (m, 1H), 2.02 (td,  $J = 12.65, 8.87$  Hz, 1H), 1.73 (m, 2H), 1.52 (m, 1H), 1.32 (qd,  $J = 12.48, 5.41$  Hz, 1H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  175.8, 175.0, 142.1, 133.0, 130.9, 128.0, 127.4, 125.3, 121.1, 113.5, 43.5, 42.2, 41.4, 32.3, 31.1, 30.1, 26.0, 24.8, 21.5; IR (film) 3463, 2959, 2867, 2249, 1775, 1709  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  349 ( $\text{M}^+$ ).

**4-Methoxy-12-methylene-16-phenyl-6,7,8,12,13,14-hexahydro-16-aza-cyclopenta[a]phenanthrene-15,17-dione (3l):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25  $^\circ\text{C}$ , TMS)  $\delta$  7.41 (m, 2H), 7.33 (m, 1H), 7.23 (m, 3H), 7.14 (t,  $J = 8.05$  Hz, 1H), 6.85 (d,  $J = 2.25$  Hz, 1H), 6.75 (d,  $J = 8.02$  Hz, 1H), 5.52 (s, 1H), 5.35 (s, 1H), 3.92 (d,  $J = 8.09$  Hz, 1H), 3.83 (s, 3H), 3.50 (dd,  $J = 8.07, 5.44$  Hz, 1H), 3.27 (td,  $J = 16.33, 3.32$  Hz, 1H), 2.88 (m, 1H), 2.55 (qd,  $J = 13.06, 3.81$  Hz, 1H), 2.33 (m, 1H), 2.13 (m, 1H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  175.0, 174.8, 155.8, 135.6, 134.8, 133.0, 130.9, 127.9, 127.3, 126.4, 125.5, 125.3, 121.1, 116.1, 114.7, 108.1, 54.5, 45.0, 43.3, 34.3, 23.0, 21.8. IR (film) 2834, 1710, 1572, 1499, 1387, 1261  $\text{cm}^{-1}$ ; m.p. 189-190  $^\circ\text{C}$ ; MS (EI)  $m/z$  371 ( $\text{M}^+$ ).

**Ethyl 4-ethyl-5-methyl-3-phenyl benzoate (4a):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25  $^\circ\text{C}$ , TMS)  $\delta$  7.84 (d,  $J = 1.33$  Hz, 1H), 7.70 (d,  $J = 1.69$  Hz, 1H), 7.46-7.37 (m, 3H), 7.30-7.27 (m, 2H), 4.35 (q,  $J = 7.09$  Hz, 2H), 2.60 (q,  $J = 7.52$  Hz, 2H), 2.44 (s, 3H), 1.36 (t,  $J = 7.18$  Hz, 3H), 0.99 (t,  $J = 7.51$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  166.8, 145.6, 142.4, 141.8, 136.6, 130.5, 129.2, 129.0, 128.0, 127.3, 127.0, 60.8, 23.3, 19.7, 14.4, 14.1; IR (film) 2971, 2930, 1717, 1223  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  268 ( $\text{M}^+$ ).

**Dimethyl 3,4-dimethyl-5-phenylphthalate (4b):** A mixture of 3-methyl-4-phenyl-1,2,4-pentatriene (**1a**) (46.86 mg, 0.3 mmol) and DMAD (42.63 mg, 0.3 mmol) was heated in toluene (1.0 mL) at 100  $^\circ\text{C}$  for 4 h. The reaction mixture was quenched with  $\text{NaHCO}_3$  (sat. aq.). The aqueous layer was extracted with  $\text{CH}_2\text{Cl}_2$  (20 mL  $\times$  3) and the combined organics were washed with water and brine, dried over  $\text{MgSO}_4$ , filtered, and concentrated under reduced pressure. It was purified by recrystallization ( $\text{CH}_2\text{Cl}_2$ :Hex) to give **4b** (73.0 mg, 82%).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ , 25  $^\circ\text{C}$ , TMS)  $\delta$  7.78 (s, 1H), 7.47-7.36 (m, 3H), 7.28-7.25 (m, 2H), 3.99 (s, 3H), 3.86 (s, 3H), 2.30 (s, 3H), 2.21 (s, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  170.5, 166.1, 143.1, 141.0, 140.4, 134.6, 134.4, 129.2, 129.1, 128.2, 127.3, 124.3, 52.6, 52.3, 17.8,

17.1; IR (film) 1730, 1432, 1323  $\text{cm}^{-1}$ ; m.p. 107-108  $^\circ\text{C}$ ; MS (EI)  $m/z$  298 ( $\text{M}^+$ ).

**Dimethyl 4-ethyl-3-methyl-5-oxo-5,6,7,8-tetrahydronaphthalene-1,2-dicarboxylate (4c):**  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ , 25  $^\circ\text{C}$ , TMS)  $\delta$  3.90 (s, 3H), 3.87 (s, 3H), 2.99 (q,  $J = 7.34$  Hz, 2H), 2.96 (t,  $J = 6.26$  Hz, 2H), 2.67 (t,  $J = 6.75$  Hz, 2H), 2.33 (s, 3H), 2.04 (quintet,  $J = 6.50$  Hz, 2H), 1.20 (t,  $J = 7.33$  Hz, 3H);  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  200.1, 169.0, 168.1, 148.4, 141.1, 136.4, 133.7, 133.4, 128.5, 52.6, 52.5, 40.7, 28.1, 23.9, 22.3, 15.6, 14.1; IR (film) 2953, 2359, 1739, 1690, 1436  $\text{cm}^{-1}$ ; MS (EI)  $m/z$  304 ( $\text{M}^+$ ).

**Acknowledgment.** This work was supported by KOSEF through the National Research Lab. Program funded by the Ministry of Science and Technology (No. M10600000203-06J0000-20310), by the Korea Science and Engineering Foundation (KOSEF, R01-2006-000-11283-0), and by the CMDS at KAIST. The NMR and mass data were obtained from the central instrumental facility in Kangwon National University.

## References

- (a) Patai, S., Ed. *The Chemistry of Ketenes, Allenes and Related Compounds*; Wiley: New York, 1980. (b) *The Chemistry of the Allenes*; Landor, S. R., Ed.; Academic: London, 1982. (c) Smadja, W. *Chem. Rev.* **1983**, *83*, 263. (d) Schuster, H. F.; Coppola, G. M. *Allenenes in Organic Synthesis*; Wiley: New York, 1984. (e) Zimmer, R.; Dinesh, C. U.; Nandan, E.; Khan, F. A. *Chem. Rev.* **2000**, *100*, 3067. (f) *Modern Allene Chemistry*; Krause, N., Hashmi, A. S. K., Eds.; Wiley-VCH: Weinheim, 2004.
- (a) Angelov, C. M.; Mondeshka, D. M.; Tancheva, T. N. *Chem. Commun.* **1985**, 647. (b) Reich, H. J.; Eisenhart, E. K.; Whipple, W. L.; Kelly, M. J. *J. Am. Chem. Soc.* **1988**, *110*, 6432. (c) Dulcere, J. P.; Agati, V.; Faure, R. *Chem. Commun.* **1993**, 270. (d) Spino, C.; Thibault, C.; Gingras, S. *J. Org. Chem.* **1998**, *63*, 5283. (e) Regás, D.; Afonso, M. M.; Rodríguez, M. L.; Palenzuela, J. A. *J. Org. Chem.* **2003**, *68*, 7845. (f) Regás, D.; Ruiz, J. M.; Afonso, M. M.; Palenzuela, J. A. *J. Org. Chem.* **2006**, *71*, 9153.
- (a) Murakami, M.; Itami, K.; Ito, Y. *Angew. Chem. Int. Ed.* **1995**, *34*, 2691. (b) Murakami, M.; Itami, K.; Ito, Y. *J. Am. Chem. Soc.* **1997**, *119*, 7163. (c) Lee, P. H.; Lee, K. *Angew. Chem. Int. Ed.* **2005**, *44*, 3253. (d) Lee, P. H.; Lee, K.; Kang, Y. *J. Am. Chem. Soc.* **2006**, *128*, 1139. (e) Funami, H.; Kusama, H.; Iwasawa, N. *Angew. Chem. Int. Ed.* **2007**, *46*, 909.
- (a) Cavallito, C. J.; Fruehauf, D. M.; Bailey, J. H. *J. Am. Chem. Soc.* **1948**, *70*, 3724. (b) Lee, K. H.; Hall, I. H.; Mar, E. C.; Starnes, C. O.; El Gebaly, S. A.; Waddell, T. G.; Hadgraft, R. I.; Ruffner, C. G.; Weidner, I. *Science* **1977**, *196*, 533. (c) Spring, O.; Albert, K.; Gradmann, W. *Phytochemistry* **1981**, *20*, 1883. (d) Park, B. K.; Nakagawa, M.; Hirota, A.; Nakayama, M. *J. Antibiot.* **1988**, *41*, 751. (e) Liu, S.; Liu, H.; Yan, W.; Zhang, L.; Bai, N.; Ho, C.-T. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1101.
- Lee, K.; Seomoon, D.; Lee, P. H. *Angew. Chem. Int. Ed.* **2002**, *41*, 3901.