

Pd-Mediated Cross-Coupling Reactions between the Bromide of Baylis-Hillman Adduct and Organostannanes

Saravanan Gowrisankar, Sung Hwan Kim, and Jae Nyong Kim*

Department of Chemistry and Institute of Basic Science, Chonnam National University, Gwangju 500-757, Korea

*E-mail: kimjn@chonnam.ac.kr

Received September 20, 2008, Accepted January 14, 2009

Key Words: Palladium, Cross-coupling, Baylis-Hillman adduct, Organostannanes


Recently numerous chemical transformations of Baylis-Hillman adducts have been published involving the synthesis of cyclic and acyclic compounds.¹ Among the reactions introduction of allyl,² vinyl³ and aryl^{4,5} moiety at the primary position of the Baylis-Hillman adducts can be regarded as an important transformation due to the usefulness of the compounds thereof as synthetic intermediates.²⁻⁵

Introduction of allyl group at the primary position of Baylis-Hillman adduct was studied by Yadav and coworkers.^{2a,b} They reported the synthesis of 1,5-diene derivatives involving the use of *in-situ* generated allylzinc reagent and Baylis-Hillman acetates.^{2a} Later, they reported another method, the reaction of allyltrimethylsilane and Baylis-Hillman alcohol in the presence of BF₃ etherate.^{2b} Introduction of alkenyl moiety at the primary position of Baylis-Hillman adduct has not been reported much.³ Isobutenyl and isopentenyl groups were introduced using the corresponding Grignard reagents and used for the synthesis of dihydronaphthalenes.^{3a} Very recently Ranu and coworkers published an elegant method, a Pd-mediated introduction of aryl and vinyl moiety to Baylis-Hillman acetate, using triorganoindium reagents.^{3b} Introduction of aryl group at the primary position of the Baylis-Hillman adduct has been studied relatively well and carried out by using Friedel-Crafts type reaction under various acidic conditions.⁴ However, regioselectivity is the principle problem in this case.^{4c,d} As an example, Friedel-Crafts reaction with toluene produced a mixture of *ortho*- and *para*-isomers (*vide infra*, Scheme 2). In addition, various Pd- and Rh-catalyzed introduction of aryl


moiety has been studied.⁵ Kabalka group reported a Pd-catalyzed cross-coupling of Baylis-Hillman acetate and organosilane^{5a} or potassium organotrifluoroborate.^{5b} Genet and coworkers examined a Rh-catalyzed arylation with arylboronic acid^{5c} or potassium aryltrifluoroborate.^{5d}

Although various methods have been reported and most of the methods provided moderate yields of products,²⁻⁵ development of an efficient and general method for the introduction of carbon nucleophile at the primary position of Baylis-Hillman adduct is still highly required, especially in a stereoselective manner. Thus we decided to examine the cross-coupling reactions between the bromide of Baylis-Hillman adduct and allyl-, aryl-, and vinylstannane (Scheme 1) as a continuous work on our recent Pd-mediated reactions with Baylis-Hillman adducts.⁶ Although cross-couplings involving the use of allyl bromides and organostannanes have been studied extensively,⁷ this is the first trial in the Baylis-Hillman chemistry to the best of our knowledge. Fortunately we obtained good results and wish to report herein the results. The results are summarized in Scheme 1.


We decided to use the bromides of Baylis-Hillman adducts and chose three representative Baylis-Hillman adducts, **2a-c**.⁸ The starting materials **2a-c** were prepared in pure form (*Z* for **2a** and **2b**, *E* for **2c**) as shown in Scheme 1.^{9,10} The reaction of cinnamyl bromide **2a** and allyltributylstannane (1.2 equiv) in the presence of Pd(Ph₃P)₄/LiCl (0.5 equiv) in THF afforded desired product **3a** in 96% yield.^{2a,7d,11} As a palladium catalyst, Pd(Ph₃P)₄ was superior than Pd(OAc)₂ and only 2



Scheme 1



Scheme 2



Scheme 3

mol% of catalyst was sufficient. The reaction time can be shortened by raising the reaction temperature with 5 mol% of Pd(0) catalyst. Similarly, **3b** and **3c** were prepared in high yields also (81-93%). The arylation and vinylation of **2a-c** were examined with phenyltributylstannane, 2-(tributylstannyl)furan, and tetravinylstannane. The corresponding products **4a-c**, **5a-c**, and **6a-c** were obtained in good yields (73-95%). The yields of nitrile-containing compounds were comparatively lower than those of the ester- or acetyl-containing compounds. The reaction of phenyltributylstannane required elevated temperature (50 °C).

Tolyl-substituted products **4d** and **4e** were synthesized in good yields (74-78%) similarly by the reactions of **2a** and *ortho*-tolyltributylstannane and *para*-tolyltributylstannane which were prepared from Pd-mediated reactions between *n*-tributyltin hydride and 2-iodo- and 4-iodotoluene, respectively,¹² while the traditional Friedel-Crafts reaction of Baylis-Hillman adduct **1a** and toluene under the influence of H₂SO₄ (0.2 equiv) afforded *ortho/para* mixture in 74% yield (55:45 ratio in ¹H NMR) as in Scheme 2.

As a next trial we prepared tributylstannyl derivative **7a** by the Pd-mediated reaction of **2a** and hexabutyltin in the presence of Pd(0) in 66% yield as in Scheme 3.^{13,14} The reaction of **7a** and allyl bromide under the influence of Pd(Ph₃P)₄/LiCl produced **3a** in low yield (48%). Similarly the reaction of **7a** and bromobenzene afforded **4a** in low yield (43%) also together with some self-coupling product **8a** (10%).¹⁵

In summary, we disclosed an efficient synthetic method of allyl-, aryl- and vinyl-attached Baylis-Hillman adducts *via* the Pd-mediated cross-coupling reactions between the bromide of Baylis-Hillman adduct and the corresponding organostannane compounds.

Experimental Section

Typical experimental procedure for the preparation of **3a**.

To a stirred mixture of **2a**¹⁰ (128 mg, 0.5 mmol) and allyltributylstannane (200 mg, 0.6 mmol) in dry THF (2 mL) was added Pd(PPh₃)₄ (12 mg, 2 mol%) and LiCl (11 mg, 0.25 mmol) and the reaction mixture was stirred at room temperature for 12 h. After removal of solvent and column chromatographic purification process (hexanes/EtOAc, 95:5) analytically pure product **3a** was obtained as colorless oil,^{2a} 104 mg (96%). Other compounds were synthesized similarly and the known compounds, **3a**,^{2a} **4a**,^{5b} **4b**,^{5a} **4c**,^{5e} **4d**,^{5b} **4e**,^{5b} **6a**,^{3b} **6c**¹⁶ and **8a**¹⁵ were identified by comparison with their ¹H NMR and IR spectroscopic data with the reported. The spectroscopic data of unknown compounds, **3b**, **3c**, **5a**, **5b**, **5c**, **6b** and **7a**, are as follows.

Compound **3b**: 93%; colorless oil; IR (film) 3077, 2927, 1667 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.20-2.24 (m, 2H), 2.45 (s, 3H), 2.60-2.63 (m, 2H), 4.95-5.05 (m, 2H), 5.78-5.87 (m, 1H), 7.34-7.43 (m, 5H), 7.51 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 25.61, 26.07, 33.00, 114.77, 128.56, 128.60, 129.20, 135.57, 137.88, 140.02, 142.06, 200.06.

Compound **3c**: 81%; colorless oil; IR (film) 3080, 2924, 2209 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.37-2.54 (m, 4H), 5.04-5.16 (m, 2H), 5.75-5.88 (m, 1H), 6.94 (s, 1H), 7.26-7.44 (m, 3H), 7.69-7.74 (m, 2H); ¹³C NMR (CDCl₃, 75 MHz) δ 32.27, 35.64, 110.66, 116.45, 118.65, 128.55, 128.78, 129.93, 133.67, 136.04, 143.84.

Compound **5a**: 84%; yellow oil; IR (film) 2951, 1715 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 3.80 (s, 3H), 3.88 (s, 2H), 6.06-6.08 (m, 1H), 6.30-6.31 (m, 1H), 7.33-7.47 (m, 6H), 7.87 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 26.89, 52.15, 106.00, 110.39, 129.23, 128.55, 128.89, 129.33, 134.99, 141.23, 141.30, 152.96, 168.22.

Compound **5b**: 86%; yellow oil; IR (film) 2957, 2924, 1669 cm⁻¹; ¹H NMR (CDCl₃, 300 MHz) δ 2.46 (s, 3H), 3.87 (s, 2H), 6.00-6.02 (m, 1H), 6.27-6.29 (m, 1H), 7.32-7.49 (m, 6H), 7.68 (s, 1H); ¹³C NMR (CDCl₃, 75 MHz) δ 25.78, 26.05, 106.01, 110.37, 128.60, 129.04, 129.36, 134.99, 137.45, 141.14, 141.45, 153.04, 199.13.

Compound **5c**: 82%; yellow oil; IR (film) 2913, 2212 cm^{-1} ; ^1H NMR (CDCl_3 , 500 MHz) δ 3.73 (s, 2H), 6.26-6.27 (m, 1H), 6.35-6.36 (m, 1H), 6.99 (s, 1H), 7.38-7.43 (m, 5H), 7.72-7.74 (m, 2H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 34.58, 107.60, 108.00, 110.63, 118.33, 128.73, 128.80, 130.28, 133.34, 142.46, 144.84, 149.82.

Compound **6b**: 93%; colorless oil; IR (film) 3079, 2978, 1668 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz) δ 2.47 (s, 3H), 3.26-3.29 (m, 2H), 4.99-5.11 (m, 2H), 5.91-6.03 (m, 1H), 7.32-7.46 (m, 5H), 7.64 (s, 1H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 26.05, 30.57, 115.51, 128.48, 128.82, 129.28, 135.34, 135.73, 139.20, 141.03, 199.54.

Compound **7a**: 66%; colorless oil; IR (film) 2956, 2925, 1709 cm^{-1} ; ^1H NMR (CDCl_3 , 300 MHz) δ 0.84-0.94 (m, 15H), 1.23-1.31 (m, 6H), 1.37-1.46 (m, 6H), 2.29 (s, 2H), 3.80 (s, 3H), 7.35-7.37 (m, 5H); ^{13}C NMR (CDCl_3 , 75 MHz) δ 10.22, 11.80, 13.67, 27.33, 28.95, 51.98, 127.47, 128.30, 129.25, 131.78, 133.89, 136.88, 169.46; ESIMS m/z 465 (M^+ +1). Anal. Calcd for $\text{C}_{23}\text{H}_{38}\text{O}_2\text{Sn}$: C, 59.37; H, 8.23. Found: C, 59.02; H, 8.44.

Acknowledgments. This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD, KRF-2007-313-C00417). Spectroscopic data was obtained from the Korea Basic Science Institute, Gwangju branch.

References and Notes

- For general review on Baylis-Hillman reaction, see: (a) Basavaiah, D.; Rao, A. J.; Satyanarayana, T. *Chem. Rev.* **2003**, *103*, 811-891. (b) Ciganek, E. In *Organic Reactions*; Paquette, L. A., Ed.; John Wiley & Sons: New York, 1997; Vol. 51, pp 201-350. (c) Basavaiah, D.; Rao, P. D.; Hyma, R. S. *Tetrahedron* **1996**, *52*, 8001-8062. (d) Kim, J. N.; Lee, K. Y. *Curr. Org. Chem.* **2002**, *6*, 627-645. (e) Lee, K. Y.; Gowrisankar, S.; Kim, J. N. *Bull. Korean Chem. Soc.* **2005**, *26*, 1481-1490. (f) Singh, V.; Batra, S. *Tetrahedron* **2008**, *64*, 4511-4574 and further references cited therein.
- For the synthesis of allylated Baylis-Hillman adducts, see: (a) Srihari, P.; Singh, A. P.; Basak, A. K.; Yadav, J. S. *Tetrahedron Lett.* **2007**, *48*, 5999-6001. (b) Yadav, J. S.; Reddy, B. V. S.; Mandal, S. S.; Singh, A. P.; Basak, A. K. *Synthesis* **2008**, 1943-1947. (c) Yadav, J. S.; Reddy, B. V. S.; Singh, A. P.; Basak, A. K. *Synthesis* **2008**, 469-473. (d) Mandal, S. K.; Paira, M.; Roy, S. C. *J. Org. Chem.* **2008**, *73*, 3823-3827.
- For the synthesis of vinyl moiety-containing Baylis-Hillman adducts, see: (a) Gowrisankar, S.; Lee, C. G.; Kim, J. N. *Tetrahedron Lett.* **2004**, *45*, 6949-6953. (b) Ranu, B. C.; Chattopadhyay, K.; Jana, R. *Tetrahedron Lett.* **2007**, *48*, 3847-3850.
- For the synthesis of arylated Baylis-Hillman adducts by Friedel-Crafts reaction, see: (a) Basavaiah, D.; Pandiaraju, S.; Padmaja, K. *Synlett* **1996**, 393-395. (b) Basavaiah, D.; Krishnamacharyulu, M.; Hyma, R. S.; Pandiaraju, S. *Tetrahedron Lett.* **1997**, *38*, 2141-2144. (c) Lee, H. J.; Seong, M. R.; Kim, J. N. *Tetrahedron Lett.* **1998**, *39*, 6223-6226. (d) Das, B.; Majhi, A.; Banerjee, J.; Chowdhury, N.; Venkateswarlu, K. *Chem. Lett.* **2005**, *34*, 1492-1493. (e) Shanmugam, P.; Rajasingh, P. *Chem. Lett.* **2005**, *34*, 1494-1495.
- For the synthesis of arylated Baylis-Hillman adducts using metal catalyst, see: (a) Kabalka, G. W.; Dong, G.; Venkataiah, B.; Chen, C. *J. Org. Chem.* **2005**, *70*, 9207-9210. (b) Kabalka, G. W.; Venkataiah, B.; Dong, G. *Org. Lett.* **2003**, *5*, 3803-3805. (c) Navarre, L.; Darses, S.; Genet, J.-P. *Chem. Commun.* **2004**, 1108-1109. (d) Navarre, L.; Darses, S.; Genet, J.-P. *Adv. Synth. Catal.* **2006**, *348*, 317-322. (e) Kantam, M. L.; Kumar, K. B. S.; Sreedhar, B. *J. Org. Chem.* **2008**, *73*, 320-322.
- For our recent contributions on Pd-mediated reactions of modified Baylis-Hillman adducts, see: (a) Gowrisankar, S.; Lee, H. S.; Kim, J. M.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 1670-1673. (b) Kim, J. M.; Kim, K. H.; Kim, T. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 3248-3251. (c) Gowrisankar, S.; Lee, H. S.; Lee, K. Y.; Lee, J.-E.; Kim, J. N. *Tetrahedron Lett.* **2007**, *48*, 8619-8622. (d) Lee, H. S.; Kim, S. H.; Kim, T. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 1773-1776. (e) Kim, H. S.; Gowrisankar, S.; Kim, S. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 3858-3861. (f) Lee, H. S.; Kim, S. H.; Gowrisankar, S.; Kim, J. N. *Tetrahedron* **2008**, *64*, 7183-7190.
- For the Pd-mediated cross-coupling reactions between allylic bromides (and acetates) and organostannanes, see: (a) Godschalx, J.; Stille, J. K. *Tetrahedron Lett.* **1980**, *21*, 2599-2602. (b) Trost, B. M.; Keinan, E. *Tetrahedron Lett.* **1980**, *21*, 2595-2598. (c) Trost, B. M.; Pietrusiewicz, K. M. *Tetrahedron Lett.* **1985**, *26*, 4039-4042. (d) Mendez, M.; Cuerva, J. M.; Gomez-Bengoa, E.; Cardenas, D. J.; Echavarren, A. M. *Chem. Eur. J.* **2002**, *8*, 3620-3628.
- At the initial stage of this work, the reaction of Baylis-Hillman alcohol **1a** and allyltributylstannane was examined, however, we did not observe the formation of **3a**. The reaction of the acetate of **1a** was also examined. Although the reaction rate of Baylis-Hillman acetate was faster than that of the bromide, appreciable amounts of side products including **3a-Z** (ca. 5%) and secondary adduct (ca. 5%) was observed.
- For the synthesis of cinnamyl bromide derivatives in a stereoselective manner, see: (a) Das, B.; Banerjee, J.; Ravindranath, N. *Tetrahedron* **2004**, *60*, 8357-8361. (b) Fernandes, L.; Bortoluzzi, A. J.; Sa, M. M. *Tetrahedron* **2004**, *60*, 9983-9989. (c) Sa, M. M.; Ramos, M. D.; Fernandes, L. *Tetrahedron* **2006**, *62*, 11652-11656. (d) Deng, J.; Hu, X.-P.; Huang, J.-D.; Yu, S.-B.; Wang, D.-Y.; Duan, Z.-C.; Zheng, Z. *J. Org. Chem.* **2008**, *73*, 2015-2017. (e) Buchholz, R.; Hoffmann, H.; Martin, R. *Helv. Chim. Acta* **1991**, *74*, 1213-1220. (f) Mazdiyasi, H.; Konopacki, D. B.; Dickman, D. A.; Zydowsky, T. M. *Tetrahedron Lett.* **1993**, *34*, 435-438. (g) Basavaiah, D.; Reddy, K. R.; Kumaragurubaran, N. *Nature Protocols* **2007**, *2*, 2665-2676. (h) Lee, K. Y.; Lee, Y. J.; Kim, J. N. *Bull. Korean Chem. Soc.* **2007**, *28*, 143-146. (i) Lee, K. Y.; Park, D. Y.; Kim, J. N. *Bull. Korean Chem. Soc.* **2006**, *27*, 1489-1492.
- Treatment of **1a** and **1b** with 48% aqueous HBr at room temperature afforded **2a** and **2b** in 81 and 76%, respectively. The stereochemistry of **2a** and **2b** was found to be very pure (*Z*-form, > 97% by ^1H NMR). The reaction of **1c** and HBr afforded **2c** as a mixture of *E/Z* isomers (84%, *E/Z* = 4:1). Thus, we first converted **1c** into the corresponding primary alcohol with aqueous H_2SO_4 (63%),^{10a} and synthesized **2c** by treatment with PBr_3 (51%, ca. 95% *E* by ^1H NMR). For the stereoselective synthesis of primary alcohol, see: (a) Basavaiah, D.; Kumaragurubaran, N.; Padmaja, K. *Synlett* **1999**, 1630-1632.
- Deng, B.-L.; Lepoivre, J. A.; Lemièrre, G. *Eur. J. Org. Chem.* **1999**, 2683-2688.
- Murata, M.; Watanabe, S.; Masuda, Y. *Synlett* **2000**, 1043-1045.
- For the similar cinnamyltin compounds, see: (a) Richter, F.; Weichmann, H. *J. Organometal. Chem.* **1994**, *466*, 77-87. (b) Fouquet, E.; Pereyre, M.; Rodriguez, A. L.; Roulet, T. *Bull. Soc. Chim. Fr.* **1997**, *134*, 959-967.
- (a) Burnett, D. A.; Caplen, M. A.; Domalski, M. S.; Browne, M. E.; Davis, H. R., Jr.; Clader, J. W. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 311-314. (b) Hervet, M.; Thery, I.; Gueffier, A.; Enguehard-Gueffier, C. *Helv. Chim. Acta* **2003**, *86*, 3461-3469.
- For the dimerization product **8a**, see: Li, J.; Qian, W.; Zhang, Y. *Tetrahedron* **2004**, *60*, 5793-5798.
- (a) Thibonnet, J.; Vu, V. A.; Berillon, L.; Knochel, P. *Tetrahedron* **2002**, *58*, 4787-4799. (b) Thibonnet, J.; Knochel, P. *Tetrahedron Lett.* **2000**, *41*, 3319-3322.